



---

## **Corrigendum to O3 and Drought Effects on Steady State Conductance and Kinetics in Pima Cotton**

Source: Air, Soil and Water Research, 15(1)

Published By: SAGE Publishing

URL: <https://doi.org/10.1177/11786221221135705>

---

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>)

# O<sub>3</sub> and Drought Effects on Steady State Conductance and Kinetics in Pima Cotton

David A Grantz<sup>1</sup> and Nancy E Grulke<sup>2</sup> 

<sup>1</sup>Department of Botany and Plant Sciences, University of California, Riverside, Parlier, CA, USA.

<sup>2</sup>Pacific Northwest Research Station, U.S.D.A. Forest Service, Bend, OR, USA.

Air, Soil and Water Research  
Volume 15: 1–10  
© The Author(s) 2022  
Article reuse guidelines:  
sagepub.com/journals-permissions  
DOI: 10.1177/11786221221114313



**ABSTRACT:** The degree to which ozone (O<sub>3</sub>) exposure and drought affect stomatal control of water loss and respond to environmental stimuli such as varying light is poorly characterized. To that end, we exposed Pima cotton to chronic O<sub>3</sub> exposure (month-long daytime exposures) with and without sufficient water, as well as short term acute O<sub>3</sub> exposure and varying light levels to understand stomatal kinetics. Chronic, month-long exposure to moderately high O<sub>3</sub> (~114 ppb) reduced daytime steady state stomatal conductance (g<sub>s</sub>), as did water deficit. Both stomatal opening and closing displayed dose specific, “sluggish” responses to step-changes in illumination with acute, 1-day, O<sub>3</sub> exposures of 0, 50, 100, and 125 ppb. At higher concentration (150 ppb), stomatal control of both opening and closing was degraded. Altered steady state and dynamic stomatal function suggest that elevated ambient O<sub>3</sub>, expected to increase in the future, may increasingly influence field water management and appropriate crop choices.

**KEYWORDS:** Ozone effects, drought stress, stomatal kinetics, crop water use

**RECEIVED:** November 16, 2021. **ACCEPTED:** June 26, 2022.

**TYPE:** Original Research

**CORRESPONDING AUTHOR:** Nancy E Grulke, Pacific Northwest Research Station, U.S.D.A. Forest Service, 63095 Deschutes Market Road, Bend, OR 97701, USA.  
Email: nancy.grulke@usda.gov

## Introduction

Although it has been broadly documented that elevated ozone (O<sub>3</sub>) exposure and uptake decreases plant productivity, alters within-plant resource allocation, and has economic repercussions in agronomic species (Tai & Val Martin, 2017), its effect on plant water balance is under-investigated. The frequency, severity, and global distribution of elevated ambient O<sub>3</sub>, water limitation, and their intersection over agronomic regions are predicted to increase due to increases in tropospheric air pollutants and ongoing changes in regional climate (Bates et al., 2008; Emberson et al., 2018; ONS, 2014; Sitch et al., 2007; The Royal Society, 2008). Stomatal conductance (g<sub>s</sub>) is sensitive to both O<sub>3</sub> exposure and water limitation and is the principal physiological regulator of carbon and water balance from plant to landscape scales. Understanding the roles of these stressors in plant water balance, particularly in crops, is important for planning and prediction of societal water consumption (Jackson et al., 2001). Further, stomatal conductance alone inadvertently regulates uptake of O<sub>3</sub> and other gaseous pollutants into the leaf, forming biologically important positive and negative feed-back loops.

Chronic and acute O<sub>3</sub> exposure (referring to sustained vs. brief duration) exert three distinct impacts on stomatal regulation: altered daytime rates of water loss (Nussbaum et al., 1995; Saurer et al., 1991; Vozzo et al., 1995), altered responsiveness to changes in other environmental stimuli (Leipner et al., 2001; McAinsh et al., 2002), and altered night time rates of water loss due to incomplete closure at night after daytime exposure (Grams et al., 1999; Grantz & Yang, 1996; Grulke et al., 2007b, 2007c; Hoshika, Omasa et al., 2013; Kellomaki & Wang, 1997; Kitao et al., 2009; Oksanen, 2003; Paoletti, 2005). All three regulatory processes have the capacity to impact total plant

water use (Booker et al., 2004) and regional hydrology (McLaughlin, Nosal et al., 2007; McLaughlin, Wullschlegel et al., 2007; Sellers et al., 1996; Sun et al., 2012).

Environmental effects on opening and closing kinetics appear to be closely related to each other (Paoletti & Grulke, 2010). Sluggish and or incomplete stomatal closure uncouples transpiration from carbon assimilation, degrades water use efficiency (WUE; Kirschbaum et al., 1988; Lawson & Blatt, 2014; Lawson et al., 2011), and depletes soil moisture (*reviewed in* Paoletti & Grulke, 2005). O<sub>3</sub>-induced, sluggish stomatal response was suspected (Keller & Häslner, 1984; Reich & Lassoie, 1984; Skarby et al., 1987) and later confirmed in other gas exchange studies. Responses of g<sub>s</sub> to many environmental stimuli were slowed by O<sub>3</sub> exposure including PPFD (Hoshika et al., 2012; Paoletti & Grulke, 2010; Tjoelker et al., 1995), VPD (Grulke et al., 2007a, 2007c; Hoshika, Omasa et al., 2013; Hoshika, Watanabe et al., 2013; Matyssek et al., 1995; Tjoelker et al., 1995; Uddling et al., 2009; Wieser & Havranek, 1995), soil moisture (Hayes et al., 2012), and CO<sub>2</sub> (Onandia et al., 2011).

The reduction of daytime g<sub>s</sub> by water limitation has long been known (Hsiao, 1973). As evaluated within a meta-analysis (Wittig et al., 2007), g<sub>s</sub> is also reduced by low to moderately high daytime O<sub>3</sub> exposure. Species with large midday g<sub>s</sub> may limit plant water loss by rapid closing responses, reducing the potential for xylem cavitation (Caird et al., 2007; Drake et al., 2013; Grantz et al., 2019; Violet-Chabrand et al., 2013; Vico et al., 2011). Both stressors provide proximate protection of the mesophyll from O<sub>3</sub> uptake at times of both peak O<sub>3</sub> concentration and evaporative demand in cotton (Temple, 1986, 1990;) and in a number of other species (Cavender-Bares et al., 2007; Grulke et al., 2003; Massman et al., 2000; Panek, 2004; Paoletti



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial

& Grulke, 2005; Temple et al., 1988). Static and kinetic stomatal responses to  $O_3$  and water limitation have been investigated separately, but their interactive effects on stomatal kinetics are not well characterized (Fuhrer & Booker, 2003; Hoshika, Omasa et al., 2013; Matyssek et al., 2006; Nikolova et al., 2010). Both antagonistic (ie, protective: Silim et al., 2009; Temple, 1986; Temple et al., 1988; Temple, 1990) and synergistic (ie, deleterious; Heggestad et al., 1985; Wagg et al., 2012, 2013) interactions of  $O_3$  and water deficit have been described in some plant species. The net effect of  $O_3 \times$  water limitation on stomatal response on the three distinct impacts on stomatal regulation (daytime, nighttime, and during changes in other environmental conditions) considered above remains unexplored.

Incomplete stomatal closure increases nighttime transpiration and uptake of  $O_3$ . Downwind of urban centers,  $O_3$  remains elevated at night (Gregg et al., 2003; Matyssek et al., 1995; Miller et al., 1972) and has been shown to inhibit growth of cottonwood, birch, and ponderosa pine (respectively). It was also suspected in the collapse of viticulture downwind of Los Angeles in the late 1950s (P.R. Miller, personal communication).  $O_3$  is also elevated at night downwind from heavily fertilized agricultural fields due to  $NO$  emissions and  $NO_x$  interconversions to  $O_3$  in sunlight. These emission plumes are transported into other agricultural areas as well as to natural ecosystems (Almaraz et al., 2018; Matson et al., 2002; Miller et al., 1972) where their effects are often unrecognized. The effects of  $O_3$  on nighttime transpiration has been reported in a number of species (Grulke et al., 2004, 2007a; Hoshika et al., 2012; Matyssek et al., 1995; Wieser & Havranek, 1993;). In birch, nighttime  $g_s$  contributed about 10% of total transpiration in low  $O_3$  (Matyssek et al., 1995), and 15% of total transpiration in high  $O_3$ . In blue oak and black oak, chronic daytime  $O_3$  exposure increased nighttime  $g_s$  to about 16% and 30% (respectively) of daytime maxima (Grulke et al., 2007b).

The study presented here evaluates  $O_3$  impacts on the three aspects of stomatal regulation: altered daytime  $g_s$ , incomplete stomatal closure at night with chronic exposure, and sluggish stomatal response to stepped increase and decrease of illumination during acute exposure of previously un-exposed (“naïve”) plants. We evaluate a perennial species of economic importance, Pima cotton (*Gossypium barbadense* L.; cv. S-6), that has been characterized with respect to responses to  $O_3$  and water limitation separately (Grantz, 2016; Temple & Grantz, 2010).

## Materials and Methods

### Plant material

Seeds of Pima cotton (*Gossypium barbadense* L.) from foundation seed stock were germinated in moist commercial potting mix (Earthgro Potting Soil, Scotts Company, Marysville, OH<sup>1</sup>) in plastic pots (870 ml; 110 mm  $\times$  110 mm  $\times$  125 mm). Plants were grown as described previously (Grantz et al., 2015) in a greenhouse at the University of California, Kearney Agricultural Center (Parlier, CA, USA; 103 masl; 36.598°N, 119.503°W).

Pots were thinned to a single plant 10 to 12 days after planting (DAP). Plants remained on the greenhouse bench in filtered air until they developed five to six leaves (May 15 and June 21, 2014, in runs 1 and 2, respectively). Single-plant pots were then available for two types of experiments: stomatal kinetics of cotton with (1) chronic  $O_3$  exposure at two water availabilities in exposure chambers; and (2) bench-top experiments with naïve (previously unexposed to  $O_3$ ) plants, using stepped high and stepped low light level to drive kinetics in different background  $O_3$  concentrations.

### Chronic $O_3$ exposure $\times$ water availability: steady state conductance

Eighteen pots were distributed among nine cylindrical, Teflon-walled,  $O_3$  exposure chambers (Continuously Stirred Tank Reactors; CSTRs; Grantz et al., 2008; Heck et al., 1978) located in the same greenhouse. The CSTRs were aligned in three blocks parallel to windows with cooling fans to reduce location effects. One CSTR per block was exposed to each of the three  $O_3$  concentration profiles delivered as daily half-sine waves with peak concentrations at 1,300 PDT; 12 hours daytime, mean  $O_3$  concentrations were 4, 59, and 114 ppb.  $O_3$  was generated by corona discharge (Model SGC-11, Pacific Ozone Technology, Brentwood, CA) from purified oxygen (Series ATF-15, Model 1242, SeQual Technologies Inc., San Diego, CA). The flow rate through the CSTRs was one air exchange  $\text{min}^{-1}$ ; air temperature was 20°C to 35°C, relative humidity was 34% to 60%, and midday PPFD from sunlight was 1,275  $\text{mmol m}^{-2} \text{s}^{-1}$  (Grantz et al., 2008; Paudel et al., 2016).

Plants in the CSTRs were grown under well-watered (WW) or water-deficit (WD) conditions. Field capacity of the soil was 21% by volume (VWC). Plants were irrigated to maintain a VWC of 15% to 18% (WW, ~80% of pot “field” capacity), and 9% to 11% (WD, ~50% of pot field capacity). All pots were fertilized twice a week with a complete fertilizer solution (Miracle Gro, Scotts Miracle-Gro Products Inc., Port Washington, NY, USA). The two runs of the experiment yielded consistent results and were pooled. The data for chronic  $O_3$  exposures in CSTRs were analyzed as a split plot, randomized complete block design.  $O_3$  was the main treatment, water application rate was the sub-treatment, and CSTR was the unit of replication. Values of  $g_s$  were log-transformed prior to analysis using PROC GLM (SAS for Windows; v. 9.2.1).

Stomatal conductance ( $g_s$ ) was measured on the youngest, fully expanded leaf, on 3 to 4 dates per experimental run, between 30 and 45 DAP ( $n=125$ ). Diurnal and nighttime  $g_s$  measurements were made, between 0730 and 1800 at 90 minutes intervals, and at 0200, respectively, within the CSTRs. Based on previous research, nighttime gas exchange minimas are achieved ~2 hours after full darkness is achieved (Grulke et al., 2004, 2007b). The intent of the nighttime  $g_s$  measurements was to capture the impact of daytime  $O_3$  exposure on  $g_s$ .

at night. Measurements of  $g_s$  were conducted with a cycling leaf porometer (AP-4; Delta-T Devices, Cambridge, UK).

#### *Acute O<sub>3</sub> exposure of naïve cotton: Dynamic stomatal responses to illumination*

Our experiments were conducted with previously unexposed (ie, “O<sub>3</sub>-naïve”), WW potted plants in the laboratory, using a custom gas exchange system modified from a system previously described (Grulke & Paoletti, 2005; Grulke et al., 2007b, 2007c). To drive stomatal response kinetics within the context of differing concurrent O<sub>3</sub> concentrations (0, 50, 100, 125, or 150 ppb), stepped changes in PPFD were imposed from low to saturating light levels (100–1600 μmol m<sup>-2</sup> s<sup>-1</sup>) and the reverse, randomizing which light level was applied first. After a 1-hour equilibration at each O<sub>3</sub> concentration at the beginning of the day,  $g_s$  was allowed to come to steady state at the initial PPFD level, and illumination was increased or decreased depending on the initial light level. O<sub>3</sub> exposure was maintained as a constant throughout the experiment.

The gas exchange system consisted of matched, custom-designed, sample and reference cuvettes (i.d. 2 × 3 × 1 cm) constructed of acrylic and lined with teflon film, fitted with low vibration micro-fans to ensure air mixing. The cuvettes were in parallel, receiving the same gas stream through the same length and diameter of Teflon tubing. Air for this open (flow-through) system was drawn from two large buffer volumes placed in series. Part of the gas stream was ozonated using an adjustable ultraviolet lamp, then cooled by passing across electronically controlled Peltier blocks, then humidified to maintain leaf to air VPD at ~2 kPa using a dewpoint generator (LI-610; LiCor Inc., Lincoln, NE, USA). Leaves were illuminated from above with a red and blue LED light array (LI-6400-18; LiCor Inc.; Lincoln, NE, USA); PPFD was measured at leaf level within the cuvette. The rest of the plant was illuminated uniformly using a bank of LEDs with a similar wavelength intensity profile (EcoSmart ECS 38 V2; 300K, 24W; 120W tungsten bulb equivalent). A portion of the youngest fully expanded leaf was exposed to the experimental O<sub>3</sub> concentrations in the sample cuvette, one concentration per day. Individual leaves were measured only once and individual plants were not measured on consecutive days.

Water vapor, CO<sub>2</sub> and O<sub>3</sub> were recorded at 15 seconds intervals, using the sample and reference cells of a steady state gas exchange system (LI-6400) in parallel with two cross-calibrated ultraviolet O<sub>3</sub> monitors (Model 41C; Thermo Fisher Scientific Inc.; Waltham, MA, USA). Leaf temperature was determined with a contact thermocouple (Type E, 76 μm dia) appressed to the abaxial surface of the leaf and monitored directly by the LI-6400 cuvette. This temperature, ~30°C, was lagged by 30 seconds in calculations of VPD and  $g_s$  to be in sync with flow from cuvettes to analyzers.

Half times ( $t_{1/2}$ ) for stomatal response to stepped changes in PPFD were calculated by fitting single exponential equations to  $g_s$  data obtained at the 15 seconds intervals. For stomatal opening:

$$g_s(t) = a - (b) e^{-\lambda t} \quad (1a)$$

and for stomatal closing:

$$g_s(t) = a + (b) e^{-\lambda t} \quad (1b)$$

where a and b are fitted parameters related to the initial and final magnitude of  $g_s$ ;  $\lambda$  is the fitted time constant, and t is time after the stepped change.

The half time was calculated from:

$$t_{1/2} = \ln(2)/\lambda \quad (2)$$

#### *Statistical analyses*

The data for chronic O<sub>3</sub> exposures in CSTRs were analyzed as a split plot, randomized complete block design. O<sub>3</sub> was the main treatment, water application rate was the sub-treatment, and CSTR was the unit of replication. Statistical analyses and graph preparation were performed with SAS for Windows, v. 9.2.1, SigmaPlot 11.0, and Systat 14.0. Statistical tests and their significance are as described in Results. Except as noted, significance is reported at  $p < .05$ .

## **Results**


### *Daytime steady state conductance under chronic O<sub>3</sub> exposure*

In all treatments, low O<sub>3</sub> (4 ppb, LO<sub>3</sub>) and moderately high O<sub>3</sub> (114 ppb, HO<sub>3</sub>), with adequate water (80% of “field” capacity in pots, WW) and limited water (50% of capacity, WD), peak daytime  $g_s$  values occurred at ~1,030 (Figure 1); end-of-day  $g_s$  was low. The peak level of  $g_s$  differed, with O<sub>3</sub> level, water regime, and their interaction significant ( $p = .010$ ;  $.002$ ; and  $.037$ , respectively). In the WW LO<sub>3</sub> treatment, diurnal  $g_s$  was characterized by a midday plateau (1030–1400). For plants in the three other treatments, the midday high was a peak diurnal value and the plateau as observed for WW LO<sub>3</sub> was truncated. WD reduced peak midday  $g_s$  by ~25% relative to WW LO<sub>3</sub> plants. High mean O<sub>3</sub> exposure (HO<sub>3</sub>, 114 ppb) reduced the midday peak  $g_s$  by ~55%, relative to WW LO<sub>3</sub>, whether plants were WW or WD.

### *Nighttime steady state conductance under chronic O<sub>3</sub> exposure*

Nighttime  $g_s$  was non-zero following daytime exposure to O<sub>3</sub> (Table 1). The effect of daytime LO<sub>3</sub> on nighttime  $g_s$  was minimal in both WW and WD treatments (both 1.9% of peak daytime  $g_s$ ). Daytime moderately high O<sub>3</sub> exposure increased the subsequent nighttime  $g_s$  to 7.7% and 7.3% of peak daytime values in WW and WD treatments, respectively. Assuming that  $g_s$  at 0600 and 1800 were those as measured at 0200 (nighttime  $g_s$ , Table 1), mean daytime and nighttime  $g_s$  were calculated. In this case, nighttime  $g_s$  was 16% to 17% of daytime  $g_s$  in high O<sub>3</sub>





**Figure 1.** The effect of chronic ozone ( $O_3$ ) exposure on the diel time course of stomatal conductance ( $g_s$ ) in Pima cotton under (a) well-watered (WW, ~80% of capacity) or (b) water deficit (WD, ~50% of capacity) conditions. Plants were exposed to low  $O_3$  (circles,  $LO_3$ ; 4 ppb, 12 hours mean) or moderately high  $O_3$  (triangles,  $HO_3$ ; 114 ppb, 12 hours mean) for 30 to 35 days (first and second run, respectively and combined here).

**Table 1.** Effect of Ozone Exposure Level ( $LO_3$ ,  $HO_3$ ) and Water Availability (WW; WD as Described in Figure 1) on Daytime Peak and Nighttime Steady State Stomatal Conductance ( $g_s$ ).

MEAN $O_3$	% "FIELD CAPACITY"	DAYTIME 1,000	NIGHTTIME 0200	DAYTIME WATER USE	NIGHTTIME WATER USE	TOTAL WATER USE
PPB		$G_s$ , MOL M <sup>-2</sup> S <sup>-1</sup>	$G_s$ , MOL M <sup>-2</sup> S <sup>-1</sup>	L/YEAR	L/YEAR	L/YEAR
4	80	$0.590 \pm 0.100$	$0.011 \pm 0.001$	59,494	1,130	60,624
4	50	$0.430 \pm 0.040$	$0.008 \pm 0.001$	38,074	990	39,064
114	80	$0.260 \pm 0.035$	$0.020 \pm 0.004$	21,001	924	21,925
114	50	$0.260 \pm 0.015$	$0.019 \pm 0.004$	19,277	578	19,855

Note. DAYTIME:  $O_3$ ,  $p = .010$ ; W,  $p = .002$ ;  $O_3 \times W$ ,  $p = .037$ . NIGHTTIME:  $O_3$ ,  $p = .26$ ; W,  $p = .057$ ;  $O_3 \times W$ ,  $p = .037$ . Acronyms as in Figure 1. Means  $\pm$  1 SD.

(WW and WD, respectively), and only 3.4% to 3.9% in low  $O_3$  (WW and WD, respectively).  $O_3$  ( $p = .025$ ) and water availability ( $p = .057$ ) individually had a significant effect on nighttime  $g_s$ , but the interaction term not significant ( $O_3 \times W$ ,  $p = .535$ ).


#### Stomatal response kinetics to acute $O_3$ exposure

The kinetics of stomatal opening (Figure 2a and c) and closing (Figure 2b and d) were evaluated following step increases (100–1600  $\mu\text{mol m}^{-2}\text{s}^{-1}$ ) and step decreases (1600–100  $\mu\text{mol m}^{-2}\text{s}^{-1}$ ) in PPFD during five levels of concurrent, acute (day of experiment)  $O_3$  exposure, with cotton plants not previously exposed to  $O_3$ . Time courses of both stomatal opening (equation (1a)) and closing (equation (1b)) were well described by exponential relationships over two levels of  $O_3$  exposure tested (50 and 100 ppb; Figure 2).


The mean  $t_{1/2}$  of stomatal opening and closing in  $O_3$ -free air did not significantly differ between replicate runs. The  $t_{1/2}$  for both stomatal opening and closing increased linearly with  $O_3$  between 0 and 125 ppb (Figure 3). On average, the  $t_{1/2}$  of stomatal opening was slower (eg, greater  $t_{1/2}$  times), and thus

more sensitive to  $O_3$  exposure, than that for closing (Figure 3; significant differences in slope,  $p = .005$ ). At 125 ppb the  $t_{1/2}$  of stomatal opening increased 3-fold and of closing by 1.7-fold, relative to no  $O_3$ . The  $t_{1/2}$  for both stomatal opening and closing was lower at 150 ppb than that at 125 ppb (Figure 3), reversing the increasingly sluggish  $g_s$  response and suggesting loss of stomatal control (Figure 3). Relative to the  $t_{1/2}$  for stomatal opening at 125 ppb  $O_3$ ,  $t_{1/2}$  at 150 ppb decreased by ~35%, and for stomatal closure, decreased by 40% (Figure 3). The greater variability in  $g_s$  observed at 125 ppb (Figure 3) may reflect variability in individual plant response, with some having reached their threshold  $O_3$  concentration and some not. Closing remained faster than opening at all non-zero  $O_3$ , even above 125 ppb.


The amplitude of stomatal opening ( $\Delta g_s$  from  $g_{i0}$  to  $g_{stf}$ ) increased linearly from 0 to 125 ppb  $O_3$  ( $p < .001$ ; Figure 4, solid circles). An abrupt decrease in the amplitude of stomatal opening was observed when  $[O_3]$  increased from 125 to 150 ppb  $O_3$  (Figure 4). The amplitude for stomatal closure with a stepped decrease in light was greater at each  $O_3$  exposure than that for opening, but was less responsive to  $O_3$ , differing



**Figure 2.** Representative stomatal responses to a stepped increase in PPFD (100–1,600  $\mu\text{mol m}^{-2}\text{s}^{-1}$ ; (a and c)), or a step decrease in PPFD (1600–100  $\mu\text{mol m}^{-2}\text{s}^{-1}$ ; (b and d)) in Pima cotton during acute exposure (same day exposure of naive plants) to 50 ppb  $\text{O}_3$  (a and b) or 100 ppb  $\text{O}_3$  (c and d). Values of  $t_{1/2}$  and solid lines were determined by fitting single exponential functions for opening (equation (1a)) and closing (equation (1b)). Curves were fitted to the open circles, beginning immediately after the stepped change in PPFD and ending when the new steady state was attained. Filled circles represent  $g_s$  before and after the response.




**Figure 3.** Effect of acute ozone ( $\text{O}_3$ ) exposure in Pima cotton on the kinetics ( $t_{1/2}$ , half time response to PPFD stimulus or reduction) of stomatal opening (filled circles; solid line) and stomatal closing (open triangles; dashed line) (means  $\pm$  1 S.E.). Linear regressions were fitted for opening and closing separately for data between 0 and 125 ppb and was assumed to linearly decrease from 125 to 150 ppb  $\text{O}_3$  (dotted lines).



**Figure 4.** Effect of 1-day ozone ( $\text{O}_3$ ) exposure of naive Pima cotton on the amplitude ( $\Delta g_{\text{st}0}$  to  $g_{\text{st}t}$ ) of stomatal opening (circles; solid line) and stomatal closing (triangles; dashed line) to a stepped change in PPFD from 100 to 1600  $\mu\text{mol m}^{-2}\text{s}^{-1}$  (opening) or 1600 to 100  $\mu\text{mol m}^{-2}\text{s}^{-1}$  (closing) (means  $\pm$  1 S.E.). A linear regression was fitted for stomatal opening for exposure between 0 and 125 ppb; a linear trend in stomatal amplitude was assumed between 125 and 150 ppb.

significantly from  $\Delta g_s$  for opening only at 50 ppb (Student's  $t$  test; Figure 4, open triangles). The amplitude of closing did not change significantly from 50 to 150 ppb. Greater  $t_{1/2}$  was

correlated with a greater response amplitude, both for stomatal opening ( $p = .003$ ; Student's  $t$ -test) and closing ( $p = .029$ ; Mann-Whitney  $U$  test).



**Figure 5.** Mathematically fit instantaneous changes in stomatal conductance to increasing (left) and decreasing light (right) from experimentally derived mean initial, half-times ( $t_{1/2}$ ) and final  $g_s$ , in 0 ppb (solid lines) and 125 ppb  $O_3$  (dashed lines). Presented graphically here to 20 minutes. Transpirational water use calculated to 60 minutes presented in Table 2.

**Table 2.**  $O_3$  Impact on the Half Time of Stomatal Response to Increasing or Decreasing Light, Cumulative  $g_s$ , and Resulting Effect on Integrated  $g_s$ , Over Periods of 5 to 60 Minutes in Pima Cotton.

[ $O_3$ ]	0 PPB				125 PPB			
Time, min	5	10	20	60	5	10	20	60
Stomatal opening								
Half time, $t_{1/2}$	2.06				6.42			
Amplitude range, $\Delta g_s$	0.12 (0.13–0.25)				0.18 (0.10–0.28)			
Cumulative $g_s$ , $\Sigma g_s$	348	387	418	440	252	306	372	454
$\Sigma g_s$ , 125–0 ppb					–96	–81	–46	14
Stomatal closing								
Half time, $t_{1/2}$	2.36				4.43			
Amplitude range, $\Delta g_s$	0.30 (0.30–0.10)				0.30 (0.30–0.10)			
Cumulative $g_s$ , $\Sigma g_s$	372	297	242	200	432	363	290	218
$\Sigma g_s$ , 125–0 ppb					60	66	48	18

#### Effect of sluggish stomatal response on transpiration

Mean values of initial  $g_s$ ,  $t_{1/2}$ , and final  $g_s$  across  $O_3$  exposures of 0 to 125 ppb  $O_3$  were used to define responses of stomatal opening and closing to stepped increased and decreased light, respectively (Figure 5). The responses were integrated to calculate the effect of changes in light with no and moderately high  $O_3$  exposure on cumulative transpiration over an hour. The cumulative values, and sign of the difference between opening and closing  $g_s$  measured in 125 versus 0 ppb, were similar to the differences in nighttime  $g_s$  in the two  $O_3$  exposures:  $\sim 22 \text{ mmol m}^{-2} \text{ s}^{-1}$  and  $20 \pm 0.4 \text{ mmol m}^{-2} \text{ s}^{-1}$ , respectively

(Table 2). Up to 10 minutes following step change in PPFD, sluggish opening reduced cumulative  $g_s$  more than sluggish closure increased it. By 60 minutes, sluggish closure slightly increased cumulative  $g_s$  slightly more than did sluggish opening. As noted above,  $t_{1/2}$  was correlated with the amplitude of stomatal response. Disregarding this influence underestimated the effect of  $O_3$  on cumulative  $g_s$  and expected transpirational losses.

#### Discussion

Well-watered pima cotton grown with negligible  $O_3$  exposure had a bell-shaped, diurnal  $g_s$  curve with a midday plateau,

consistent with the pattern described in the analysis of the diel sensitivity in the same variety of cotton (Grantz et al., 2015). Grown in elevated  $O_3$  and or droughted, the maximum  $g_s$  occurred at 1030, and  $g_s$  then declined with incomplete stomatal closure by 1800. Moderately high  $O_3$  exposure reduced  $g_s$  and expected transpirational water loss to a greater extent than did drought stress, and in combination, the two did not act synergistically. Similar effects of  $O_3$  and WD were observed in the congeneric upland cotton, *G. hirsutum* (Temple, 1986, 1990; Temple et al., 1988). Although antagonistic (ie, protective) interactions (Silim et al., 2009; Temple, 1986, 1990; Temple et al., 1988) and synergistic (ie, deleterious) interactions (Heggstad et al., 1985; Wagg et al., 2012, 2013) have been described in this and other species (*Populus* spp.; *Glycine max* L. Merr., *Dactylis glomerata* L., *Ranunculus acris* L.).

Stomata did not completely close in any of the growth treatments tested ( $O_3$  level  $\times$  water availability) by the end of the day (1,800) or following several hours of darkness during the night (0200). Grown in  $LO_3$ , instantaneous nighttime  $g_s$  was <4% of daytime mean  $g_s$ ; in  $HO_3$ , instantaneous  $g_s$  was <18% of daytime values and slightly lower with concurrent water deficit. Nighttime  $g_s$  has rarely been measured in crop species exposed to elevated  $O_3$ .

Pima cotton has much higher  $g_s$  than is observed in many forest species (Lu et al., 1994) which could alter the relative impact of daytime and nighttime stomatal responses. In trees, nighttime  $g_s$  was ~15% of daytime maximum  $g_s$  values (beech (*Fagus crenata*), Hoshika, Watanabe et al. (2013); spruce (*Picea abies*), Wieser and Havranek (1993); larch (*Larix decidua*), Wieser and Havranek (1993, 1995) birch (*Betula pendula*), Matyssek et al. (1995); ponderosa pine (*Pinus ponderosa*), Grulke et al. (2004); blue oak (*Quercus douglasii*), Grulke et al., 2007b). After daytime high  $O_3$  exposure, nighttime  $g_s$  increased to 30% of daytime maxima in black oak (*Q. kelloggii*), Grulke et al., 2007b). Significant nighttime  $g_s$  has also been observed in the absence of  $O_3$  (Dawson et al., 2007) with drought or in areas of low soil nutrient availability (Caird et al., 2007). Other pollutants such as  $NO_x$  (Grulke et al., 2004) may also result in nighttime transpirational losses. Similar to responses of daytime  $g_s$ , nighttime  $g_s$  also responds to VPD,  $CO_2$ , and WD (Cavender-Bares et al., 2007; Daley & Phillips, 2006; Dawson et al., 2007; Donovan et al., 2003; Zeppel et al., 2011), potentially enhancing nutrient and oxygen availability (Caird et al., 2007) but reducing hydraulic lift and water redistribution between soil horizons (Dawson, 1996).

In comparison to those grown in WW  $LO_3$ , plants exposed to high  $O_3$  with either adequate water or a water deficit would lose less water despite the small increases due to sluggish closure and increased nocturnal  $g_s$  following  $O_3$  exposure. Although this suggests that exposure to  $HO_3$  would offer field water savings, both above- and below-ground biomass was significantly decreased in exposure levels comparable to the  $HO_3$  treatment described here (Paudel et al., 2016), suggesting that

$O_3$  toxicity is a greater determinant of growth than effects of water budgets. Decreased root biomass as reported would exacerbate the impact of soil water deficits (in cotton, Grantz et al., 2006; in ponderosa pine, Grulke & Balduman, 1999).

The lack of complete stomatal closure at the end of the day and at night suggested  $O_3$ -induced, mechanistic inhibition of stomatal kinetics (Caird et al., 2007; Paoletti & Grulke, 2010; Torsethaugen et al., 1999). The  $t_{1/2}$  linearly increased from 0 to 125 ppb for both opening and closing in short term daytime  $O_3$  exposures. At all exposures,  $t_{1/2}$  times were greater for opening than closing in this controlled experiment, for example, less water was transpired with sluggish stomatal opening than with sluggish stomatal closure at the same  $O_3$  exposure. Slower stomatal opening restricts  $CO_2$  uptake under potentially more favorable conditions such as in the morning when VPDs may be lower, limiting transpirational losses. Slower stomatal closure potentially increases transpirational water losses under unfavorable conditions such as in the afternoon when VPDs are higher, increasing the rate of transpirational losses (Panek, 2004; Patterson & Rundel, 1989). An increase in the time for stomatal response to stimulus at high  $O_3$  has been described as “sluggish” (Kaiser & Paoletti, 2014). The “sluggish” stomatal response with  $O_3$  exposure reported here is consistent with previous results (Handley & Grulke, 2008; Hetherington & Woodward, 2003; Paoletti & Grulke, 2010).

High  $O_3$  concentrations (150 ppb) decreased  $t_{1/2}$ , and if the change in  $t_{1/2}$  from 125 to 150 ppb can be assumed to be linear, the rate of decline for both opening and closing is similar, suggesting that either antioxidant capacities were exceeded at that concentration (Wieser & Matyssek, 2007), and or that guard or subsidiary cell turgor was altered due to changes in membrane ion exchange (Dumont et al., 2013; Moldau et al., 1990). These data provide evidence of a fundamental change in stomatal regulation (or loss thereof) of a 1 day, high  $O_3$  exposure in cotton. A similar threshold was observed in experimental exposures of deciduous oak (*Q. kelloggii*, Grulke & Paoletti, 2005), European beech (*Fagus sylvatica*, Paoletti et al., 2020), and hybrid poplar (*Populus*, spp., Grantz, unpubl. data). In each case, an apparent loss of stomatal control occurs above a species-specific  $O_3$  concentration.

Greater  $t_{1/2}$  was correlated with a greater amplitude in stomatal response to a step change in PPFD as  $O_3$  increased for stomatal opening, but the  $t_{1/2}$  for stomatal closure was not related to the amplitude of response. These kinetic parameters can be sensitive to plant stress history (Assmann & Grantz, 1990; Lawson & Blatt, 2014; Percy & Way, 2012) and an interaction at the level of signaling has been suggested (Wilkinson & Davies, 2010), in which stress ethylene induced by  $O_3$  antagonizes responses to abscisic acid (ABA). ABA is known to induce stomatal closure under conditions of soil or root water deficit. However, in this cultivar of Pima cotton ethylene emission was not induced by  $O_3$  over this range of exposure (Grantz & Vu, 2012). Disregarding the amplitude of the change in  $g_s$  to environmental stimuli will underestimate transpirational losses.



The decline in  $g_s$  after a mid-morning peak to high  $O_3$  and drought dominated the daytime integrated response so that effects of  $O_3$  or WD on steady state stomatal responses do not provide a mechanism for increased stand water loss of Pima cotton. This contrasts with previous results in a deciduous forest (eg, McLaughlin, Nosal et al., 2007; McLaughlin, Wullschlegel et al., 2007;). Our calculations were based on a well-characterized daytime course of  $g_s$  and representative nighttime  $g_s$ , both obtained under controlled conditions. Extrapolation to canopy-scale water loss or  $O_3$  uptake requires more complete characterization of the diel course of  $g_s$ , environmental conditions, and plant species composition. Steady state conditions are not common outside of greenhouses or the laboratory, so that the impacts on steady state conductance do not reflect stomatal kinetics in variable environmental conditions (eg, Kaiser & Paoletti, 2014). Our estimates of cotton water use were estimated with “full sun” leaves which dominate gas exchange. Similarly, shade leaves of poplar also had altered stomatal kinetics but the steady state and kinetic  $g_s$  of full sun leaves dominated whole tree water balance (Paoletti et al., 2020). Response rate has been shown to be negatively related to stomatal size (Drake et al., 2013), and the stomata of Pima cotton are larger than those of many tree species.

## Conclusions

The effects of  $O_3$  exposure on fundamental processes of plant water balance were investigated in Pima cotton, an economically important species grown in the Central Valley of California, U.S.A. We report here that low to moderately elevated  $O_3$  exposure (up to 125 ppb): (1) reduced daytime and nighttime steady state  $g_s$  responses; (2) increased sluggish stomatal opening with increasing light; (3) increased sluggish stomatal closure with decreased light; and (3) incomplete nighttime closure. As the magnitude of stomatal conductance is much larger in the daytime, and daytime stomatal closure was faster than that of opening, the net result of  $O_3$  exposure was reduction of transpirational losses in Pima cotton up to 125 ppb. The current results suggest that neither nighttime stomatal opening nor sluggish stomatal closure are likely to increase whole plant, or agricultural field water use. Our results are consistent with those obtained with soybean under open-air  $O_3$  exposure in field conditions, in which a consistent decrease in canopy water loss was observed with increasing  $O_3$ . However, acute  $O_3$  exposures in the range of 125 to 150 ppb affected two aspects of stomatal kinetics: a decrease in half times for both stomatal opening and closing, and a decrease in amplitude of stomatal opening, suggesting a loss of stomatal control. If ambient or hourly spikes in  $O_3$  concentrations were to increase, the water balance of agricultural fields would need to be further considered.

## Author Note

This manuscript evaluates the effect of realistic tropospheric ozone exposure and drought stress on stomatal control of an important crop, Pima cotton, and how it may influence field water management.

## Acknowledgements

The authors are grateful for the skilled technical assistance of V.-B. Vu and R. Paudel during the performance of the gas exchange experiments. The authors declare no competing financial interests.

## Author Contributions

Conceived and designed the experiments: DAG, NEG. Analyzed the data: DAG. Wrote the first draft of the manuscript: DAG. Contributed to the writing of the manuscript: DAG, NEG. Agree with the manuscript results and conclusions: DAG, NEG. Jointly developed the structure and arguments for the paper: DAG, NEG. Made critical revisions and approved final version: DAG, NEG. All authors reviewed and approved of the final manuscript.

## Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

## Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This research was supported by the University of California at Riverside Research Allocation Process, the U.S. Department of Agriculture, Forest Service through Cooperative Agreement Order AG-04T0-P-13-0052.

## Disclosures and Ethics

Author(s) have provided to the publisher signed confirmation of compliance with legal and ethical obligations including but not limited to the following: authorship and contributorship, conflicts of interest, privacy and confidentiality. The authors have read and confirmed their agreement with the ICMJE authorship and conflict of interest criteria. This article is unique and not under consideration or published in any other publication.

## ORCID iD

Nancy E. Grulke  <https://orcid.org/0000-0002-6317-5073>

## NOTE

1. Mention of commercial names are for convenience only and do not constitute promotion of product.

## REFERENCES

- Almaraz, M., Bai, E., Wang, C., Trousdell, J., Conley, S., Faloona, I., & Houlton, B. Z. (2018). Agriculture is a major source of  $NO_x$  pollution in California. *Science Advances*, 4(1), 3aao3477.
- Assmann, S. M., & Grantz, D. A. (1990). Stomatal response to humidity in sugarcane and soybean; effect of vapour pressure difference on the kinetics of the blue light response. *Plant Cell & Environment*, 13, 163–169.
- Bates, B. C., Kundzewicz, Z. W., Wu, S., & Palutik, J. P. (2008). *Climate Change and water. Technical Paper, Intergovernmental Panel on Climate Change* (pp. 13–31). IPCC Secretariat.



- McAinsh, M. R., Evans, N. H., Montgomery, L. T., & North, K. A. (2002). Calcium signalling in stomatal responses to pollutants. *New Phytologist*, 153, 441–447.
- McLaughlin, S. B., Nosal, M., Wullschlegler, S. D., & Sun, G. (2007). Interactive effects of ozone and climate on tree growth and water use in a southern Appalachian forest in the USA. *New Phytologist*, 174, 109–124.
- McLaughlin, S. B., Wullschlegler, S. D., Sun, G., & Nosal, M. (2007). Interactive effects of ozone and climate on water use, soil moisture content, and streamflow in a southern Appalachian forest in the USA. *New Phytologist*, 174, 125–136.
- Miller, P. R., McCutchan, M. H., & Milligan, H. P. (1972). Oxidant air pollution in the Central Valley, Sierra Nevada foothills, and Mineral King Valley of California. *Atmospheric Environment*, 6(9), 623–633.
- Moldau, H., Sober, J., & Sober, A. (1990). Differential sensitivity of stomata and mesophyll to sudden exposure of bean shoots to ozone. *Photosynthesis*, 24, 446–458.
- Nikolova, P. S., Andersen, C. P., Blaschke, H., Matyssek, R., & Häberle, K. H. (2010). Belowground effects of enhanced tropospheric ozone and drought in a beech/spruce forest (*Fagus sylvatica* L./*Picea abies* [L.] Karst). *Environmental Pollution*, 158, 1071–1078.
- Nussbaum, S., Geissmann, M., & Fuhrer, J. (1995). Effects of nitric oxide and ozone on spring wheat (*Triticum aestivum*). *Water Air & Soil Pollution*, 85, 1449–1454.
- Oksanen, E. (2003). Responses of selected birch (*Betula pendula* Roth) clones to ozone change over time. *Plant Cell & Environment*, 26, 875–886.
- Onandia, G., Olsson, A.-K., Barth, S., King, J. S., & Uddling, J. (2011). Exposure to moderate concentrations of tropospheric ozone impairs tree stomatal response to carbon dioxide. *Environmental Pollution*, 159, 2350–2354.
- Panek, J. A. (2004). Ozone uptake, water loss and carbon exchange dynamics in annually drought-stressed *Pinus ponderosa* forests: Measured trends and parameters for uptake modeling. *Tree Physiology*, 24, 277–290.
- Paoletti, E. (2005). Ozone slows stomatal response to light and leaf wounding in a Mediterranean evergreen broadleaf (*Arbutus unedo*). *Environmental Pollution*, 134, 439–445.
- Paoletti, E., & Grulke, N. E. (2005). Does living in elevated CO<sub>2</sub> ameliorate tree response to ozone? A review on stomatal responses. *Environmental Pollution*, 137, 483–493.
- Paoletti, E., & Grulke, N. E. (2010). Ozone exposure and stomatal sluggishness in different plant physiognomic classes. *Environmental Pollution*, 158, 2664–2671.
- Paoletti, E., Grulke, N. E., & Matyssek, R. (2020). Ozone amplifies short-term water loss from mature trees by slowing stomatal dynamics and decreases long-term water loss because of lower steady-state stomatal conductance. *Forests*, 11(1), 1–15.
- Patterson, M. T., & Rundel, P. W. (1989). Seasonal physiological responses of ozone stressed Jeffrey pine in Sequoia National Park, California. In R. K. Olson & A. S. Lefohn (Eds.), *Effects of air pollution on Western forests* (pp. 419–428). Air Pollution Control Association. Transaction Series.
- Paudel, R., Grantz, D. A., Vu, H. B., & Shrestha, A. (2016). Tolerance of elevated ozone and water stress in a California population of Palmer amaranth (*Amaranthus palmeri*). *Weed Science*, 64, 276–284.
- Pearcy, R. W., & Way, D. A. (2012). Two decades of sunfleck research: looking back to move forward. *Tree Physiology*, 32, 1059–1061.
- Reich, P. B., & Lassoie, J. P. (1984). Effects of low-level O<sub>3</sub> exposure on leaf diffusive conductance and water-use efficiency in hybrid poplar. *Plant Cell & Environment*, 7, 661–668.
- Saurer, M., Führer, J., & Siegenthaler, U. (1991). Influence of ozone on the stable carbon isotope composition, <sup>13</sup>C, of leaves and grain of spring wheat (*Triticum aestivum* L.). *Plant Physiology*, 97, 313–316.
- Sellers, P. J., Bounoua, L., Collatz, G. J., Randall, D. A., Dazlich, D. A., Los, S. O., Berry, J. A., Fung, I., Tucker, C. J., Field, C. B., & Jensen, T. G. (1996). Comparison of radiative and physiological effects of doubled atmospheric CO<sub>2</sub> on climate. *Science*, 271, 1402–1406.
- Silim, S., Nash, R., Reynard, D., White, B., & Schroeder, W. (2009). Leaf gas exchange and water potential responses to drought in nine poplar (*Populus* spp.) clones with contrasting drought tolerance. *Trees*, 23, 959–969.
- Sitch, S., Cox, P. M., Collins, W. J., & Huntingford, C. (2007). Indirect radiative forcing of climate change through ozone effects on the land-carbon sink. *Nature*, 448, 791–794.
- Skarby, L., Troeng, E., & Bostrom, C. A. (1987). Ozone uptake and effects on transpiration, net photosynthesis, and dark respiration in Scots pine. *Forest Science*, 33, 801–808.
- Sun, G., McLaughlin, S. B., Porter, J. H., Uddling, J., Mulholland, P. J., Adams, M. B., & Pederson, N. (2012). Interactive influences of ozone and climate on streamflow of forested watersheds. *Global Change Biology*, 18, 3395–3409.
- Tai, A. P. K., & Val Martin, M. (2017). Impacts of ozone air pollution and temperature extremes on crop yields: Spatial variability, adaptation and implications for future food security. *Atmospheric Environment*, 169, 11–21.
- Temple, P. J. (1986). Stomatal conductance and transpirational responses of field-grown cotton to ozone. *Plant Cell & Environment*, 9, 315–321.
- Temple, P. J. (1990). Water relations of differentially irrigated cotton exposed to ozone. *Agriculture Journal*, 82, 800–805.
- Temple, P. J., & Grantz, D. A. (2010). Air pollution stress. In J. M. Stewart, D. M. Oosterhuis, J. J. Heithold, & J. R. Mauney (Eds.), *Physiology of Cotton* (pp. 162–173). Springer.
- Temple, P. J., Kupper, R. S., Lennox, R. W., & Rohr, K. (1988). Injury and yield responses of differentially irrigated cotton to ozone. *Agriculture Journal*, 80, 751–755.
- The Royal Society. (2008). *Ground-level ozone in the 21st century: Future trends, impacts and policy implications*. Royal Society Policy. Retrieved August 19, 2012, from [http://royalsociety.org/uploadedFiles/Royal\\_Society\\_Content/policy/publications/2008/7925.pdf](http://royalsociety.org/uploadedFiles/Royal_Society_Content/policy/publications/2008/7925.pdf).
- Tjoelker, M. G., Volin, J. C., Oleksyn, J., & Reich, P. B. (1995). Interaction of ozone pollution and light effects on photosynthesis in a forest canopy experiment. *Plant Cell & Environment*, 18, 895–905.
- Torsethagen, G., Pell, E. J., & Assmann, S. M. (1999). Ozone inhibits guard cell K<sup>+</sup> channels implicated in stomatal opening. *Proceedings of the National Academy of Sciences*, 96, 13577–13582.
- Uddling, J., Teclaw, R. M., Pregitzer, K. S., & Ellsworth, D. S. (2009). Leaf and canopy conductance in aspen and aspen-birch forests under free-air enrichment of carbon dioxide and ozone. *Tree Physiology*, 29, 1367–1380.
- Violet-Chabrand, S., Dreyer, E., & Brendel, O. (2013). Performance of a new dynamic model for predicting diurnal time courses of stomatal conductance at the leaf level. *Plant Cell & Environment*, 36, 1529–1546.
- Vico, G., Manzoni, S., Palmroth, S., & Katul, G. (2011). Effects of stomatal delays on the economics of leaf gas exchange under intermittent light regimes. *New Phytologist*, 192, 640–652.
- Vozzo, S. F., Miller, J. E., Pursley, W. A., & Heagle, A. S. (1995). Atmospheric pollutants and trace gases – effects of ozone and water-deficit on field-grown soybean. 1. Leaf gas-exchange. *Journal of Environmental Quality*, 24, 663–670.
- Wagg, S., Mills, G., Hayes, F., Wilkinson, S., Cooper, D., & Davies, W. J. (2012). Reduced soil water availability did not protect two competing grassland species from the negative effects of increasing background ozone. *Environmental Pollution*, 165, 91–99.
- Wagg, S., Mills, G., Hayes, F., Wilkinson, S., & Davies, W. J. (2013). Stomata are less responsive to environmental stimuli in high background ozone in *Dactylis glomerata* and *Ranunculus acris*. *Environmental Pollution*, 175, 82–91.
- Wieser, G., & Havranek, W. (1993). Ozone uptake in the sun and shade crown of spruce: Quantifying the physiological effects of ozone exposure. *Trees*, 7, 227–232.
- Wieser, G., & Havranek, W. M. (1995). Environmental control of ozone uptake in *Larix decidua* Mill.: A comparison between different altitudes. *Tree Physiology*, 15, 253–258.
- Wieser, G., & Matyssek, R. (2007). Linking ozone uptake and defense towards a mechanistic risk assessment for forest trees. *New Phytologist*, 174, 7–9.
- Wilkinson, S., & Davies, W. J. (2010). Drought, ozone, ABA and ethylene: New insights from cell to plant to community. *Plant Cell & Environment*, 33, 510–525.
- Wittig, V. E., Ainsworth, E. A., & Long, S. P. (2007). To what extent do current and projected increases in surface ozone affect photosynthesis and stomatal conductance of trees? A meta-analytic review of the last 3 decades of experiments. *Plant Cell & Environment*, 30, 1150–1162.
- Zeppel, M. J., Lewis, J. D., Medlyn, B., Barton, C. V., Duursma, R. A., Eamus, D., Adams, M. A., Phillips, N., Ellsworth, D. S., Forster, M. A., & Tissue, D. T. (2011). Interactive effects of elevated CO<sub>2</sub> and drought on nocturnal water fluxes in *Eucalyptus saligna*. *Tree Physiology*, 31, 932–944.



# Corrigendum to O<sub>3</sub> and Drought Effects on Steady State Conductance and Kinetics in Pima Cotton

Air, Soil and Water Research  
Volume 15: 1  
© The Author(s) 2022  
Article reuse guidelines:  
sagepub.com/journals-permissions  
DOI: 10.1177/11786221221135705



Grantz, David A. and Grulke, Nancy E. (2022). O<sub>3</sub> and Drought Effects on Steady State Conductance and Kinetics in Pima Cotton. *Air, Soil and Water Research*. Advance online publication. <https://doi.org/10.1177/1178622122114313>

In the above-mentioned article, the authors missed including 'Rama Paudel' as the co-author. She was integral in the implementation of the research and the collection of the data.

Also, Grantz, David A. and Grulke, Nancy E. shared she is a second author for this article.

The author list has been updated to:

Dr. David A. Grantz, Rama Paudel, Dr. Nancy E. Grulke. Rama Paudel is affiliated with Department of Crop Science, University of Illinois at Urbana-Champaign, Assistant Professor, Institute of Agriculture and Animal Science, Tribhuvan University.

The following are the corrected text for Section paragraph, Acknowledgements, Author Contributions and Reference.

## Chronic O<sub>3</sub> exposure × water availability: steady state conductance

Eighteen pots were distributed among nine cylindrical, Teflon-walled, O<sub>3</sub> exposure chambers (Continuously Stirred Tank Reactors; CSTRs; Grantz et al., 2008; Heck et al., 1978) located in the same greenhouse. The CSTRs were aligned in three blocks parallel to windows with cooling fans to reduce location effects. One CSTR per block was exposed to each of the three O<sub>3</sub> concentration profiles delivered as daily half-sine waves with peak concentrations at 1,300 PDT; 12 hours daytime, mean O<sub>3</sub> concentrations were 4, 59, and 114 ppb. O<sub>3</sub> was generated by corona discharge (Model SGC-11, Pacific Ozone Technology, Brentwood, CA) from purified oxygen (Series ATF-15, Model 1242, SeQual Technologies Inc., San Diego, CA). The flow rate through the CSTRs was one air exchange min<sup>-1</sup>; air temperature was 20°C to 35°C, relative humidity was 34% to 60%, and midday PPFD from sunlight was 1,275 mmol m<sup>-2</sup> s<sup>-1</sup> (Grantz et al., 2008; Paudel et al., 2016; Paudel, 2015).

## Acknowledgements

The authors are grateful for the skilled technical assistance of H.-B. Vu during the performance of the gas exchange experiments.

## Author Contributions

Conceived and designed the experiments: DAG, NEG. Analyzed the data: DAG, RP. Wrote the first draft of the manuscript: DAG and RP. Contributed to the writing of the manuscript: DAG, RP and NEG. Agree with the manuscript results and conclusions: DAG, RP and NEG. Jointly developed the structure and arguments for the paper: DAG, NEG. Made critical revisions and approved final version: DAG, NEG. All authors reviewed and approved of the final manuscript.

## REFERENCE

Paudel, R. 2015. Effects of ozone and water deficit on stomatal kinetics, diel trends in stomatal conductance, growth and allometry in crops and weeds. California State University, Fresno, California. M.Sc. Thesis, pp. 19-35.

This has been corrected in the online version of the paper.



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without