

Species Removed from the Cicada Fauna of North America North of Mexico

Several species have been incorrectly attributed to the cicada fauna of North America north of Mexico. These species were generally identified in synonymic lists, and the error is perpetuated when new lists are constructed (e.g., Poole et al. 1997). Some of the records may have been made before subsequent species were described; and the incorrect, but available, description was applied to specimens at hand. We have listed the species that are incorrectly reported from North America north of Mexico and the actual distribution for the species based on the majority of localities given in Metcalf (1963a, b) and Duffels and van der Laan (1985). The multiple authors who identify these species as being distributed in North America north of Mexico are found in Metcalf (1963a, b) and Duffels and van der Laan (1985). We provide further discussion for species that require additional information to support this contention or for species that have been described from several regions of North America north of Mexico.

Family Cicadidae Latreille Subfamily Cicadinae Latreille Tribe Zammarini Distant

192. *Odopoea funesta* (Walker). The species is actually from Central and South America.

Zammaria funesta Walker 1858b: 2.

Odopoea funesta = *Zammaria funesta*, Distant 1906a: 97.

193. *Zammaria smaragdina* Walker. The species is found in Central and South America.

Zammaria smaragdina Walker 1850: 33.

Zammaria angulosa Walker 1850: 34.

Zammaria smaragdina = *Zammaria angulosa*, Metcalf 1963a: 116.