

## KEYS TO THE GENERA

## Males

- 1 Posterior margin of metanotum partially covered by mesonotum; timbal covers completely or partially covering the timbal organ ..... 2
- 1' Posterior margin of metanotum completely visible behind mesonotum; timbal covers absent ..... 10
- 2 (1) Timbal covers complete dorsally ..... 6
- 2' Timbal covers incomplete dorsally (Fig. 75, 79, 82, 84)..... 3
- 3 (2') Timbal covers triangular and concave medially; opercula meeting or near meeting (Fig. 79)..... *Pacarina* Distant
- 3' Timbal covers not triangular or concave medially; opercula separated (Fig. 74)..... 4
- 4 (3') 1<sup>st</sup> and 2<sup>nd</sup> crossveins of forewing parallel (Fig. 84) *Neocicada* Kato
- 4' 1<sup>st</sup> and 2<sup>nd</sup> crossveins of forewing not parallel ..... 5
- 5 (4) Lateral edges of pronotum smooth (Fig. 75)..... *Beameria* Davis
- 5' Lateral edges of pronotum jagged or irregularly toothed (Fig. 82)....  
..... *Quesada* Distant
- 6 (2) Pygofer with dorsal beak ..... 7
- 6' Pygofer without a dorsal beak (Fig. 9) ..... 9
- 7 (6) Width of head including eyes much narrower than the width of the mesonotum; length of the body less than two and one-half times the width of the mesonotum; abdomen broadly rounded caudally (Fig. 2) . *Cacama* Distant
- 7' Width of head including eyes about as wide or wider than the mesonotum; body length at least 2 1/2 times the width of the mesonotum; abdomen tapering caudally ..... 8
- 8 (7') Pygofer with a pair of spines protruding backward from the ventral edge (Fig. 73) ..... *Cornuplura* Davis
- 8' Pygofer lacking ventral spines (Fig. 34) ..... *Tibicen* Latreille
- 9 (6') Uncus simple; body length >35 mm (Fig. 34) ..... *Tibicen*
- 9' Uncus bifurcated; body length <35 mm (Fig. 9) ..... *Diceroprocta* Stål
- 10 (1') Uncus retractable ..... 11
- 10' Uncus exposed; sternite VIII elongated ..... 12
- 11 (10') Medial and cubital veins joined when emerging from basal cell of forewing (Fig. 96) ..... *Cicadetta* Kolenati
- 11' Medial and cubital veins emerging separately from basal cell of forewing (Fig. 88)..... *Magiccicada* Davis
- 12 (10') Timbal organs absent (Fig. 102)..... 13
- 12' Timbal organs present ..... 14
- 13 (12') Eight apical cells in forewing (Fig. 102) ..... *Platypedia* Uhler
- 13' Seven apical cells in forewing (Fig. 128)..... *Neoplatypedia* Davis
- 14 (12) Pronotum flared laterally forming longitudinal ridge or carina when viewed from the side; ocelli variously colored ..... 15
- 14' Pronotum constricted, rounding laterally; ocelli color ruby red (Fig. 201)..... *Okanagodes* Davis