

## Acknowledgements

I am indebted to the following botanical specialists for expert plant identifications: L. Anderson, J. H. Beaman, S. F. Blake, E. C. Leonard, T. Morley, G. B. Ownbey, K. F. Parker, S. G. Shetler, and L. M. Smith. I am also indebted to R. W. Hodges and the late J.F.G. Clarke for the initial *Gnorimoschema* identifications in this study. I thank the following entomological curators and specialists for *Gnorimoschema* specimen loans: B. Mather (BM), J.-F. Landry (CNC), C. C. Hsiung (LEM), J. W. Brown and R. W. Hodges (USNM), R. L. Brown (MSU), and E. C. Knudson (ECK). I also thank L. K. Miller for unflagging field assistance; L. Slabaugh, A. M. Fallon, and R. L. Talerico for collecting galls; R. A. Watkins and S. Passoa for searching the Ohio State University Museum of Biodiversity for Kellicott specimens and specimen records; R. W. Holzenthal for the use of photomicrographic equipment; R. D. Moon for advice on describing patchiness; J.-F. Landry for geographic informational assistance; M. Epstein for drawing my attention to, and forwarding information from, the archived Riley notebooks; R. O. Kendall for specimen data; and J. A. Powell and D. Povolny for generously providing a copy of their unpublished manuscript. I benefitted greatly from written comments by J. A. Powell and an unnamed reviewer on an early version of the manuscript, as well as from postreview discussions with J. A. Powell.