



Lectotypification of the name *Achillea brachyphylla* (Anthemideae, Asteraceae)

Author: Inceer, Huseyin

Source: *Candollea*, 79(1) : 171-173

Published By: The Conservatory and Botanical Garden of the City of Geneva (CJBG)

URL: <https://doi.org/10.15553/c2024v791a9>

Lectotypification of the name *Achillea brachyphylla* (Anthemideae, Asteraceae)

Huseyin Inceer

Abstract

INCEER, H. (2024). Lectotypification of the name *Achillea brachyphylla* (Anthemideae, Asteraceae). *Candollea* 79: 171–173. In English, English abstract. DOI: <http://dx.doi.org/10.15553/c2024v791a9>

Within the genus *Achillea* L. (Anthemideae, Asteraceae) from Turkey, the name *Achillea brachyphylla* Boiss. & Hausskn. was originally described by Edmond Boissier (1810–1885) in his *Flora Orientalis* based on two syntypes collected by Carl Haussknecht (1838–1903). A lectotype is designated here for this name. The specimen is deposited in Boissier's herbarium in Geneva (G-BOIS).

Keywords

ASTERACEAE – *Achillea* – Lectotype – Nomenclature

Address of the author:

Karadeniz Technical University, Faculty of Science, Department of Biology, 61080 Trabzon, Turkey. E-mail: inceer@ktu.edu.tr

Submitted on February 23, 2024 – Accepted on May 21, 2024 – First published online on June 28, 2024
ISSN: 0373-2967 (print), 2235-3658 (online) – Published by the Conservatoire et Jardin botaniques de Genève
Open access article under Creative Commons Attribution Licence (CC BY 4.0)

Introduction

The genus *Achillea* L. (*Anthemideae*, *Asteraceae*) includes c. 110–140 species, mostly distributed in Europe, Asia, and North Africa (EHRENDORFER & GUO, 2006; OBERPRIELER et al., 2007). Its centre of diversity is located in SE Europe and SW Asia, and the taxa are highly variable in morphology, life cycle, and ecology (GUO et al., 2004; BALTISBERGER & WIDMER, 2016). The genus *Achillea* encompasses c. 60 accepted taxa in Turkey, 33 of which are endemic (SEMİZ et al., 2022).

Achillea brachyphylla Boiss. & Hausskn. is a perennial herb with thick, woody rootstocks, primary root strongly developed, caespitose (ARABACI & YILDIZ, 2008). It is morphologically related to *A. membranacea* (Labill.) DC. and *A. oligocephala* DC. (HUBER-MORATH, 1975). *Achillea brachyphylla* differs from *A. membranacea* in having a shorter stem, tomentose involucre, ovate-lanceolate outer phyllaries apically acute, and obtuse-oblong paleae; it differs from *A. oligocephala* in having a larger involucre and longer ray florets (ARABACI & YILDIZ, 2008).

Achillea brachyphylla was described by Edmond Boissier (1810–1885) in his *Flora Orientalis* (BOISSIER, 1875: 273) based on two collections of Carl Haussknecht (1838–1903) from Turkey. Those collections are to be considered as syntypes. The taxonomic treatment of the genus *Achillea* in the *Flora of Turkey and the East Aegean Islands* did not designate a lectotype for this name (HUBER-MORATH, 1975). Hence, a lectotype for the name *A. brachyphylla* is designated here on the best-preserved material. Scan of the lectotype can be found in the CHG (2024) website.

Typification

Achillea brachyphylla Boiss & Hausskn. in Boiss., *Fl. Orientalis* 3: 273. 1875.

Type: “Hab. in calcareis Mesopotamiae prope Surug et Orfa et in monte Gebel Taktak (Haussk!). Fl. Mai”.

Lectotypus (designated here): **TURKEY. Prov. Şanlıurfa:** “inter Surug et Orfa”, V.1865, *Haussknecht s.n.* (G-BOIS [G00150025] image!; isolecto-: B [B100093148] image!, E [E00414097] image!, JE [JE00011029] image!, K [K000928408] image!, W [W0027047, W0027048] images!). **Syntypus:** **TURKEY. Prov. Şanlıurfa:** “Dhebel Taktak”, V.1867, *Haussknecht 536* (BM [BM000945997] image!, G-BOIS [G00150024] image!; G [G00412361] image!; JE [JE00011029] image!, K [K000928409] image!, W [W0027049] image!).

Distribution and ecology. – *Achillea brachyphylla* is endemic to Turkey and it has a restricted distribution in SE Anatolia. The species is an Irano-Turanian element and grows on calcareous rocky slopes and steppes (HUBER-MORATH, 1975; ARABACI & YILDIZ, 2008).

Notes. – In the protologue of *Achillea brachyphylla* BOISSIER (1875: 273) cited: “Hab. in calcareis Mesopotamiae prope Surug et Orfa et in monte Gebel Taktak (Haussk!). Fl. Mai”. Two Haussknecht collections are deposited in Boissier’s herbarium (G-BOIS). One specimen was collected in May 1865 between Suruç and Şanlıurfa (G-BOIS [G00150025]) and the other in May 1867 in the region of Tektek Dağları (G-BOIS [G00150024]).

Since G00150025 (*Haussknecht s.n.*) is taxonomically informative and better preserved than G00150024, it is designated here as the lectotype of *Achillea brachyphylla*. Several duplicates of both the lectotype and the remaining syntype have been located. A specimen in Herbarium Haussknecht in Jena (JE00011028) bears the collection number “578” and was collected by Haussknecht on limestones on 6 May 1865 between “Surug & Kifri”. This specimen is not considered as original material of *A. brachyphylla*.

The colour of the ray florets of *Achillea brachyphylla* was reported as ochroleucous (off-white colour or white tinted with yellow) by BOISSIER (1875), whereas HUBER-MORATH (1975) inaccurately described them as white. On the other hand, ARABACI & YILDIZ (2008) stated the colour as yellow, becoming ochroleucous at the time of fruiting; this information was based on field observations in the type locality of the species.

Acknowledgements

The author thanks Joel Calvo (G) and Martin Callmänder (G) for their thorough reviews of the original manuscript, which resulted in numerous improvements to this article.

References

- ARABACI, T. & B. YILDIZ (2008). Taxonomy and threatened categories of three *Achillea* L. (Asteraceae-Anthemideae) species previously cited in the Data Deficient (DD) category. *Turk. J. Bot.* 32: 311–317.
- BALTISBERGER, M. & A. WIDMER (2016). Chromosome numbers and karyotypes within the genus *Achillea* (Asteraceae: Anthemideae). *Willdenowia* 46: 121–135.
- BOISSIER, E. (1875). *Fl. Orientalis* 3.
- CHG [CATALOGUE DES HERBIERS DE GENÈVE] (2024). *Base de données des herbiers en ligne du Conservatoire et Jardin botaniques de Genève*. [<http://www.ville-ge.ch/musinfo/bd/cjb/chg>]
- EHRENDORFER, F. & Y.P. GUO (2006). Multidisciplinary studies on *Achillea* sensu lato (Compositae-Anthemideae): New data on systematics and phylogeography. *Willdenowia* 36: 1–19.
- GUO, Y.P., F. EHRENDORFER & R. SAMUEL (2004). Phylogeny and systematics of *Achillea* (Asteraceae-Anthemideae) inferred from nrITS and plastid *trnL-F* DNA sequences. *Taxon* 53: 657–672.

- HUBER-MORATH, A. (1975). *Achillea* L. In: DAVIS, P.H. (ed.), *Fl. Turkey* 5: 224–252.
- OBERPRIELER, C., R. VOGT & L.E. WATSON (2007). Tribe Anthemideae Cass. In: KADEREIT, J.W. & C. JEFFREY (ed.), *The families and genera of vascular plants* 8: 342–374. Springer-Verlag, Berlin.
- SEMİZ, G., T. UYSAL, M. BOZKURT & B. GUNAL (2022). A new *Achillea* (Asteraceae) species from southwestern Turkey. *Turk. J. Bot.* 46: 406–415.