



A new checklist of the mosses of the continental United States and Canada¹

Authors: Buck, William R., and Goffinet, Bernard

Source: *The Bryologist*, 127(4) : 484-549

Published By: The American Bryological and Lichenological Society

URL: <https://doi.org/10.1639/0007-2745-127.4.484>

A new checklist of the mosses of the continental United States and Canada¹

William R. Buck^{2,4} and Bernard Goffinet^{3,4}

² *Institute of Systematic Botany, The New York Botanical Garden, Bronx, NY 10458-5126, U.S.A.*; ³ *Department of Ecology and Evolutionary Biology, 75 North Eagleville Road, Unit 3043, University of Connecticut, Storrs, CT 06269-3043, U.S.A.*

ABSTRACT. The checklist includes a listing of the genera and species of North American Bryophyta thought to occur in the continental United States and Canada. The floras of Mexico, Hawaii and Greenland are not included. The current list recognizes 1565 species, 12 subspecies, 34 varieties and one form (for a total of 1612 taxa) in 366 genera and 100 families. As a preface to the list, a systematic arrangement of the families and included genera for North America is presented. Many changes from the previous checklist are documented via footnotes that provide references to where changes were made. Only synonymy since the previous checklist is included. Twenty nomenclatural changes are made. These include 19 new combinations: *Bryum brassicoides* (\equiv *Gemmabryum brassicoides*), *B. pacificum* (\equiv *Ptychostomum pacificum*), *B. torenii* (\equiv *Imbribryum torenii*), *B. vinosum* (\equiv *Gemmabryum vinosum*), *Chionoloma maragniphyllum* (\equiv *Oxystegus maragniphyllus*), *Lescurea tribulosa* (\equiv *Pseudoleskea tribulosa*), *Pterygoneurum* \times *kieneri* (\equiv *P. subsessile* var. *kieneri* Habeeb), *Pylaisiadelphina canadensis* (\equiv *Brotherella canadensis*), *Streblotrichum convolutum* var. *eustegium* (\equiv *Barbula eustegia*), *Streblotrichum convolutum* var. *gallinula* (\equiv *Barbula convoluta* var. *gallinula*), *Voitia angustata* (\equiv *Splachnum angustatum*), *V. mnioides* (\equiv *Splachnum mnioides*), *V. pallida* (\equiv *Tetraplodon pallidus*), *V. paradoxa* (\equiv *Splachnum paradoxum*), *V. urceolata* (\equiv *Splachnum urceolatum*), *Warnstorfia badia* (\equiv *Hypnum badium*), *W. straminea* (\equiv *Hypnum stramineum*), *W. straminea* var. *patens* (Lindb.) (\equiv *Amblystegium stramineum* var. *patens*), *W. wickesiae* (\equiv *Calliargon wickesiae*). A new order is also introduced: Rhizogemmales W.R.Buck & Goffinet (\equiv Rhizogemmaceae Bonfim Santos, Siebel & Fedosov).

KEYWORDS. Bryophyta, inventory, North America, systematics, classification.



We present here the newest version of the North American moss checklist. In doing so, we stand on the shoulders of previous North American bryologists. Beginning with A. J. Grout (1940), who published the first comprehensive checklist of North American mosses, the task was then taken over by Howard Crum with the collaboration of Lewis Anderson and Bill Steere (Anderson 1990; Anderson et al. 1990; Crum et al. 1965, 1973). It has now been

almost 35 years since the last checklist and our understanding of moss diversity, species delineation and phylogeny and its accompanying nomenclatural and classification changes have moved ahead faster than at any time in the past. Molecular data have revealed relationships previously undetected by morphological observations, and similarly may reveal that phenotypic divergences may be infraspecific rather than diagnostic of speciation events. Moss taxonomy is a dynamic field and hence changes are far from being finished and names will continue to change for the foreseeable future, as we can attest by the changes encountered during the two-year preparation of this list. Furthermore, some concepts remain contentious among authors (e.g., taxonomy

¹ We dedicate this checklist to the memory of Howard A. Crum and Lewis E. Anderson on whose shoulders we stand.

⁴ Corresponding authors' e-mails: bbuck@nybg.org; bernard.goffinet@uconn.edu

DOI: 10.1639/0007-2745-127.4.484

and systematics of the Pottiaceae), and consequently we have had to make choices. While we aimed at considering the evidence of alternative hypotheses (e.g., strength of the data, whether morphological or molecular, in supporting lineages), in various cases the final decision may have been shaped by our own general concepts. However, the impacts of our perhaps subjective decisions affect primarily systematic rather than taxonomic concepts at the species level, and hence should have a minimal impact on the recognized diversity of species and infraspecific taxa in the continental United States and Canada.

Moss systematics historically has been driven by phenotypic divergences, which is obviously the first step in an effort to organize diversity. The difficulty arose from the weight attributed to specific characters to diagnose lineages. A good example is the genus *Physcomitrella* Bruch & Schimp. (Funariaceae). Distinguished from *Physcomitrium* (Brid.) Brid. by its indehiscent, immersed capsules, its circumscription varied in terms of species and rank (Tan 1979) and even its affinities to *Aphanorhegma* Sull., a unispecific genus also characterized by immersed but operculate capsules. Inferences from DNA sequence data, first from about ten loci (Liu et al. 2012) and then hundreds of organellar (Medina et al. 2018) and then nuclear loci (Medina et al. 2019a), revealed that immersed sporangia evolved multiple times within *Physcomitrium*, and hence recognizing *Physcomitrella* or *Aphanorhegma* resulted in a paraphyletic *Physcomitrium*. Consequently, all species with immersed capsules were transferred to *Physcomitrium* (Medina et al. 2019a).

When a genus diagnosed by morphological traits is resolved as paraphyletic, based on DNA data, for example, due to the inclusion of a small genus, a single genus should be recognized if monophyly is a criterion, such as in the above case of *Physcomitrella* and *Physcomitrium*. The single genus recognized should bear the earliest name available (e.g., *Physcomitrium*) as dictated by the *International Code of Nomenclature for algae, fungi, and plants* (Turland et al. 2018). If the older name applies to a lesser-known genus, such priority can be overruled by a decision to conserve the younger name. Such is the recent case of *Plagiomnium* T.J.Kop. (Koponen 1968) being conserved over *Orthomnion* Wilson (Wilson 1857) at the recent congress in Madrid

(Wilson 2024) on the basis that the former is “a well-established name for a widely distributed genus and therefore conservation would promote nomenclatural stability for the broad community of users of botanical names.” Whether to conserve a name is to some extent a subjective decision based on the perceived “usage” of the younger name. However, one can argue that the sooner a name is adopted the sooner it will gain in usage; the time ahead of us is much longer than the past two centuries of Linnean classification. The genus *Voitia* Hornsch. (1818) has traditionally accommodated two species, and phylogenetic inferences drawn from many nuclear loci (Lewis et al. 2014) reveal that their common ancestor arose from within *Tetraplodon* Bruch & Schimp. (1844), which was thus resolved as paraphyletic. *Tetraplodon* occurs primarily in the Northern Hemisphere with disjunctions in southern South America, and high elevations in Africa and New Guinea, whereas *Voitia* is confined to Laurasia, where it is largely sympatric with species of *Tetraplodon* (Marino 2014). Both genera belong to the Splachnaceae and have been studied in this context (see A. Koponen [1982, 1990] and Marino [1988, 1991a, 1991b] and Marino et al. [2009]). Both genera have been accepted since their protologues and their species have been included in all subsequent inventories. Merging the two genera translates morphologically into species with indehiscent capsules (diagnostic of *Voitia*) having arisen from *Tetraplodon*-like ancestor with dehiscent sporangia. Such a scenario has been accepted for *Voitia grandis* D.G.Long transferred to *Tayloria* (Goffinet & Shaw 2002) and is recurrent in other genera such as *Physcomitrium* (Medina et al. 2019a) or *Weissia* Hedw. (Reese & Lemmon 1965; Zander 1993).

Other genera, primarily speciose ones, have been broken up based on their polyphyletic resolution; examples include *Brachythecium* Schimp. (Brachytheciaceae; e.g., Ignatov & Huttunen 2002) and *Hypnum* Hedw. (Hypnaceae; Schlesak et al. 2018). The outcome may be the recognition of new unispecific genera (e.g., *Buckia* D.Ríos, M.T.Gallego & J.Guerra), which serve as placeholders pending completion of a more comprehensive taxonomic sampling of the speciose genera. We have reluctantly accepted all these segregates. In other cases, the splitting of genera reflects perhaps more subjective preferences, and

may not be supported by phylogenetic evidence. For example, *Racomitrium* Brid. (Grimmiaceae) has been divided into several genera (i.e., the resurrected *Bucklandiella* Roiv. and *Codriophorus* P.Beauv. [Ochyra et al. 2003] and the newly proposed *Dilutineuron* Bedn.-Ochyra, Sawicki, Ochyra, Szczecińska & Plášek [Bednarek-Ochyra et al. 2015], *Frisvollia* Sawicki, Szczecińska, Bedn.-Ochyra & Ochyra [Sawicki et al. 2015] and *Niphotrichum* (Bedn.-Ochyra) Bedn.-Ochyra & Ochyra [Ochyra et al. 2003]). Although these generic segregates may be monophyletic, they all arose from a unique shared common ancestor, i.e., *Racomitrium* s.l. is itself monophyletic (Larraín et al. 2013). Hence, phylogenetically the recognition of smaller genera is not justified. Furthermore, these newer segregate genera lack unambiguous diagnostic morphological traits, given that their circumscription was dependent on DNA versus morphological character analyses (e.g., Bednarek-Ochyra et al. 2015, Sawicki et al. 2015). Hence, we recommend treating these lineages within *Racomitrium* as subgenera as originally proposed by Bednarek-Ochyra (1995). A similar case pertains to the *Warnstorfia* Loeske complex (Calliergonaceae), which comprises four genera, each accommodating only a few species: *Loeskypnum* H.K.G.Paul (2 spp.), *Sarmentypnum* Tuom. & T.J.Kop. (6 spp.), *Straminergon* Hedenäs (1 sp.) and *Warnstorfia* Loeske (4 spp.; Hedenäs 1993). Together these genera compose a monophyletic group (Hedenäs et al. 2005), which we propose recognizing at the generic level.

Another case of segregation of a new genus that can be avoided if a broader concept of the core genus is followed, is *Anomodon* Hook. & Taylor. Ignatov et al. (2019a) divided *Anomodon* into several genera. They erected *Pseudanomodon* for *Anomodon attenuatus* (Hedw.) Huebener because it was only distantly related to the clade comprising the type of *Anomodon*, i.e., *A. viticulosus* (Hedw.) Hook. & Taylor. We support recognizing *Pseudanomodon*. They also established the genus *Anomodontopsis* for *Anomodon rugelii* (Müll.Hal.) Keissl. because the latter is resolved as sister to a clade composed of *Anomodon* s.str. and *Haplohymenium* Dozy & Molk. Given the paraphyly of *Anomodon*, their proposal could be justified on phylogenetic grounds. However, Granzow-de la Cerda (1997) had previously argued for the merger of *Haplohymenium* with *Anomodon*; such a concept would restore the monophyly of *Anomodon*

and hence make the segregation of *Anomodontopsis* unnecessary.

In its traditional circumscription, *Bryum* Hedw. is among the most speciose genera of mosses (Crosby et al. 1999). Some of its diversity was accommodated early on in distinct genera (e.g., *Ptychostomum* Hornsch. [Hornschuch 1824]; *Plagiobryum* Lindb. [Lindberg 1862]), but most have been transferred more recently into new generic segregates (i.e., *Gemabryum* J.R.Spence & H.P.Ramsay [Spence & Ramsay 2005]; *Imbribryum* N.Pedersen [Pedersen 2005; Pedersen & Hedenäs 2005]; *Plagiobryoides* J.R.Spence [Spence 2005]; *Rosulabryum* J.R.Spence [Spence 1996]). The most recent phylogenetic analyses based on DNA data led Holyoak & Pedersen (2007) to merge *Plagiobryum* with *Ptychostomum* as well as other phylogenetic inferences, whether based on DNA (Pedersen & Hedenäs 2003; Wang & Zhao 2009) or morphological traits (Pedersen et al. 2007), thus raising doubts on the monophyly of the *Bryum* segregates, as recently also discussed by Toren et al. (2024). Consequently, we recognize a single, albeit morphologically diverse, genus, *Bryum*, pending a comprehensive taxonomic and data sampling to resolve the ambiguities.

Finally, generic concepts may be based on the integration of criteria other than strict monophyly. Zander (2024), for example, integrated functional information in the analysis of complexity leading him to recognize monothetic and fractal genera. Admittedly, we do not yet fully understand the conceptual basis for this approach. Therefore, we decided to not recognize the one new generic name that Zander (2023) applied to a taxon present in our flora, i.e., *Neotrichostomum* for *Trichostomum crispulum* Bruch.

Overall, we aimed for consistency in applying criteria for recognizing genera and understand that our concepts depart from those proposed or adopted by others, such as contributors to the *Bryophyte Flora of North America* (Flora of North America Editorial Committee 2007, 2014). In all cases, we sought to provide the basis or explanation for our systematic decisions. Systematic concepts will continue to be vulnerable to change as investigations continue to resolve relationships, methodological approaches evolve, and theoretical principles underlying classifications are developed. However, the generic or systematic concepts adopted have no bearing on the

actual inventory of moss species diversity. Our list provides a significant update to the previous one by Anderson et al. (1990). This list recognizes an additional 183 species and infraspecific taxa, an additional 54 genera, and an additional 28 families. For all additions, deletions, or synonymies we provide documentation in the form of footnotes. We have not reprinted the synonymy of previous lists but are only including names in the list of synonyms that differ from previous lists. All authority names have been checked and updated as needed. The supra-generic classification is built on that proposed by Goffinet et al. (2009a,b), and integrates proposals by Liu et al. (2019), Bechteler et al. (2023), Li et al. (2024) and many others as referenced in the footnotes throughout the classification. The list of taxa occurring in the United States and Canada is available as a spreadsheet (**Supplementary Table S1**), along with an alphabetical list of families of North American mosses and their generic inclusions (**Supplementary Table S2**), an alphabetical list of families of North American mosses and their ordinal affiliations (**Supplementary Table S3**), and an alphabetical list of orders of North American mosses and their familial inclusions (**Supplementary Table S4**).

SYSTEMATIC ARRANGEMENT OF GENERA OF NORTH AMERICAN MOSSES

SUBDIVISION TAKAKIOPHYTINA M.Stech & W.Frey

CLASS TAKAKIOPSIDA M.Stech & W.Frey

TAKAKIALES M.Stech & W.Frey

Takakiaceae M.Stech & W.Frey

Takakia S. Hatt. & Inoue

SUBDIVISION SPHAGNOPHYTINA Doweld

CLASS SPHAGNOPSIDA Schimp.

SPHAGNALES Limpr.

Sphagnaceae Dumort.

Sphagnum L.

SUBDIVISION ANDREAEOPHYTINA Goffinet, W.R.Buck & A.J.Shaw

CLASS ANDREAEOPSIDA Trevis.

ANDREAEALES Limpr.

Andreaeaceae Dumort.

Andreaea Hedw.

CLASS ANDREAEOBRYOPSIDA Goffinet & W.R. Buck¹

ANDREAEOBRYALES B.M.Murray

Andreaeobryaceae Steere

Andreaeobryum Steere & B.M.Murray

SUBDIVISION BRYOPHYTINA Engl.

CLASS POLYTRICHOPSIDA Doweld

OEDIPODIALES Goffinet & W.R.Buck

Oedipodiaceae Schimp.

Oedipodium Schwägr.

POLYTRICHALES Cavers

Polytrichaceae Schwägr.

Atrichum P.Beauv., *nom. cons.*

Bartramiopsis Kindb.

Lyellia R.Br.

Oligotrichum DC., *nom. cons.*

Pogonatum P.Beauv.

Polytrichastrum G.L.Sm.

Polytrichum Hedw.

Psilopilum Brid.

TETRAPHIDALES Cavers

Tetraphidaceae Schimp.

Tetraphis Hedw.

Tetradontium Schwägr.

CLASS BRYOPSIDA Ritgen

SUBCLASS BUXBAUMIIDAE Doweld

BUXBAUMIALES Cavers

Buxbaumiaceae Schwägr.

Buxbaumia Hedw.

SUBCLASS DIPHYSCIIDAE Ochyra

DIPHYSCIALES M.Fleisch.

Diphysciaceae M.Fleisch.

Diphyscium D.Mohr

SUBCLASS GIGASPERMIDAE M.Stech & W.Frey

GIGASPERMALES Goffinet, Wickett, O.Werner, Ros,

A.J.Shaw & C.J.Cox

Gigaspermaceae A.Jaeger & Sauerb.

Lorentziella Müll.Hal. *ex* Besch.

¹ The Andreaeobryophytina were erected by Goffinet et al. (2009a), but recent phylogenetic evidence (Bechteler et al. 2024, Liu et al. 2019) supports that this lineage is sister to the Andreaeopsida, and this combined monophyletic lineage is best treated as the Andreaeophytina.

SUBCLASS FUNARIIDAE Ochyra

DISCELIALES Ignatov, Ignatova & Fedosov

Disceliaceae Schimp.*Discelium* Brid.

ENCALYPTALES Dixon

Encalyptaceae Schimp.*Bryobrittonia* R.S. Williams*Encalypta* Hedw.

FUNARIALES M.Fleisch.

Funariaceae Schwägr.*Entosthodon* Schwägr.*Funaria* Hedw.*Physcomitrium* (Brid.) Brid.*Pyramidula* Brid.

SUBCLASS TIMMIIDAE Ochyra

TIMMIALES Ochyra

Timmiaceae Schimp.*Timmia* Hedw., *nom. cons.*

SUBCLASS DICRANIDAE Doweld

DISTICHIALES D.Bell & Goffinet

Distichiaceae Schimp.*Distichium* Bruch & Schimp., *nom. cons.***Timmiellaceae Y.Inoue & H.Tsubota***Luisierella* Thér. & P.de la Varde²*Timmiella* (De Not.) Schimp.

PSEUDODITRICHIALES Ignatov & Fedosov

Pseudoditrichaceae Steere & Z.Iwats.*Pseudoditrichum* Steere & Z.Iwats.

FLEXITRICHIALES D.Bell & Goffinet

Flexitrichaceae Ignatov & Fedosov ex D.Bell & Goffinet*Flexitrichum* Ignatov & Fedosov

CATOSCOPIALES Ignatov & Ignatova

Catoscopiaceae Boulay ex Broth.*Catoscopium* Brid.

SCOULERIALES Goffinet & W.R.Buck

Hymenolomataceae Ignatov & Fedosov³*Hymenoloma* Dusén**Scouleriaceae S.P.Churchill***Scouleria* Hook.**Drummondiaceae Goffinet***Drummondia* Hook., *nom. cons.*

BRYOXIPHIALES Á.Löve & D.Löve

Bryoxiphiaceae Besch., nom. cons.*Bryoxiphium* Mitt., *nom. cons.*RHIZOGEMMALES W.R.Buck & Goffinet⁴**Rhizogemmaceae Bonfim Santos, Siebel & Fedosov***Rhizogemma* Bonfim Santos, Siebel & Fedosov

GRIMMIALES M.Fleisch.

Saelaniaceae Ignatov & Fedosov*Saelania* Lindb.**Grimmiaceae Arn.***Coscinodon* Spreng.*Dryptodon* Brid.*Grimmia* Hedw.*Jaffuelobryum* Thér.*Racomitrium* Brid.*Schistidium* Bruch & Schimp., *nom. cons.***Ptychomitriaceae Schimp.***Brachydontium* Fürnr.*Campylostelium* Bruch & Schimp.*Indusiella* Broth. & Müll.Hal.⁵*Ptychomitrium* Fürnr., *nom. cons.***Seligeriaceae Schimp.***Blindia* Bruch & Schimp.*Blindiadelphus* (Lindb.) Fedosov & Ignatov*Seligeria* Bruch. & Schimp.ARCHIDIALES Limpr.⁶**Archidiaceae Schimp.***Archidium* Brid.**Micromitriaceae Smyth ex Goffinet & Budke***Micromitrium* Austin**Leucobryaceae Schimp.***Atractylocarpus* Mitt., *nom. cons.*² See Inoue & Tsubota (2014).³ See Fedosov et al. (2016a).⁴ The Rhizogemmaceae were erected by Bonfim Santos et al. (in Fedosov et al. 2023b), and resolved as sister to a clade comprising Archidiales, Seligeriales, Grimmiales and Dicranales. We propose to accommodate this family in its own order, Rhizogemmales W.R.Buck & Goffinet comb. et stat. nov. ≡ Rhizogemmaceae Bonfim Santos, Siebel & Fedosov, *Plants* 12: 1360 [13]. 2023.⁵ See Hernández-Maqueda et al. (2008).⁶ See Bechteler et al. (2023).

Brothera Müll.Hal.
Campylopus Brid.
Leucobryum Hampe

AMPHIDIALES D.Bell & Goffinet

Amphidiaceae M.Stech

Amphidium Schimp., *nom. cons.*

DICRANALES M.Fleisch.

Ruficaulaceae Bonfim Santos & Fedosov

Ruficaulis Bonfim Santos & Fedosov

Dicranellopsidaceae Bonfim Santos, Siebel & Fedosov

Dicranellopsis Bonfim Santos, Siebel & Fedosov

Fissidentaceae Schimp.

Fissidens Hedw.

Schistostegaceae Schimp.

Schistostega D.Mohr

Dicranaceae Schimp., *nom. cons.*

Campylopodia Cardot

Dicranodontium Bruch & Schimp.

Dicranoweisia Lindb. *ex Milde*

Dicranum Hedw.

Oncophorus (Brid.) Brid.

Oreas Brid., *nom. cons.*

Paraleucobryum (Lindb. *ex* Limpr.) Péterfi

Aongstroemiaceae De Not.

Aongstroemia Bruch & Schimp., *nom. cons.*

Calcidicranella Bonfim Santos, Fedosov & Jan Kučera

Dichodontium Schimp.

Diobelonella Ochyra

Dicranellaceae M.Stech

Dicranella (Müll.Hal.) Schimp., *nom. cons.*

Eccremidium Wilson

Octoblepharaceae A.Eddy *ex* M.Menzel

Octoblepharum Hedw.

Calymperaceae Kindb.

Calymperes Sw.

Syrrhopodon Schwägr.

ERPODIALES Goffinet

Erpodiaceae Broth.

Erpodium (Brid.) Brid.

Solmsiella Müll.Hal.

Venturiella Müll.Hal.

BRUCHIALES Goffinet

Bruchiaceae Schimp.

Bruchia Schwägr.

Trematodon Michx.

RHABDOWEISIALES D.Bell & Goffinet

Rhabdoweisiaceae Limpr., *nom. cons.*

Arctoa Bruch & Schimp.

Brideliella Fedosov, M.Stech & Ignatov

Cnestrum I.Hagen

Cynodontium Bruch & Schimp., *nom. cons.*

Kiaeria I.Hagen

Rhabdoweisia Bruch & Schimp.

Symblepharis Mont.

Rhachithecaceae H.Rob.

Rhachithecium Broth. *ex* Le Jol.

DITRICHIALES D.Bell & Goffinet

Ditrichaceae Limpr., *nom. cons.*

Ceratodon Brid.

Cleistocarpidium Ochyra & Bedn.-Ochyra

Ditrichum Timm *ex* Hampe, *nom. cons.*

Pleuridium Rabenh., *nom. cons.*

Pseudephemerum (Lindb.) I.Hagen

Trichodon Schimp.

POTTIALES M.Fleisch.

Pottiaceae Hampe, *nom. cons.*

Acaulon Müll.Hal.

Aloina Kindb., *nom. cons.*

Anoetangium Schwägr., *nom. cons.*

Aschisma Lindb.

Barbula Hedw., *nom. cons.*

Bellibarbula P.C.Chen

Bryoerythrophyllum P.C.Chen

Chenia R.H.Zander

Chionoloma Dixon

Crossidium Jur., *nom. cons.*

Crumia W.B.Schofield

Didymodon Hedw.

Ephemerum Hampe, *nom. cons.*

Eucladium Bruch & Schimp.

Geheebia Schimp.

Gertrudiella Broth.

Globulinella Steere

Gymnostomiella M.Fleisch.

Gymnostomum Nees & Hornsch., *nom. cons.*

Gyroweisia Schimp., *nom. cons.*

Henediella Paris

Hilpertia R.H.Zander

Husnotiella Cardot

Hydrogonium (Müll.Hal.) A.Jaeger

Hymenostylium Brid.

Hyophila Brid., *nom. cons.*

Hyophiladelphus (Müll.Hal.) R.H.Zander
Leptodontium (Müll.Hal.) Lindb.
Merceyopsis Broth. & Dixon
Microbryum Schimp.
Molendoa Lindb.
Ozobryum G.L.Sm.Merr.
Plaubelia Brid.
Pleurochaete Lindb.
Pseudocrossidium R.S.Williams
Pterygoneurum Jur., *nom. cons.*
Rhexophyllum Herzog
Scopelophila (Mitt.) Lindb.
Splachnobryum Müll.Hal.
Stegonia Venturi
Streblotrichum P.Beauv.
Syntrichia Brid.
Tortella (Müll.Hal.) Limpr., *nom. cons.*
Tortula Hedw., *nom. cons.*
Trichostomopsis Cardot
Trichostomum Bruch, *nom. cons.*
Triquetrella Müll.Hal.
Tuerckheimia Broth.
Vinealobryum R.H.Zander
Weissia Hedw.
Zanderella J.A.Jiménez & M.J.Cano

SUBCLASS BRYIDAE Engl.

SUPERORDER **BRYANAE** Goffinet & W.R.Buck

SPLACHNALES Ochyra

Splachnaceae Grev. & Arn.

Aplodon R.Br.
Splachnum Hedw.
Tayloria Hook.
Voitia Hornsch.

Meesiaceae Schimp.

Amblyodon P.Beauv., *nom. cons.*
Leptobryum (Schimp.) Wilson⁷
Meesia Hedw., *nom. cons.*
Paludella Ehrh. *ex* Brid.

HEDWIGIALES Ochyra

Hedwigiaceae Schimp.

Braunia Bruch & Schimp., *nom. cons.*
Hedwigia P.Beauv., *nom. cons.*
Pseudobraunia (Lesq. & James) Broth.

BARTRAMIALES M.Menzel

Bartramiaceae Schwägr.

Anacolia Schimp., *nom. cons.*
Bartramia Hedw., *nom. cons.*
Conostomum Sw.
Philonotis Brid.
Plagiopus Brid.

BRYALES Limpr.

Bryaceae Rchb.

Anomobryum Schimp.
Brachymenium Schwägr.
Bryum Hedw.
Haplodontium Hampe
Plagiobryum Lindb.
Rhodobryum (Schimp.) Limpr., *nom. cons.*

Roellobryaceae Ochyra

Roellobryon Ochyra

Mniaceae Schwägr.

Cinclidium Sw.
Cyrtomnium Holmen
Leucolepis Lindb.
Mnium Hedw., *nom. cons.*
Plagiomnium T.J.Kop., *nom. cons.*
Pseudobryum (Kindb.) T.J.Kop.
Rhizomnium (Mitt. *ex* Broth.) T.J.Kop.
Trachycystis Lindb.

Mielichhoferiaceae Schimp.

Epipterygium Lindb.
Mielichhoferia Nees & Hornsch.
Pohlia Hedw.

ORTHOTRICHALES Dixon

Orthotrichaceae Arn.

Codonoblepharon Schwägr.
Grottiella Steere
Leratia Broth. & Paris
Lewinskya F.Lara, Garilleti & Goffinet
Macrocoma (Hornsch. *ex* Müll.Hal.) Grout
Macromitrium Brid.
Nyholmiella Holmen & E.Warncke
Orthotrichum Hedw.
Plenogemma Plášek, Sawicki & Ochyra
Pulviger Plášek, Sawicki & Ochyra
Schlotheimia Brid.
Ulota D.Mohr
Zygodon Hook. & Taylor

⁷ See Cox & Hedderson (1999) and Cox et al. (2000).

RHIZOGONIALES Goffinet & W.R.Buck

Rhizogoniaceae Broth.

Pyrrhobryum Mitt.

AULACOMNIALES N.E.Bell, A.E.Newton & D.Quandt

Aulacomniaceae Schimp.

Aulacomnium Schwägr., *nom. cons.*

ORTHODONTIALES N.E.Bell, A.E.Newton & D.Quandt

Orthodontiaceae Goffinet

Orthodontium Schwägr.

SUPERORDER **HYPNANAE** W.R.Buck, Goffinet & A.J.Shaw

HYPNODENDRALES N.E.Bell, A.E.Newton & D.Quandt

Racopilaceae Kindb.

Racopilum P.Beauv.

HYPOPTERYGIALES Goffinet

Hypopterygiaceae Mitt.

Hypopterygium Brid.

HOOKERIALES M.Fleisch.

Daltoniaceae Schimp.

Daltonia Hook. & Taylor, *nom. cons.*

Hookeriaceae Schimp.

Hookeria Sm., *nom. cons.*

Pilotrichaceae Kindb.

Callicostella (Müll.Hal.) Mitt., *nom. cons.*

Cyclodictyon Mitt.

Lepidopilum (Brid.) Brid., *nom. cons.*

Trachyxiphium W.R.Buck

HYPNALES W.R.Buck & Vitt

Rutenbergiaceae M.Fleisch.

Pseudocryphaea E.Britton *ex* Broth.

Fontinalaceae Schimp.

Brachelyma Schimp. *ex* Cardot

Dichelyma Myrin

Fontinalis Hedw.

Climaciaceae Kindb.

Climacium F.Weber & D.Mohr

Pleuroziopsis Kindb. *ex* E.Britton

Amblystegiaceae Kindb.

Amblystegium Schimp.

Anacamptodon Brid.

Campylium (Sull.) Spruce

Campylophyllopsis W.R.Buck

Conardia H.Rob.

Cratoneuron (Sull.) Spruce

Drepanium (Schimp.) Spruce

Drepanocladus (Müll.Hal.) G.Roth, *nom. cons.*

Hygroamblystegium Loeske, *nom. cons.*

Hygrohypnella Ignatov & Ignatova

Hygrohypnum Lindb.

Leptodictyum (Schimp.) Warnst.

Limbella (Müll.Hal.) Renauld & Cardot

Palustriella Ochyra

Platyhypnum Loeske

Platylomella A.L.Andrews

Pseudoamblystegium Vanderp. & Hedenäs

Pseudocampylium Vanderp. & Hedenäs

Sanionia Loeske

Tomentypnum Loeske

Calliergonaceae Vanderp., Hedenäs, C.J.Cox & A.J.Shaw

Calliergon (Sull.) Kindb.

Hamatocaulis Hedenäs

Scorpidium (Schimp.) Limpr.

Warnstorfia Loeske

Helodiaceae Ochyra

Helodium Warnst., *nom. cons.*

Leskeaceae Schimp.

Claopodium (Lesq. & James) Renauld & Cardot

Haplocladium (Müll.Hal.) Müll.Hal., *nom. cons.*

Lescuraea Schimp.

Leskea Hedw.

Lindbergia Kindb.

Pseudoleskeella Kindb.

Thuidiaceae Schimp.

Abietinella Müll.Hal.

Echinophyllum T.J.O'Brien

Herpetineuron (Müll.Hal.) Cardot

Pelekium Mitt., *nom. cons.*

Raiiella Reimers

Thuidium Schimp.

Stereophyllaceae W.R.Buck & Ireland

Entodontopsis Broth.

Pilosium (Müll.Hal.) M.Fleisch.

Stereophyllum Mitt.

Brachytheciaceae Schimp.

Brachytheciastrum Ignatov & Huttunen

Brachythecium Schimp.

Bryoandersonia H.Rob.

Cirriphyllum Grout

Clasmatodon Hook. & Wilson

Donrichardsia H.A.Crum & L.E.Anderson

Eurhynchiastrum Ignatov & Huttunen
Homalotheciella (Cardot) Broth.
Homalothecium Schimp.
Kindbergia Ochyra
Koponeniella Huttunen & Ignatov
Myuroclada Besch.
Oxyrrhynchium (Schimp.) Warnst., *nom. cons.*
Palamocladium Müll.Hal.
Pseudoscleropodium (Limpr.) M.Fleisch.
Rhynchostegium Schimp., *nom. cons.*
Sciuro-hypnum (Hampe) Hampe
Scleropodium Schimp.
Zelometeorium Manuel

Meteoriaceae Kindb.

Meteorium Dozy & Molk., *nom. cons.*
Neodicladiella (Nog.) W.R.Buck

Myriniaceae Schimp.

Myrinia Schimp., *nom. cons.*

Fabroniaceae Schimp.

Fabronia Raddi

Stereodontaceae Hedenäs, Schlesak & D.Quandt

Stereodon (Brid.) Brid.

Pylaisiaceae Schimp.

Aquilonium Hedenäs, Schlesak & D.Quandt
Buckia D.Ríos, M.T.Gallego & J.Guerra
Callicladium H.A.Crum
Calliergonella Loeske
Calliergonellopsis Jan Kučera & Ignatov
Pseudohygrohypnum Kanda
Pseudostereodon (Broth.) M.Fleisch.
Ptilium De Not.
Pylaisia Schimp., *nom. cons.*
Roaldia P.E.A.S.Câmara & Carv.-Silva

Heterocladellaceae Ignatov & Fedosov

Heterocladiella Ignatov & Fedosov

Hypnaceae Schimp.

Bryocrumia L.E.Anderson
Campylophyllum (Schimp.) M.Fleisch.
Chryso-hypnum Hampe
Dacryophyllum Ireland
Gollania Broth.
Hageniella Broth.
Homomallium (Schimp.) Loeske
Hypnum Hedw., *nom. cons.*
Vesicularia (Müll.Hal.) Müll.Hal., *nom. cons.*

Pterigynandraceae Schimp.

Heterocladium Schimp.
Iwatsukiella W.R.Buck & H.A.Crum
Pterigynandrum Hedw.

Hylocomiaceae M.Fleisch.

Hylocomiadelphus Ochyra & Stebel
Hylocomiastrum M.Fleisch. *ex* Broth.
Hylocomium Schimp., *nom. cons.*
Leptohymenium Schwägr.
Loeskeobryum M.Fleisch. *ex* Broth.
Pleurozium (Sull.) Mitt., *nom. cons.*
Rhytidiadelphus (Limpr.) Warnst.
Rhytidiopsis Broth.

Rhytidiaceae Broth.

Rhytidium (Sull.) Austin *ex* C.F.Parker

Plagiotheciaceae M.Fleisch.

Herzogiella Broth.
Isopterygiella Ignatov & Ignatova
Isopterygiopsis Z.Iwats.
Myurella Schimp.
Ortholimnobium Dixon
Orthothecium Schimp., *nom. cons.*
Plagiothecium Schimp.
Platydictya Berk.
Pseudotaxiphyllum Z.Iwats.
Rectithecium Hedenäs & Huttunen
Redfearnia J.T.Wynns

Entodontaceae Kindb., nom. cons.

Entodon Müll.Hal.

Jocheniaceae Jan Kučera & Ignatov⁸

Jochenia Hedenäs, Schlesak & D.Quandt

Pylaisiadelphaceae Goffinet & W.R.Buck

Isopterygium Mitt.
Platygyrium Schimp., *nom. cons.*
Pylaisiadelpha Cardot
Serpoleskea (Hampe *ex* Limpr.) Loeske
Taxithelium Spruce *ex* Mitt.
Trochophyllohypnum Jan Kučera & Ignatov
Wijkia H.A.Crum

Sematophyllaceae Broth., nom. cons.

Acroporium Mitt.
Brittonodoxa W.R.Buck, P.E.A.S.Câmara & Carv.-Silva
Donnellia Austin
Heterophyllum (Schimp.) Müll.Hal. *ex* Kindb.
Sematophyllum Mitt.

Cryphaeaceae Schimp.

Cryphaea F.Weber
Dendroalsia E.Britton
Schoenobryum Dozy & Molk.

⁸ See Kučera et al. (2019).

Leucodontaceae Schimp.*Antitrichia* Brid.*Leucodon* Schwägr.*Nogopterium* Crosby & W.R.Buck**Pterobryaceae Kindb.***Henicodinium* (Müll.Hal.) Kindb.*Jaegerina* Müll.Hal.*Pirella* Cardot**Neckeraceae Schimp.***Alleniella* S.Olsson, Enroth & D.Quandt*Bryolawtonia* D.H.Norris & Enroth*Dannorrisia* Enroth*Forsstroemia* Lindb.⁹*Homalia* Brid., *nom. cons.**Metaneckera* Steere*Neckera* Hedw., *nom. cons.**Neckeropsis* Reichardt*Neomacounia* Ireland*Pseudanomodon* (Limpr.) Ignatov & Federov*Thamnobryum* Nieuwl.**Leptodontaceae Schimp.***Leptodon* D.Mohr, *nom. cons.***Lembophyllaceae Broth.***Bestia* Broth.*Isothecium* Brid.*Pseudisothecium* Grout*Tripterocladium* (Müll.Hal.) A.Jaeger**Anomodontaceae Kindb.***Anomodon* Hook. & Taylor*Anomodontella* Ignatov & Fedosov*Schwetschkeopsis* Broth.**Taxiphyllaceae Ignatov¹⁰***Leptopterigynandrum* Müll.Hal.*Taxiphyllum* M.Fleisch.**Ctenidiaceae Qinghua Wang & Y.Jia¹¹***Ctenidium* (Schimp.) Mitt.**Miyabeaceae Enroth, S.Olsson, Buchb., Hedenäs,****Huttunen & D.Quandt***Homaliadelphus* Dixon & P.de la Varde**Theliaceae M.Fleisch.***Thelia* Sull.

ALPHABETICAL LIST OF TAXA

Abietinella Müll.Hal. (THUIDIACEAE)*abietina* (Hedw.) M.Fleisch.***Acaulon*** Müll.Hal. (POTTIACEAE)*mediterraneum* Limpr.¹²*muticum* (Hedw.) Müll.Hal.*rufescens* A.Jaeger¹³*schimperianum* (Sull.) Sull.*triquetrum* (Spruce) Müll.Hal.***Acroporium*** Mitt. (SEMATOPHYLLACEAE)*smallii* (R.S.Williams) H.A.Crum & L.E.Anderson***Alleniella*** S.Olsson, Enroth & D.Quandt¹⁴ (NECKERACEAE)*besseri* (Łobarz.) S.Olsson, Enroth & D.Quandt*complanata* (Hedw.) S.Olsson, Enroth & D.Quandt***Aloina*** Kindb., *nom. cons.* (POTTIACEAE)*ambigua* (Bruch & Schimp.) Limpr.*bifrons* (De Not.) Delgad.*brevirostris* (Hook. & Grev.) Kindb.*hamulus* (Müll.Hal.) Broth.*rigida* (Hedw.) Limpr.***Amblyodon*** P.Beauv., *nom. cons.* (MEESIACEAE)*dealbatus* (Hedw.) P.Beauv.***Amblystegium*** Schimp. (AMBLYSTEGIACEAE)*serpens* (Hedw.) Schimp.***Amphidium*** Schimp., *nom. cons.* (AMPHIDIACEAE)*californicum* (Hampe ex Müll.Hal.) Broth.*lapponicum* (Hedw.) Schimp.*mougeotii* (Bruch & Schimp.) Schimp.***Anacamptodon*** Brid. (AMBLYSTEGIACEAE)*splachnoides* (Brid.) Brid.***Anacolia*** Schimp., *nom. cons.* (BARTRAMIACEAE)*baueri* (Hampe) Paris¹⁵*laevisphaera* (Taylor) Flowers*menziesii* (Turner) Paris***Andreaea*** Hedw. (ANDREAEACEAE)*alpestris* (Thed.) Schimp.*blyttii* Schimp.*crassinervia* Bruch*heinemannii* Hampe & Müll.Hal.*megistospora* B.M.Murray var. *megistospora*var. *epapillosa* (B.M.Murray) H.A.Crum &

L.E.Anderson

mutabilis Hook.f. & Wilson⁹ See Enroth et al. (2022).¹⁰ See Ignatov et al. (2012).¹¹ See Li et al. (2024).¹² See GBIF.org (2024a).¹³ See McLaughlin (2024).¹⁴ See Olsson et al. (2011).¹⁵ See McLaughlin & Carter (2024).

- nivalis* Hook.
rothii F.Weber & D.Mohr subsp. *rothii*
 subsp. *falcata* (Schimp.) Lindb.
rupestris Hedw. var. *rupestris*
 var. *papillosa* (Lindb.) Podp.
schofieldiana B.M.Murray
sinuosa B.M.Murray
Andraeobryum Steere & B.M.Murray (ANDRAEOBRYACEAE)
macrosporum Steere & B.M.Murray
Anoetangium Schwägr., *nom. cons.* (POTTIACEAE)
aestivum (Hedw.) Spruce
sikkimense M.N.Aziz & Vohra¹⁶
stracheyanum Mitt.¹⁷
Anomobryum Schimp. (BRYACEAE)
concinatum (Spruce) A.Jaeger
julaceum (P.Gaertn., B.Mey. & Scherb.) Schimp.
Anomodon Hook. & Taylor¹⁸ (ANOMODONTACEAE)
minor (Hedw.) Lindb.
rugelii (Müll.Hal.) Keissl.
thraustus Müll.Hal.
tristis (Ces.) Sull.
viticulosus (Hedw.) Hook. & Taylor
Anomodontella Ignatov & Fedosov (ANOMODONTACEAE)¹⁹
longifolius (Schleich. ex Brid.) Ignatov & Fedosov²⁰
Antitrichia Brid. (LEUCODONTACEAE)
californica Sull. ex Lesq.
curtipendula (Hedw.) Brid.
gigantea (Sull. & Lesq.) Kindb.²¹
Aongstroemia Bruch & Schimp. (AONGSTROEMIIACEAE)
canadensis (Mitt.) Siebel & Fedosov
grevilleana (Brid.) Müll.Hal.
longipes (Sommerf.) Bruch & Schimp.
schreberiana (Hedw.) Bonfim Santos & Fedosov²²

¹⁶ See Zander (2017).

¹⁷ See Zander & Eckel (2007).

¹⁸ The generic circumscription of the Anomodontaceae was revised by Ignatov et al. (2019a) based on phylogenetic inferences from organellar loci. We accept the recognition of *Pseudanomodon* and *Anomodontella* but not the segregate unispecific genus *Anomodontopsis*, and maintain the merger of *Haplohymenium* with *Anomodon* (Granzow-de la Cerda 1997).

¹⁹ See Ignatov et al. (2019a).

²⁰ *Anomodon longifolius* and *A. thraustus* were reported as new to North America by Granzow-de la Cerda & Düll (2009).

²¹ See Hedenäs (2008a).

²² See Fedosov et al. (2023a).

- Aplodon** R.Br. (SPLACHNACEAE)
wormskioldii (Hornem.) R.Br.
Aquilonium Hedenäs, Schlesak & D.Quandt²³
 (PYLAISIACEAE)
adscendens (Lindb.) Hedenäs, Schlesak & D.Quandt
plicatulum (Lindb.) Hedenäs, Schlesak & D.Quandt
Archidium Brid. (ARCHIDIACEAE)
alternifolium (Dicks. ex Hedw.) Mitt.
crassicostatum D.R.Toren, Kellman & Shevock²⁴
donnellii Austin
hallii Austin
minus (Renauld & Cardot) Snider
ohioense Schimp.
tenerrimum Mitt.
Arctoa Bruch & Schimp. (RHABDOWEISIIACEAE)
anderssonii Wich.
blyttii (Bruch & Schimp.) Loeske
fulvella (Dicks.) Bruch & Schimp.
glacialis (Berggr.) Fedosov, Jan Kučera & M.Stech²⁵
hyperborea (Gunnerus ex Dicks.) Bruch & Schimp.
starkei (F.Weber & D.Mohr) Loeske
Aschisma Lindb. (POTTIACEAE)
kansanum A.L.Andrews
Atractylocarpus Mitt., *nom. cons.* (LEUCOBRYACEAE)
subporodictyon (Broth.) Bonfim Santos & M.Stech²⁶
Atrichum P.Beauv., *nom. cons.* (POLYTRICHACEAE)
altecristatum (Renauld & Cardot) Smyth & L.C.
 R.Smyth
angustatum (Brid.) Bruch & Schimp.
crispulum Schimp. ex Besch.²⁷
crispum (James) Sull.
cylindricum (Willd. ex F.Weber) G.L.Sm.
flavisetum Mitt.
selwynii Austin
tenellum (Röhl.) Bruch & Schimp.
undulatum (Hedw.) P.Beauv.
Aulacomnium Schwägr., *nom. cons.* (AULACOMNIACEAE)
acuminatum (Lindb. & Arnell) Kindb.
androgynum (Hedw.) Schwägr.
heterostichum (Hedw.) Bruch & Schimp.
palustre (Hedw.) Schwägr.
turgidum (Wahlenb.) Schwägr.

²³ *Aquilonium* was established by Schlesak et al. (2018) based on phylogenetic evidence.

²⁴ See Toren et al. (2016).

²⁵ See Fedosov et al. (2021).

²⁶ See Bonfim Santos & Stech (2017).

²⁷ See Merrill & Ireland (2007).

Barbula Hedw., *nom. cons.* (POTTIACEAE)
orizabensis Müll.Hal.
unguiculata Hedw.

Bartramia Hedw., *nom. cons.* (BARTRAMIACEAE)
aprica Müll.Hal.²⁸
brevifolia Brid.²⁹
halleriana Hedw.
ithyphylla Brid.
pomiformis Hedw.
potosica Mont.
subulata Bruch & Schimp.

Bartramiopsis Kindb. (POLYTRICHACEAE)
lescurii (James) Kindb.

Bellibarbula P.C.Chen³⁰ (POTTIACEAE)
recurva (Griff.) R.H.Zander

Bestia Broth. (LEMBOPHYLLACEAE)
longipes (Sull. & Lesq.) Broth.

Blindia Bruch & Schimp. (SELIGERACEAE)
acuta (Hedw.) Bruch & Schimp.

Blindiadelphus (Lindb.) Fedosov & Ignatov³¹
 (SELIGERACEAE)
campylopodus (Kindb.) Fedosov & Ignatov
diversifolius (Lindb.) Fedosov & Ignatov
polaris (Berggr.) Fedosov & Ignatov
recurvatus (Hedw.) Fedosov & Ignatov
subimmersus (Lindb.) Fedosov & Ignatov

Brachelyma Schimp. ex Cardot (FONTINALACEAE)
subulatum (P.Beauv.) Schimp. ex Cardot

Brachydontium Fűrnr. (PTYCHOMITRIACEAE)
olympicum (E.Britton) T.T.McIntosh & J.R.Spence
trichodes (F.Weber) Milde

Brachymenium Schwägr. (BRYACEAE)
andersonii H.A.Crum
exile (Dozy & Molk.) Bosch & Sande Lac.³²
erectum (Hook.) Wilson ex Mont.
macrocarpum Cardot
mexicanum Mont.
niveum Besch.
speciosum (Hook.f. & Wilson) Steere
systylium (Müll.Hal.) A.Jaeger
vinosulum Cardot³³

²⁸ See Müller (2014).²⁹ See Fransén (1995).³⁰ See Zander (1993).³¹ See Fedosov et al. (2017).³² See Spence et al. (2022).³³ See Spence (2007).

Brachytheciastrum Ignatov & Huttunen³⁴
 (BRACHYTHECIACEAE)
collinum (Schleicher ex Müll.Hal.) Ignatov & Huttunen
delicatulum (Flowers) Ignatov
fendleri (Sull.) Ochyra & Żarnowiec
leibergii (Grout) Ignatov & Huttunen
salicinum (Schimp.) Orgaz, M.J.Cano & J.Guerra³⁵
trachypodium (Brid.) Ignatov & Huttunen
velutinum (Hedw.) Ignatov & Huttunen var.
velutinum

Brachythecium Schimp. (BRACHYTHECIACEAE)
acuminatum (Hedw.) Austin
acutum (Mitt.) Sull.
albicans (Hedw.) Schimp.
asperrimum (Mitt. ex Müll.Hal.) Sull.³⁶
boreale Ignatov³⁷
brandegeei (Austin) H.Rob.
campestre (Müll.Hal.) Schimp.
capillaceum (F.Weber & D.Mohr) Giacom.³⁸
cirrosum (Schwägr.) Schimp.
erythrorrhizon Schimp. var. *erythrorrhizon*
 var. *alpinum* Kosovich-Anderson & Ignatov³⁹
falcatum (Grout) H.A.Crum⁴⁰
frigidum (Müll.Hal.) Besch.
jacuticum Ignatov⁴¹
laetum (Brid.) Schimp.
novae-angliae (Sull. & Lesq.) Austin subsp.
novae-angliae
 subsp. *hultenii* (E.B.Bartram) Huttunen
rivulare Schimp.
ruderale (Brid.) W.R.Buck⁴²
rutabulum (Hedw.) Schimp.
salebrosum (Hoffm. ex F.Weber & D.Mohr) Schimp.
tauriscorum Molendo & Lorentz⁴³
turgidum (Hartm.) Kindb.
udum I.Hagen⁴⁴

³⁴ See Ignatov & Huttunen (2002).³⁵ See Orgaz et al. (2013).³⁶ See Norris & Shevock (2004a,b).³⁷ See Ignatov (2014).³⁸ See Hodgetts et al. (2020).³⁹ See Kosovich-Anderson & Ignatov (2010).⁴⁰ See Crum (2001).⁴¹ See Ignatov (2014).⁴² See Buck (1998).⁴³ See Hedenäs (2017).⁴⁴ See Ignatov (2014).

- Braunia** Bruch & Schimp., *nom. cons.* (HEDWIGIACEAE)
andrieuxii Lorentz⁴⁵
secunda (Hook.) Bruch & Schimp.
- Brideliella** Fedosov, M.Stech & Ignatov⁴⁶
 (RHABDOWEISIACEAE)
wahlenbergii (Brid.) Fedosov, M.Stech & Ignatov
- Brittonodoxa** W.R.Buck, P.E.A.S.Câmara & Carv.-Silva⁴⁷ (SEMATOPHYLLACEAE)
subpinnata (Brid.) W.R.Buck, P.E.A.S.Câmara & Carv.-Silva
- Brothera** Müll.Hal. (DICRANACEAE)
leana (Sull.) Müll.Hal.
- Bruchia** Schwägr. (BRUCHIACEAE)
bolanderi Lesq.
brevifolia Sull.
carolinae Austin
drummondii Hampe ex E.Britton
flexuosa (Schwägr.) Müll.Hal.
fusca E.Britton
hallii Austin
queenslandica I.G.Stone
ravenelii Wilson ex Sull.
texana Austin
vogesiaca Nest. ex Schwägr.
- Bryoandersonia** H.Rob. (BRACHYTHECIACEAE)
illecebra (Hedw.) H.Rob.
- Bryobrittonia** R.S.Williams (ENCALYPTACEAE)
longipes (Mitt.) D.G.Horton
- Bryocrumia** L.E.Anderson (HYPNACEAE)
vivicolor (Broth. & Dixon) W.R.Buck
- Bryoerythrophyllum** P.C.Chen (POTTIACEAE)
columbianum (F.J.Herm. & E.Lawton) R.H.Zander
ferruginascens (Stirt.) Giacom.
inaequalifolium (Taylor) R.H.Zander
recurvirostrum (Hedw.) P.C.Chen
wallichii (Mitt.) P.C.Chen⁴⁸
- Bryolawtonia** D.H.Norris & Enroth⁴⁹ (NECKERACEAE)
vancouveriensis (Kindb.) D.H.Norris & Enroth
- Bryoxiphium** Mitt., *nom. cons.* (BRYOXIPHIACEAE)
norvegicum (Brid.) Mitt.
- Bryum** Hedw. (BRYACEAE)
aenum Blytt ex Bruch & Schimp.
algovicum Sendtn. ex Müll.Hal. var. *algovicum*⁵⁰

⁴⁵ See Allen (2010).

⁴⁶ See Fedosov et al. (2021).

⁴⁷ See Carvalho-Silva et al. (2017).

⁴⁸ See Allen (2022).

⁴⁹ See Norris & Enroth (1990).

- var. *rutheanum* (Warnst.) Crundw.
alpinum Huds. ex With.
amblyodon Müll.Hal.
andicola Hook.
apiculatum Schwägr.
archangelicum Bruch & Schimp.
arcticum (R.Br.) Bruch & Schimp.
argenteum Hedw. var. *argenteum*
 var. *majus* Schwägr.
 var. *muticum* Brid.
austriacum Köckinger, Holyoak & Suanjak⁵¹
biforme R.S.Williams⁵²
billarderii Schwägr.
bimum (Schreb.) Turner⁵³
blindii Bruch & Schimp.
bornholmense Wink. & R.Ruthe
brachyneuron Kindb.
brassicoides (J.R.Spence & Kellman) W.R.Buck
 & Goffinet⁵⁴
caespiticium Hedw.
californicum Sull.⁵⁵
calobryoides J.R.Spence
calophyllum R.Br.
canariense Brid.
capillare Hedw. var. *capillare*
 var. *barbatum* (C.E.O.Jensen) Podp.
cellulare Hook.⁵⁶
chryseum Mitt.⁵⁷
coronatum Schwägr.
creberrimum Taylor
cryophilum Mårtensson⁵⁸

⁵⁰ Spence (2014) treated *B. algovicum* as a synonym of *Ptychostomum pendulum*; in *Bryum* the correct name should be *B. algovicum* as the combination *B. pendulum* (Hornsch.) Schimp. has earlier homonyms.

⁵¹ See Spence et al. (2022).

⁵² See Spence et al. (2022).

⁵³ See Spence (2014) as *Ptychostomum bimum* (Schreb.) J.R.Spence.

⁵⁴ Reported as *Gemmabryum brassicoides* by Spence & Kellman (2015) but accommodated here in *Bryum*; *Bryum brassicoides* (J.R.Spence & Kellman) W.R.Buck & Goffinet comb. nov. ≡ *Gemmabryum brassicoides* J.R.Spence & Kellman, *Madroño* 62(2):124. 2015. TYPE: U.S.A., California, Santa Cruz Co.: Quail Hollow Ranch County Park, 37°04'57"N, 122°03'50"W, 18 Mar. 2006, K.M. Kellman 5104 (holotype, CAS).

⁵⁵ See Spence (2014).

⁵⁶ See Spence (2014).

⁵⁷ See Toren & Heise (2009).

curvatum Kaurin & Arnell
cyclophyllum (Schwägr.) Bruch & Schimp.
demaretianum Arts⁵⁹
dichotomum Hedw.
elegans Nees ex Brid.
eremaeum Catches. ex J.R.Spence & H.P.Ramsay⁶⁰
erythroloma (Kindb.) Syed
flabelliforme D.R.Toren, J.R.Spence & Shevock⁶¹
flaccidum Brid.
gemmiferum R.Wilczek & Demaret
gemmilucens R.Wilczek & Demaret
gemmiparum De Not.
hagenii Limpr.
imbricatulum Müll.Hal.⁶²
incrassatolimbatum Cardot
intermedium (Brid.) Turton
klinggraeffii Schimp.
knowltonii Barnes
kunzei Hornsch.⁶³
lanatum (P.Beauv.) Brid.⁶⁴
lonchocaulon Müll.Hal.
longisetum Blandow ex Schwägr.
marratii Hook. & Wilson
meesioides Kindb.
microchaeton Hampe
mildeanum Jur.⁶⁵
miniaturum Lesq.
moravicum Podp.
muehlenbeckii Bruch & Schimp.
neodamense Itzigs.⁶⁶
nitidulum Lindb.
oblongum Lindb.
pacificum (J.R.Spence & Shevock) W.R.Buck & Goffinet⁶⁷

pallens Sw.
pallescens Schleich. ex Schwägr.
pseudocapillare Besch.
pseudotriquetrum (Hedw.) P.Gaertn., B.Mey. & Scherb.
purpurascens (R.Br.) Bruch & Schimp.
radiculosum Brid.
reedii H.Rob.
riparium I.Hagen
rubens Mitt.
ruderales Crundw. & Nyholm
rutilans Brid. & Schimp.
salinum I.Hagen ex Limpr.
sanguilentum Renauld & Cardot⁶⁸
schleicheri Schwägr.
subapiculatum Hampe
subneodamense Kindb.
tenuisetum Limpr.
teres Lindb.
torenia (J.R.Spence & Shevock) W.R.Buck & Goffinet⁶⁹
torquescens Bruch & Schimp.⁷⁰
turbinatum (Hedw.) Turner
uliginosum (Brid.) Bruch & Schimp.
valparaisense Thér.
veronense De Not.⁷¹
vinosum (J.R.Spence & Kellman) W.R.Buck & Goffinet⁷²

J.R.Spence & Shevock, *Madroño* 59(3): 156. 2012. TYPE: U.S.A. California, Fresno Co.: Sierra National Forst, Highway 168 above Huntington Lake, 37°12'59.7"N, 119°11'30.4"W, 2 Sep. 2002, J.R. Shevock 22887 & B. Ertter (holotype, CAS; isotypes, H, KRAM, MO, NY).

⁶⁸ See Spence et al. (2022).

⁶⁹ Described as *Imbribryum torenii* by Spence & Shevock (2015) and accommodated here in *Bryum* under the new combination *Bryum torenii* (J.R.Spence & Shevock) W.R.Buck & Goffinet comb. nov. ≡ *Imbribryum torenii* J.R.Spence & Shevock, *Madroño* 62(1): 68. 2015. TYPE: U.S.A., California, Monterey Co.: Fort Hunter Liggett Military Reservation, along Gabilan Road at bridge crossing Nacimiento River, southern end of the Palisades, 35°51'05"N, 121°11'51"W, 10 Apr. 2004, J. R. Shevock et al. 24830 (holotype, CAS; isotypes, MO, NY).

⁷⁰ See Spence (2014).

⁷¹ See Spence (2014).

⁷² Described as *Gemmabryum vinosum* by Spence & Kellman (2015) and accommodated here in *Bryum* under the new combination *Bryum vinosum* (J.R.Spence & Kellman) W.R.Buck & Goffinet comb. nov. ≡ *Gemmabryum vinosum* J.R.Spence & Kellman, *Madroño* 62(2):127. 2015. TYPE:

⁵⁸ See Spence (2014).

⁵⁹ See Whitehouse (2001).

⁶⁰ See Kellman (2017).

⁶¹ See Toren et al. (2024).

⁶² Spence (2014) recognized *Gemmabryum badium*, which is considered conspecific with *Bryum imbricatulum* by Holyoak & Pedersen (2007), which at the species level takes priority.

⁶³ See Spence (2014).

⁶⁴ See Spence (2014).

⁶⁵ See Spence (2014).

⁶⁶ See Spence (2005).

⁶⁷ Described by Spence & Shevock (2012) as *Ptychostomum pacificum*; accommodated here in *Bryum*, under the new combination *Bryum pacificum* (J.R.Spence & Shevock) W.R.Buck & Goffinet comb. nov. ≡ *Ptychostomum pacificum*

- violaceum* Crundw. & Nyholm
warneum (Schwägr. ex Steud.) Brid.
weigeli Biehler
wrightii Sull. & Lesq.
- Buckia** D.Ríos, M.T.Gallego & J.Guerra⁷³ (PYLAISIACEAE)
vaucheri (Lesq.) D.Ríos, M.T.Gallego & J.Guerra
- Buxbaumia** Hedw. (BUXBAUMIACEAE)
aphylla Hedw.
minakatae S.Okamura
piperi Best
viridis (Moug. ex DC.) Brid. ex Moug. & Nestl.
- Calcidicranella** Bonfim Santos, Fedosov & Jan Kučera⁷⁴ (AONGSTROEMIACEAE)
howei (Renauld & Cardot) Bonfim Santos, Fedosov & Jan Kučera
pacifica (W.B.Schofield) Jan Kučera & Fedosov
varia (Hedw.) Bonfim Santos, Fedosov & Jan Kučera
- Callicladium** H.A.Crum⁷⁵ (PYLAISIACEAE)
fujiyamae (Broth.) Jan Kučera & Ignatov⁷⁶
haldaneanum (Grev.) H.A.Crum
imponens (Hedw.) Hedenäs, Schlesak & D.Quandt
- Callicostella** (Müll.Hal.) Mitt., *nom. cons.* (PILOTTRICHACEAE)
pallida (Hornsch.) Spruce
- Calliergon** (Sull.) Kindb. (CALLIERGONACEAE)
cordifolium (Hedw.) Kindb.
giganteum (Schimp.) Kindb.
megalophyllum Mikut., *nom. cons.*
orbicularicordatum (Renauld & Cardot) Broth.⁷⁷
richardsonii (Mitt.) Kindb. ex G.Roth
- Calliergonella** Loeske (PYLAISIACEAE)
curvifolia (Hedw.) B.H.Allen⁷⁸
- U.S.A., California, Fresno Co.: BLM San Joaquin River Gorge Management Area, along river via a 0.25 mi trail from powerhouse and parking lot, 37°04'37"N, 119°33'43"W, 26 Mar. 2009, J. R. Shevock et al. 32676 (holotype, CAS; isotypes, H, KRAM, MO, NY, UC).
- ⁷³ See Câmara et al. (2018).
- ⁷⁴ *Calcidicranella* was erected and segregated from *Dicranella* by Bonfim Santos et al. (Fedosov et al. 2023b) based on phylogenetic evidence.
- ⁷⁵ *Callicladium* comprised a single species until Kučera et al. (2019) and Schlesak et al. (2018) transferred *Hypnum fujiyamae* and *H. imponens*, respectively.
- ⁷⁶ The species was reported from North America by Schofield (2014) as *Hypnum fujiyamae* (Broth.) Paris.
- ⁷⁷ Ignatova et al. (2021) argued that *C. orbicularicordatum* falls within the variation of *C. cordifolium*; we tentatively retain it as distinct until further data are available.
- cuspidata* (Hedw.) Loeske
lindbergii (Mitt.) Hedenäs var. *lindbergii*⁷⁹
 var. *americana* (Renauld & Cardot) J.J.Atwood & Brinda⁸⁰
- Calliergonellopsis** Jan Kučera & Ignatov⁸¹ (PYLAISIACEAE)
dieckii (Renauld & Cardot) Jan Kučera & Ignatov
- Calymperes** Sw. ex F.Weber (CALYMPERACEAE)
afzelii Sw.
erosum Müll.Hal.
palisotii Schwägr.
pallidum Mitt.⁸²
tenerum Müll.Hal.
- Campylium** (Sull.) Spruce (AMBLYSTEGIACEAE)
bambergeri (Schimp.) Hedenäs, Schlesak & D.Quandt
chrysophyllum (Brid.) Lange
laxifolium Engelmark & Hedenäs
longicuspis (Lindb. & Arnell) Hedenäs
protensum (Brid.) Kindb.
stellatum (Hedw.) Lange & C.E.O.Jensen
- Campylophyllopsis** W.R.Buck⁸³ (AMBLYSTEGIACEAE)
hispidula (Brid.) Ochyra
sommerfeltii (Myrin) Ochyra
- Campylophyllum** (Schimp.) M.Fleisch.⁸⁴ (HYPNACEAE)
halleri (Hedw.) M.Fleisch.
montanum (Lindb.) B.H.Allen⁸⁵
- Campylopodiella** Cardot (DICRANACEAE)
flagellacea (Müll.Hal.) J.-P.Frahm & Isov.⁸⁶
stenocarpa (Wilson) P.Müll. & J.-P.Frahm
- Campylopus** Brid. (DICRANACEAE)
angustiretis (Austin) Lesq. & James
arctocarpus (Hornsch.) Mitt.
atrovirens De Not. var. *atrovirens*⁸⁷
 var. *cucullatifolius* J.-P.Frahm
carolinae Grout
flexuosus (Hedw.) Brid.
fragilis (Brid.) Bruch & Schimp.
gracilis (Mitt.) A.Jaege⁸⁸
- ⁷⁸ See Allen (2014).
- ⁷⁹ See Hedenäs (1990).
- ⁸⁰ See Atwood & Brinda (2020).
- ⁸¹ See Kučera et al. (2019).
- ⁸² See Buck (1994).
- ⁸³ See Goffinet et al. (2009b).
- ⁸⁴ See Kučera et al. (2019).
- ⁸⁵ See Allen (2014).
- ⁸⁶ See Shevock (2000), as *Atractylocarpus flagellaceus*.
- ⁸⁷ See Frahm (2007).
- ⁸⁸ See Frahm (2007).

- introflexus* (Hedw.) Brid.
oerstedianus (Müll.Hal.) Mitt.
pilifer Brid.
pyriformis (Schultz) Brid.
schimperi Milde
schmidii (Müll.Hal.) A.Jaeger⁸⁹
sinensis (Müll.Hal.) J.-P.Frahm⁹⁰
subulatus Schimp. ex Milde
surinamensis Müll.Hal.
tallulensis Sull. & Lesq.
- Campylostelium** Bruch & Schimp. (PTYCHOMITRIACEAE)
brachycarpum (Nog.) Z.Iwats., Tateishi & Tad. Suzuki⁹¹
laegerae Brinda, D.R.Toren & Shevock⁹²
pitardii (Corb.) E.Maier⁹³
saxicola (F.Weber & D.Mohr) Bruch & Schimp.
- Catosciopium** Brid. (CATOSCOPIACEAE)
nigratum (Hedw.) Brid.
- Ceratodon** Brid. (DITRICHACEAE)
 × *conicus* (Hampe ex Müll.Hal.) Lindb.
heterophyllus Kindb.
purpureus (Hedw.) Brid.
stenocarpus Bruch & Schimp.⁹⁴
- Chenia** R.H.Zander⁹⁵ (POTTIACEAE)
leptophylla (Müll.Hal.) R.H.Zander
- Chionoloma** Dixon⁹⁶ (POTTIACEAE)
cylindrotheca (Mitt.) M.Alonso, M.J.Cano & J. A.Jiménez
maragniphyllum (R.H.Zander & Eckel) W.R. Buck & Goffinet⁹⁷
orthodontum (Müll.Hal.) M.Alonso, M.J.Cano & J.A.Jiménez
- recurvifolium* (Taylor) M.Alonso, M.J.Cano & J. A.Jiménez
stenocarpum (Thér.) M.Alonso, M.J.Cano & J.A. Jiménez
tenuirostre (Hook. & Taylor) M.Alonso, M.J. Cano & J.A.Jiménez
- Chryso-hypnum** Hampe (HYPNACEAE)
diminutivum (Hampe) W.R.Buck
- Cinclidium** Sw. (MNIACEAE)
alaskanum R.E.Wyatt & A.H.Stoneb.⁹⁸
arcticum (Bruch & Schimp.) Schimp.
latifolium Lindb.
minutifolium Broth.⁹⁹
stygium Sw.
subrotundum Lindb.
- Cirriphyllum** Grout (BRACHYTHECIACEAE)
piliferum (Hedw.) Grout
- Claopodium** (Lesq. & James) Renaud & Cardot (LESKEACEAE)
bolanderi Best
crispifolium (Hook.) Renaud & Cardot
pellucinerve (Mitt.) Best
rostratum (Hedw.) Ignatov¹⁰⁰
whippleanum (Sull.) Renaud & Cardot
- Clasmatodon** Hook. & Wilson (BRACHYTHECIACEAE)
parvulus (Hampe) Sull.
- Cleistocarpidium** Ochyra & Bedn.-Ochyra¹⁰¹ (DITRICHACEAE)
palustre (Bruch & Schimp.) Ochyra & Bedn.-Ochyra
- Climacium** F.Weber & D.Mohr (CLIMACIACEAE)
americanum Brid.
dendroides (Hedw.) F.Weber & D.Mohr
kindbergii (Renaud & Cardot) Grout
- Cnestrum** I.Hagen¹⁰² (RHABDOWEISIACEAE)
alpestre (Wahlenb.) Nyholm
glaucescens (Lindb. & Arnell) Holmen ex Mogensen & Steere
schisti (F.Weber & D.Mohr) I.Hagen
- Codonoblepharon** Schwägr.¹⁰³ (ORTHOTRICHACEAE)
menziesii Schwägr.¹⁰⁴

⁸⁹ See Frahm (2007).⁹⁰ See Frahm (2007).⁹¹ See Iwatsuki et al. (1999).⁹² See Brinda et al. (2016).⁹³ See Brinda et al. (2016).⁹⁴ We retain this taxon as a distinct species following Shevock et al. (2021).⁹⁵ See Zander (1989).⁹⁶ See Alonso-García et al. (2016, 2019).⁹⁷ Described as *Oxystegus maragniphyllum* by Zander & Eckel (2019) and here accommodated in *Chionoloma* under the new combination *Chionoloma maragniphyllum* (R.H.Zander & Eckel) W.R.Buck & Goffinet comb. nov. ≡ *Oxystegus maragniphyllum* R.H.Zander & Eckel, *The Bryologist* 122(4): 569. 2019. TYPE: Canada, British Columbia: West Vancouver Island, Ocluelet, 7 May 1969, W.B. Schofield 89947 (holotype, UBC).⁹⁸ See Wyatt et al. (2021b).⁹⁹ See Wyatt & Stoneburner (2021).¹⁰⁰ See Ignatov et al. (2006).¹⁰¹ See Ochyra & Bednarek-Ochyra (1996).¹⁰² See Fedosov et al. (2021).¹⁰³ See Goffinet & Vitt (1998).¹⁰⁴ See Shevock (2000).

- Conardia** H.Rob. (AMBLYSTEGIACEAE)
compacta (Müll.Hal.) H.Rob.
- Conostomum** Sw. (BARTRAMIACEAE)
tetragonum (Hedw.) Lindb.
- Coscinodon** Spreng. (GRIMMIACEAE)
arctolimnius (Steere) Steere
calyptratus (Drumm.) C.E.O.Jensen.
cribrosus (Hedw.) Spruce
hartzii C.E.O.Jensen¹⁰⁵
yukonensis Hastings¹⁰⁶
- Cratoneuron** (Sull.) Spruce (AMBLYSTEGIACEAE)
filicinum (Hedw.) Spruce
- Crossidium** Jur., *nom. cons.* (POTTIACEAE)
aberrans Holz. & E.B.Bartram
crassinervium (De Not.) Jur. var. *crassisnervium*
seriatum H.A.Crum & Steere
squamiferum (Viv.) Jur. var. *squamiferum*
var. *pottioideum* (De Not.) Mönk.
- Crumia** W.B.Schofield (POTTIACEAE)
latifolia (Kindb.) W.B.Schofield
- Cryphaea** F.Weber (CRYPHAEACEAE)
filiformis (Hedw.) Brid.
glomerata Schimp. ex Sull.
nervosa (Hook. & Wilson) Müll.Hal.
ravenelii Austin
- Ctenidium** (Schimp.) Mitt. (CTENIDIACEAE)
schofieldii N.Nishim.
subrectifolium (Brid.) Pedano ex W.R.Buck & B.
H.Allen¹⁰⁷
- Cyclodictyon** Mitt. (PILOTRICHACEAE)
varians (Sull.) Kuntze
- Cynodontium** Bruch & Schimp., *nom. cons.*
(RHABDOWEISIACEAE)
gracilescens (F.Weber & D.Mohr) Schimp.
jenneri (Schimp.) Stirt.
polycarpon (Hedw.) Schimp.
strumiferum (Hedw.) Lindb.
strumulosum Müll.Hal. & Kindb.
tenellum (Schimp.) Limpr.
- Cyrtomnium** Holmen (MNIACEAE)
hymenophylloides (Huebener) T.J.Kop.
hymenophyllum (Bruch & Schimp.) Holmen
- Dacryophyllum** Ireland (HYPNACEAE)
falcifolium Ireland¹⁰⁸
- Daltonia** Hook. & Taylor, *nom. cons.* (DALTONIACEAE)
splachnoides (Sm.) Hook. & Taylor
- Dannorrhisia** Enroth (NECKERACEAE)
bigelovii (Sull.) Enroth¹⁰⁹
- Dendroalsia** E.Britton (CRYPHAEACEAE)
abietina (Hook.) E.Britton
- Dichelyma** Myrin (FONTINALACEAE)
capillaceum (Dicks.) Myrin
falcatum (Hedw.) Myrin
pallescens Bruch & Schimp.
uncinatum Mitt.
- Dichodontium** Schimp. (AONGSTROEMIACEAE)
olympicum Renaud & Cardot
pellucidum (Hedw.) Schimp.
- Dicranella** (Müll.Hal.) Schimp., *nom. cons.*
(DICRANELLACEAE)
cerviculata (Hedw.) Schimp.
heteromalla (Hedw.) Schimp.
hilariana (Mont.) Mitt.
lindigiana (Hampe) Mitt.
- Dicranellopsis** Bonfim Santos, Siebel & Fedosov¹¹⁰
(DICRANELLOPSIDACEAE)
crispa (Hedw.) Bonfim Santos, Siebel & Fedosov
subulata (Hedw.) Bonfim Santos, Siebel & Fedosov
- Dicranodontium** Bruch & Schimp. (DICRANACEAE)
asperulum (Mitt.) Broth.
denudatum (Brid.) E.Britton
uncinatum (Harv.) A.Jaeger
- Dicranowesia** Lindb. ex Milde (DICRANACEAE)
cirrata (Hedw.) Lindb. ex Milde
- Dicranum** Hedw. (DICRANACEAE)
acutifolium (Lindb. & Arnell) C.E.O.Jensen
bonjeanii De Not.
brevifolium (Lindb.) Lindb.
condensatum Hedw.
dispersum Engelmark¹¹¹
drummondii Müll.Hal.
elongatum Schleich. ex Schwägr.
flagellare Hedw.
flexicaule Brid.¹¹²
fragilifolium Lindb.
fulvum Hook.
fuscescens Turner

¹⁰⁸ The genus and species were established by Ireland (2004).¹⁰⁹ See Enroth et al. (2019).¹¹⁰ See Fedosov et al. (2023b).¹¹¹ See Afonina & Breen (2009).¹¹² See Tubanova et al. (2010).¹⁰⁵ See Hastings (1999).¹⁰⁶ See Hastings (1999).¹⁰⁷ See Buck & Allen (2004).

groenlandicum Brid.
hakkodense Cardot¹¹³
howellii Renaud & Cardot
leioneuron Kindb.
majus Turner var. *majus*
 var. *orthophyllum* A.Braun ex Milde
montanum Hedw.
muehlenbeckii Bruch & Schimp.
ontariense W.L.Peterson
pallidisetum (J.W.Bailey) Ireland
polysetum Sw.
rhabdocarpum Sull.
scoparium Hedw.
spadiceum J.E.Zetterst.
spurium Hedw.
tauricum Sapjegin
undulatum Schrad. ex Brid.
viride (Sull. & Lesq.) Lindb.
Didymodon Hedw. (POTTIACEAE)
acutus (Brid.) K.Saito
anserinocapitatus (X.J.Li) R.H.Zander¹¹⁴
cordatus Jur.¹¹⁵
ditrichoides (Broth.) X.J.Li & S.He
icmadophilus (Schimp. ex Müll.Hal.) K.Saito
maschalogena (Renaud & Cardot) Broth.
nigrescens (Mitt.) K.Saito
novae-hispaniae J.A.Jiménez & M.J.Cano¹¹⁶
perobtusus Broth.
rigidulus Hedw.
subandreaeoides (Kindb.) R.H.Zander
tectorum (Müll.Hal.) K.Saito
Diobelonella Ochyra¹¹⁷ (AONGSTROEMICEAE)
palustris (Dicks.) Ochyra
Diphyscium D.Mohr (DIPHYSICACEAE)
foliosum (Hedw.) D.Mohr
mucronifolium Mitt. ex Dozy & Molk.¹¹⁸
Discelium Brid. (DISCELIACEAE)
nudum (Dicks.) Brid.

Distichium Bruch & Schimp., *nom. cons.* (DISTICHIACEAE)
capillaceum (Hedw.) Bruch & Schimp.
hagenii Ryan ex H.Philib.
inclinatum (Hedw.) Bruch & Schimp.
Ditrichum Timm ex Hampe, *nom. cons.* (DITRICHACEAE)
ambiguum Best
divaricatum Mitt.¹¹⁹
heteromallum (Hedw.) E.Britton
lineare (Sw.) Lindb.
montanum Leiberg
pallidum (Hedw.) Brockm.
pusillum (Hedw.) Hampe
rhynchostegium Kindb.
schimperi (Lesq.) Kuntze
septentrionale Fedosov, Jan Kučera & Ignatova¹²⁰
tortuloides Grout
zonatum (Brid.) Braithw.¹²¹
Donnellia Austin (SEMATOPHYLLACEAE)
commutata (Müll.Hal.) W.R.Buck
Donrichardia H.A.Crum & L.E.Anderson
 (BRACHYTHECIACEAE)
macroneuron (Grout) H.A.Crum & L.E.Anderson
pringlei (Cardot) Huttunen & Ignatov¹²²
Drepanium (Schimp.) Spruce (AMBLYSTEGICEAE)
fastigiatum (Brid.) Lange & C.E.O.Jensen¹²³
Drepanocladus (Müll.Hal.) G.Roth, *nom. cons.*
 (AMBLYSTEGICEAE)
aduncus (Hedw.) Warnst.
angustifolius (Hedenäs) Hedenäs & Rosborg
arcticus (R.S.Williams) Hedenäs
brevifolius (Lindb.) Warnst.
capillifolius (Warnst.) Warnst.¹²⁴
cardotii (Thér.) Hedenäs¹²⁵
latinervis Warnst.¹²⁶
polygamus (Schimp.) Hedenäs
sordidus (Müll.Hal.) Hedenäs
trifarius (F.Weber & D.Mohr) Broth.
turgescens (T.Jensen) Broth.

¹¹³ According to Huang et al. (2024) and Ignatova & Fedosov (2008) specimens from the West Coast of North America considered to belong to *D. viride* are most likely populations of *D. hakkodense*.

¹¹⁴ See Zander & Weber (1997).

¹¹⁵ See Allred et al. (2024).

¹¹⁶ See Jiménez et al. (2022).

¹¹⁷ Established for *Dicranella palustris* (Dicks.) Crundw. ex E.F.Warb. by Ochyra et al. (2003).

¹¹⁸ See Magombo (2003).

¹¹⁹ See Fedosov et al. (2023a).

¹²⁰ See Fedosov et al. (2024).

¹²¹ See Matsui & Iwatsuki (1990).

¹²² See Huttunen et al. (2007).

¹²³ See Schlesak et al. (2018).

¹²⁴ See Saługa et al. (2018); treated as *D. longifolius* earlier by Hedenäs (2014b).

¹²⁵ See Hedenäs (1997).

¹²⁶ See Hedenäs (1998).

Drummondia Hook., *nom. cons.* (DRUMMONDIACEAE)
prorepens (Hedw.) Trevis.

Dryptodon Brid.¹²⁷ (GRIMMIACEAE)
anomalus (Hampe ex Schimp.) Loeske
arcuatifolius (Kindb.) Ochyra & Żarnowiec¹²⁸
fuscoluteus (Hook.) Ochyra & Żarnowiec¹²⁹
hartmanii (Schimp.) Limpr.
leibergii (Paris) Ochyra & Żarnowiec
lisae (De Not.) Loeske¹³⁰
muehlenbeckii (Schimp.) Loeske
patens (Dicks. ex Hedw.) Brid.
torquatus (Hook.) Brid.
trichophyllus (Grev.) Brid.

Eccremidium Wilson (DICRANELLACEAE)
floridanum H.A.Crum

Echinophyllum T.J.O'Brien (THUIDIACEAE)
sachalinensis (Lindb.) T.J.O'Brien¹³¹

Encalypta Hedw. (ENCALYPTACEAE)
affinis R.Hedw. subsp. *affinis*
subsp. *macounii* (Austin) D.G.Horton
alpina Sm.
brevicolla (Bruch & Schimp.) Ångstr. subsp. *brevicolla*
subsp. *crumiana* D.G.Horton
brevipes Schljakov
ciliata Hedw.
flowersiana D.G.Horton
longicolla Bruch
mutica I.Hagen
pilifera Funck¹³²
procera Bruch
rhaptocarpa Schwägr.
spathulata Müll.Hal.
texana Magill
vittiana D.G.Horton
vulgaris Hedw.

Entodon Müll.Hal. (ENTODONTACEAE)
beyrichii (Schwägr.) Müll.Hal.
brevisetus (Hook. & Wilson) Lindb.

¹²⁷ *Dryptodon* was recovered as a clade of ambiguous affinities (Hernández-Maqueda et al. 2008, Larraín et al. 2013) and is here treated sensu Ochyra et al. (2003).

¹²⁸ This name has been proposed as a synonym of either *Grimmia lisae* (Hastings & Greven 2007) or *G. decipiens* (Maier 2010); we propose to retain the name, albeit in *Dryptodon*, pending clarification.

¹²⁹ See Muñoz (1999) and Muñoz & Pando (2000).

¹³⁰ See Streiff (2006).

¹³¹ See O'Brien & Horton (2000).

¹³² See Fedosov (2012).

challengeri (Paris) Cardot
cladorrhizans (Hedw.) Müll.Hal.
concinus (DeNot.) Paris
hampeanus Müll.Hal.
macropodus (Hedw.) Müll.Hal.
schleicheri (Schimp.) Demet.
seductrix (Hedw.) Müll.Hal.
sullivantii (Müll.Hal.) Lindb.

Entodontopsis Broth. (STEREOPHYLLACEAE)
leucostega (Brid.) W.R.Buck & Ireland

Entosthodon Schwägr. (FUNARIACEAE)
americanus (Lindb.) Fife
apiculatopilosus (Cardot) Fife
attenuatus (Dicks.) Bryhn
bolanderi Lesq.
californicus (Sull. & Lesq.) H.A.Crum & L.E.
Anderson

convexus (Spruce) Brugués
drummondii Sull.
fascicularis (Hedw.) Müll.Hal.
kochii H.A.Crum & L.E.Anderson
muhlenbergii (Turner) Fife
planoconvexus (E.B.Bartram) Grout
rubiginosus (R.S.Williams) Grout
rubrisetus (E.B.Bartram) Grout
serratus (Brid.) Fife
sonorae (Cardot) Steere
tucsonii (E.B.Bartram) Grout
wigginsii Steere

Ephemerum Hampe, *nom. cons.* (POTTIACEAE)
cohaerens (Hedw.) Hampe
crassinervium (Schwägr.) Hampe var. *crassinervium*
var. *texanum* (Grout) V.S.Bryan & L.E.
Anderson

recurvifolium (Dicks.) Boulay¹³³
serratum (Schreb. ex Hedw.) Hampe
spinulosum Bruch & Schimp.

Epipterygium Lindb. (MIELICHHOFERIACEAE)
biauratum Hanusch¹³⁴

Erpodium (Brid.) Brid. (ERPODIACEAE)
domingense (Spreng.) Brid.

Eucladium Bruch & Schimp. (POTTIACEAE)
verticillatum (With.) Bruch & Schimp.

¹³³ See Snyder (2023).

¹³⁴ This name replaces *E. tozeri* for the North American flora (Hanush et al. 2020).

Eurhynchiastrum Ignatov & Huttunen¹³⁵
(BRACHYTHECIACEAE)

pulchellum (Hedw.) Ignatov & Huttunen var.
pulchellum
var. *barnesii* (Renauld & Cardot) Ignatov

Fabronia Raddi (FABRONIACEAE)

ciliaris (Brid.) Brid. var. *ciliaris*
var. *polycarpa* (Hook.) W.R.Buck
var. *wrightii* (Sull.) W.R.Buck
pusilla Raddi

Fissidens Hedw. (FISSIDENTACEAE)

adanthoides Hedw.
amoenus Müll.Hal.
aphelotaxifolius Pursell
appalachensis R.H.Zander
arcticus Bryhn
asplenioides Hedw.
bryoides Hedw.
bushii (Cardot & Thér.) Cardot & Thér.
closteri Austin
crispus Mont.
curvatus Hornsch.
dubius P.Beauv.
elegans Brid.
exilis Hedw.
fontanus (Bach.Pyl.) Steud.
grandifrons Brid.
hallianus (Sull. & Lesq.) Mitt.
hyalinus Wilson & Hook.
leptophyllus Mont.
littlei (R.S.Williams) Grout
minutulus Sull.
obtusifolius Wilson
osmundioides Hedw.
pallidinervis Mitt.
pauperculus M.Howe
pellucidus Hornsch.
polypodioides Hedw.
santa-clarensis Thér.
scalaris Mitt.
serratus Müll.Hal.
subbasilaris Hedw.
sublimbatus Grout
submarginatus Bruch
taxifolius Hedw.
taylorii Müll.Hal.

¹³⁵ See Ignatov & Huttunen (2002).

ventricosus Lesq.
zollingeri Mont.

Flexitrichum Ignatov & Fedosov¹³⁶ (FLEXITRICHACEAE)

flexicaule (Schwägr.) Ignatov & Fedosov
gracile (Mitt.) Ignatov & Fedosov

Fontinalis Hedw. (FONTINALACEAE)

antipyretica Hedw.
dalecarlica Bruch & Schimp.
howellii Renauld & Cardot
hypnoides Hartm.
neomexicana Sull. & Lesq.
novae-angliae Sull.
redfearnii B.H.Allen¹³⁷
sphagnifolia (Müll.Hal.) Wijk & Margad.
sullivantii Lindb.
welchiana B.H.Allen

Forsstroemia Lindb. (NECKERACEAE)

producta (Hornsch.) Paris
trichomitria (Hedw.) Lindb.

Funaria Hedw. (FUNARIACEAE)

arctica (Berggr.) Kindb.
calvescens Schwägr.
convexa Spruce¹³⁸
flavicans Michx.
hygrometrica Hedw.
microstoma Bruch ex Schimp.
polaris Bryhn

Geheebia Schimp.¹³⁹ (POTTIACEAE)

constricta (Mitt.) R.H.Zander & Caners¹⁴⁰
fallax (Hedw.) R.H.Zander
ferruginea (Schimp. ex Besch.) R.H.Zander
gigantea (Funck) Boulay
leskeoides (K.Saito) R.H.Zander
lurida (Hornsch. ex Spreng.) J.A.Jiménez & M.J.
Cano
maxima (Syed & Crundw.) R.H.Zander
tophacea (Brid.) R.H.Zander

Gertrudiella Broth.¹⁴¹ (POTTIACEAE)

nevadensis (R.H.Zander) J.A.Jiménez & M.J.Cano

Globulinella Steere (POTTIACEAE)

globifera (Hampe) Steere

¹³⁶ See Fedosov et al. (2016b).¹³⁷ *Fontinalis redfearnii* was described by Allen (1991) and is known from Oklahoma.¹³⁸ See Toren (2008).¹³⁹ See Jiménez et al. (2022).¹⁴⁰ See Zander & Caners (2017).¹⁴¹ See Jiménez et al. (2022).

Gollania Broth. (HYPNACEAE)*turgens* (Müll.Hal.) Ando**Grimmia** Hedw. (GRIMMIACEAE)*alpestris* (F.Weber & D.Mohr) Schleich.*americana* E.B.Bartram*anodon* Bruch & Schimp.*arizonae* Renauld & Cardot*atrata* Miel. ex Hornsch.*brittoniae* R.S.Williams*caespiticia* (Brid.) Jur.*crinitoleucophaea* Cardot*donniana* Sm.*elatior* Bruch ex Bals.-Criv. & De Not.*elongata* Kaulf.*funalis* (Schwägr.) Bruch ex De Not.*hamulosa* Lesq.*incurva* Schwägr.*insolita* J.Muñoz, I.Solano & D.Quandt¹⁴²*laevigata* (Brid.) Brid.*lesherae* Greven*longirostris* Hook.*mariniana* Sayre*mollis* Bruch & Schimp.*montana* Bruch & Schimp.*moxleyi* R.S.Williams*nevadensis* Greven*olneyi* Sull.*orbicularis* Bruch ex Wilson*ovalis* (Hedw.) Lindb.*pilifera* P.Beauv.*plagiopodia* Hedw.*pulvinata* (Hedw.) Sm.*reflexidens* Müll.Hal.*serrana* J.Muñoz, Shevock & D.R.Toren*sessitana* De Not.*shastae* Greven*shevockii* J.Muñoz, I.Solano & D.Quandt¹⁴³*tergestina* Tomm. ex Bruch & Schimp.¹⁴⁴*texicana* Greven¹⁴⁵*toreni* Hastings¹⁴⁶¹⁴² See Solano et al. (2023).¹⁴³ See Solano et al. (2023).¹⁴⁴ See Allred et al. (2024).¹⁴⁵ See Greven (2010).¹⁴⁶ See Hastings (2008). However, Maier (2010) synonymized this name with *G. tergestina*, but more recently it has been recognized by California botanists (Toren 2015; Rae 2021; McLaughlin 2024).*ungeri* Jur.¹⁴⁷*unicolor* Hook. ex Grev.**Groutiella** Steere (ORTHOTRICHACEAE)*tomentosa* (Hornsch.) Wijk & Margad.*tumidula* (Mitt.) Vitt**Gymnostomiella** M.Fleisch. (POTTIACEAE)*vernica* (Hook. ex Harv.) M.Fleisch.¹⁴⁸**Gymnostomum** Nees & Hornsch., *nom. cons.* (POTTIACEAE)*aeruginosum* Sm.*calcareum* Nees & Hornsch.*viridulum* Brid.**Gyroweisia** Schimp., *nom. cons.* (POTTIACEAE)*tenuis* (Hedw.) Schimp.**Hageniella** Broth. (HYPNACEAE)*micans* (Mitt.) B.C.Tan & Y.Jia¹⁴⁹**Hamatocaulis** Hedenäs (CALLIERGONACEAE)*lapponicus* (Norrl.) Hedenäs*vernicosus* (Mitt.) Hedenäs**Haplocladium** (Müll.Hal.) Müll.Hal., *nom. cons.*¹⁵⁰

(LESKEACEAE)

angustifolium (Hampe & Müll.Hal.) Broth.*microphyllum* (Sw. ex Hedw.) Broth.*virginianum* (Brid.) Broth.**Haplodontium** Hampe¹⁵¹ (MNIACEAE)*macrocarpum* (Drumm.) J.R.Spence*tehamense* (Showers) J.R.Spence**Hedwigia** P.Beauv., *nom. cons.* (HEDWIGIACEAE)*ciliata* (Hedw.) Boucher*detonsa* (M.Howe) W.R.Buck & D.H.Norris*emodica* Hampe ex Müll.Hal.¹⁵²*filiformis* (Michx.) P.Beauv.¹⁵³*nivalis* (Müll.Hal.) Mitt.¹⁵⁴*stellata* Hedenäs**Helodium** Warnst., *nom. cons.* (HELODIACEAE)*blandovii* (F.Weber & D.Mohr) Warnst.¹⁴⁷ See Muñoz (1999).¹⁴⁸ Pending further studies, we follow Redfearn (1991), Arts (1998) and Allen (2002), versus Zander (2007b), who argued for retaining the name *G. orcuttii* E.B.Bartram for the North American populations.¹⁴⁹ See Tan & Jia (1999).¹⁵⁰ See Wilson (2021).¹⁵¹ See Spence (2005).¹⁵² The species may be present in North America based on herbarium records (Gbif 2024b) although it has not formally been reported from North America.¹⁵³ See De Luna (2022).¹⁵⁴ Reported from the southwestern and south-central United States by Allen (2010); see Gbif.org (2024c).

- elodioides* (Renauld & Cardot) B.H.Allen
paludosum (A.Jaeger) Austin ex C.F.Parker
Henicodidium (Müll.Hal.) Kindb. (PTEROBRYACEAE)
geniculatum (Mitt.) W.R.Buck
Hennediella Paris¹⁵⁵ (POTTIACEAE)
heimii (Hedw.) R.H.Zander
stanfordensis (Steere) Blockeel
Herpetineuron (Müll.Hal.) Cardot (THUIDIACEAE)
toccoae (Sull. & Lesq.) Cardot
Herzogiella Broth. (PLAGIOTHECIACEAE)
seligeri (Brid.) Z.Iwats.
striatella (Brid.) Z.Iwats.
turfacea (Lindb.) Z.Iwats.
Heterocladiella Ignatov & Fedosov¹⁵⁶
 (HETEROCLADIACEAE)
dimorpha (Brid.) Ignatov & Fedosov
procurrens (Mitt.) Ignatov & Fedosov
Heterocladium Schimp. (PTERIGYNANDRACEAE)
macounii Best
Heterophyllum (Schimp.) Müll.Hal. ex Kindb.
 (SEMATOPHYLLACEAE)
nemorosum (W.D.J.Koch ex Brid.) Kindb.
Hilpertia R.H.Zander¹⁵⁷ (POTTIACEAE)
velenovskii (Schiffn.) R.H.Zander
Homalia Brid., *nom. cons.* (NECKERACEAE)
trichomanoides (Hedw.) Brid.
Homaliadelphus Dixon & P.de la Varde (MIYABACEAE)
sharpii (R.S.Williams) Sharp
Homalotheciella (Cardot) Broth. (BRACHYTHECIACEAE)
subcapillata (Hedw.) Broth.
Homalothecium Schimp. (BRACHYTHECIACEAE)
aeneum (Mitt.) E.Lawton
arenarium (Lesq.) E.Lawton
aureum (Spruce) H.Rob.
californicum Hedenäs
fulgescens (Mitt. ex Müll.Hal.) A.Jaeger
megaptilum (Sull.) H.Rob.
nevadense (Lesq.) Renauld & Cardot
nuttallii (Wilson) A.Jaeger
sericeum (Hedw.) Schimp.
Homomallium (Schimp.) Loeske (HYPNACEAE)
adnatum (Hedw.) Broth.
incurvatum (Schrad. ex Brid.) Loeske
mexicanum Cardot var. *mexicanum*
 var. *latifolium* Cardot
- Hookeria** Sm., *nom. cons.* (HOOKERIAEAE)
acutifolia Hook. & Grev.
lucens (Hedw.) Sm.
Husnotiella Cardot¹⁵⁸ (POTTIACEAE)
asperifolia (Mitt.) J.A.Jiménez & M.J.Cano
fragilicuspis (Broth) J.A.Jiménez & M.J.Cano
johansenii (R.S.Williams) J.A.Jiménez & M.J.Cano
revoluta Cardot
Hydrogonium (Müll.Hal.) A.Jaeger¹⁵⁹ (POTTIACEAE)
amplexifolium (Mitt.) P.C.Chen
bolleanum (Müll.Hal.) A.Jaeger
consanguineum (Thwaites & Mitt.) Hilp. var.
cancellatum (Müll.Hal.) Jan Kučera
cruegeri (Sond. ex Müll.Hal.) Jan Kučera
gregarium (Mitt.) Jan Kučera
orientale (F.Weber) Jan Kučera
Hygroamblystegium Loeske, *nom. cons.*¹⁶⁰
 (AMBLYSTEGIACEAE)
fluviatile (Hedw.) Loeske
humile (P.Beauv.) Vanderp., Hedenäs & Goffinet
noterophilum (Sull. & Lesq.) Warnst.
tenax (Hedw.) Jenn.
varium (Hedw.) Mönk.
Hygrohypnella Ignatov & Ignatova¹⁶¹ (AMBLYSTEGIACEAE)
ochracea (Turner ex Wilson) Ignatov & Ignatova
polaris (Lindb.) Ignatov & Ignatova
Hygrohypnum Lindb. (AMBLYSTEGIACEAE)
closteri (Austin) Grout
luridum (Hedw.) Jenn.
styriacum (Limpr.) Broth.
Hylocomiadelphus Ochyra & Stebel¹⁶² (HYLOCOMIACEAE)
triquetrus (Hedw.) Ochyra & Stebel
Hylocomiastrum M.Fleisch. ex Broth. (HYLOCOMIACEAE)
pyrenaicum (Spruce) M.Fleisch. ex Broth.
umbratum (Hedw.) M.Fleisch. ex Broth.
Hylocomium Schimp., *nom. cons.* (HYLOCOMIACEAE)
splendens (Hedw.) Schimp.
Hymenoloma Dusén¹⁶³ (HYMENOLOMATAEAE)
crispulum (Hedw.) Ochyra
mulahaceni (Höhn.) Ochyra¹⁶⁴

¹⁵⁵ See Cano (2008).¹⁵⁶ See Ignatov et al. (2019a).¹⁵⁷ See Zander (1989).¹⁵⁸ See Jiménez et al. (2022).¹⁵⁹ See Kučera et al. (2013).¹⁶⁰ *Hygroamblystegium* is, albeit not robustly, resolved as a sister group to *Amblystegium* (Vanderpoorten et al. (2002), which holds a single species. The two could be merged into a single genus, as adopted by Anderson et al. (1990).¹⁶¹ See Ignatov & Ignatova (2004).¹⁶² See Ochyra & Stebel (2008) and Ignatov et al. (2019b).¹⁶³ See Ochyra et al. (2003).

- Hymenostylium** Brid.¹⁶⁵ (POTTIACEAE)
annotinum Mitt. ex Dixon
aurantiacum Mitt.¹⁶⁶
recurvirostrum (Hedw.) Dixon
- Hyophila** Brid., *nom. cons.* (POTTIACEAE)
involuta (Hook.) A.Jaeger
- Hyophiladelphus** (Müll.Hal.) R.H.Zander (POTTIACEAE)
agrarius (Hedw.) R.H.Zander¹⁶⁷
- Hypnum** Hedw., *nom. cons.* (HYPNACEAE)
andoi A.J.E.Sm.¹⁶⁸
cupressiforme Hedw. var. *cupressiforme*
var. *filiforme* Brid.
var. *julaceum* Brid.
jutlandicum Holmen & E.Warncke
resupinatum Taylor
subjulaceum (Molendo) Hedenäs, Schlesak & D.
Quandt
- Hypopterygium** Brid. (HYPOPTERYGIACEAE)
flavolimbatum Müll.Hal.
tamarisci (Sw.) Brid. ex Müll.Hal.
- Indusiella** Broth. & Müll.Hal. (PTYCHOMITRIACEAE)
thianschanica Broth. & Müll.Hal.
- Isopterygiella** Ignatov & Ignatova¹⁶⁹ (PLAGIOTHECIACEAE)
alpicola (Lindb.) Ignatov & Ignatova
pulchella (Hedw.) Ignatov & Ignatova
- Isopterygiopsis** Z.Iwats. (PLAGIOTHECIACEAE)
catagonioides (Broth.) Ignatov & Ignatova¹⁷⁰
muelleriana (Schimp.) Z.Iwats.
- Isopterygium** Mitt. (PYLAIADDELPHACEAE)
tenerifolium Mitt.¹⁷¹
tenerum (Sw.) Mitt.
- Isothecium** Brid. (LEMBOPHYLLACEAE)
alopecuroides (Lam. ex Dubois) Isov.
- Iwatsukiella** W.R.Buck & H.A.Crum (PTERIGYNANDRACEAE)
leucotricha (Mitt.) W.R.Buck & H.A.Crum
- Jaegerina** Müll.Hal. (PTEROBRYACEAE)
scariosa (Lorentz) Arzeni
- Jaffuelobryum** Thér. (GRIMMIACEAE)
raui (Austin) Thér.
wrightii (Sull.) Thér.
- Jochenia** Hedenäs, Schlesak & D.Quandt¹⁷² (JOCHENIACEAE)
pallescens (Hedw.) Hedenäs, Schlesak & D.Quandt
protuberans (Brid.) Jan Kučera & Ignatov¹⁷³
- Kiaeria** I.Hagen (RHABDOWEISIACEAE)
falcata (Hedw.) I.Hagen¹⁷⁴
- Kindbergia** Ochya¹⁷⁵ (BRACHYTHECIACEAE)
oregana (Sull.) Ochya
praelonga (Hedw.) Ochya
- Koponeniella** Huttunen & Ignatov¹⁷⁶ (BRACHYTHECIACEAE)
bolanderi (Lesq.) Huttunen & Ignatov
graminicolor (Brid.) Huttunen, Ignatov, M.Li &
Y.F.Wang
- Lepidopilum** (Brid.) Brid., *nom. cons.* (PILOTRICHACEAE)
polytrichoides (Hedw.) Brid.
- Leptobryum** (Schimp.) Wilson (MEESIACEAE)
pyriforme (Hedw.) Wilson
- Leptodictyum** (Schimp.) Warnst. (AMBLYSTEGIACEAE)
riparium (Hedw.) Warnst.
wallacei B.H.Allen & Magill¹⁷⁷
- Leptodon** D.Mohr, *nom. cons.* (LEPTODONTACEAE)
smithii (Dicks. ex Hedw.) F.Weber & D.Mohr
- Leptodontium** (Müll.Hal.) Hampe ex Lindb.
(POTTIACEAE)
excelsum (Sull.) E.Britton
flexifolium (With.) Hampe ex Lindb.
- Leptohymenium** Schwägr. (HYLOCOMIACEAE)
sharpii (H.A.Crum & L.E.Anderson) W.R.Buck
& H.A.Crum
- Leptopterigynandrum** Müll.Hal. (TAXIPHYLLACEAE)
austroalpinum Müll.Hal.
subintegrum (Mitt.) Broth.¹⁷⁸

¹⁶⁴ Reported from North America by Werner et al. (2013).

¹⁶⁵ Zander & Hedderson (2016) segregated North American and other species of *Hymenostylium* in their new genus *Ardeuma*. However, the phylogenetic study by Cano et al. (2022) highlighted ambiguous monophyly of this and other related genera, and hence we prefer to retain a conservative, i.e., broader, circumscription of *Hymenostylium*. Hodgetts et al. (2020) rejected recognizing *Ardeuma* on the basis of unpublished data, which may be forthcoming.

¹⁶⁶ Reported from North America by Zander & Eckel (2017).

¹⁶⁷ See Zander (1995).

¹⁶⁸ Potentially doubtful, see Allen (2014).

¹⁶⁹ See Ignatova et al. (2020a).

¹⁷⁰ See Ignatova et al. (2020a).

¹⁷¹ See Buck (1994).

¹⁷² *Jochenia* was erected by Schlesak et al. (2018) to accommodate *Hypnum pallescens*.

¹⁷³ See Kučera et al. (2019).

¹⁷⁴ See Brinda & Fedosov (2023).

¹⁷⁵ See Li et al. (2015).

¹⁷⁶ *Koponeniella* was established based on phylogenetic evidence (Huttunen et al. 2015).

¹⁷⁷ Described by Allen & Magill (2004) for material from Texas.

¹⁷⁸ Reported by He (2005).

Leratia Broth. & Paris (ORTHOTRICHACEAE)*exigua* (Sull.) Goffinet¹⁷⁹**Lescuraea** Schimp.¹⁸⁰ (LESKEACEAE)*arizonae* (R.S.Williams) P.S.Wilson & D.H.Norris*atricha* (Kindb.) E.Lawton*baileyi* (Best & Grout) E.Lawton*incurvata* (Hedw.) E.Lawtonvar. *gigantea* E.Lawtonvar. *tenuiretis* (Culm.) E.Lawton*patens* (Lindb.) Arnell & C.E.O.Jensen*radicosa* (Mitt.) Mönk. var. *radicosa*var. *compacta* (Best) E.Lawtonvar. *denudata* (Kindb.) E.Lawtonvar. *pallida* (Best) E.Lawton*saviana* (De Not.) E.Lawton*saxicola* (Schimp.) Molendo*stenophylla* (Renauld & Cardot) Kindb.*tribulosa* (Shevock & W.R.Buck) W.R.Buck & Goffinet¹⁸¹**Leskea** Hedw. (LESKEACEAE)*australis* Sharp*gracilescens* Hedw.*obscura* Hedw.*polycarpa* Hedw.**Leucobryum** Hampe (LEUCOBRYACEAE)*albidum* (Brid. ex P.Beauv.) Lindb.*antillarum* Schimp. ex Besch.*glaucum* (Hedw.) Ångstr.**Leucodon** Schwägr. (LEUCODONTACEAE)*brachypus* Brid.*julaceus* (Hedw.) Sull.*sciuroides* (Hedw.) Schwägr.¹⁸²**Leucolepis** Lindb. (MNIACEAE)*acanthoneura* (Schwägr.) Lindb.**Lewinskya** F.Lara, Garilleti & Goffinet¹⁸³
(ORTHOTRICHACEAE)*affinis* (Brid.) F.Lara, Garilleti & Goffinet var. *affinis*
var. *bohemica* (Plášek & Sawicki) Plášek¹⁸⁴*arida* Vigalondo, F.Lara & Garilleti¹⁸⁵*bolanderi* (Sull.) F.Lara, Garilleti & Goffinet*elegans* (Schwägr. ex Hook. & Grev.) F.Lara, Garilleti
& Goffinet*fenestrata* (Cardot & Thér.) F.Lara, Garilleti &
Goffinet*holzingeri* (Renauld & Cardot) F.Lara, Garilleti
& Goffinet*keeverae* (H.A.Crum & L.E.Anderson) F.Lara,
Garilleti & Goffinet*laevigata* (J.E.Zetterst.) F.Lara, Garilleti & Goffinet*pacifica* Vigalondo, F.Lara & Garilleti¹⁸⁶*praemorsa* (Venturi) F.Lara, Garilleti & Goffinet*pseudoaffinis* Vigalondo, F.Lara & Garilleti¹⁸⁷*pycnophylla* (Schimp.) F.Lara, Garilleti & Goffinet*pylaisii* (Brid.) F.Lara, Garilleti & Goffinet*rupestris* (Schleich. ex Schwägr.) F.Lara, Garilleti
& Goffinet*shawii* (Wilson) F.Lara, Garilleti & Goffinet¹⁸⁸*sordida* (Sull. & Lesq.) F.Lara, Garilleti &
Goffinet*speciosa* (Nees) F.Lara, Garilleti & Goffinet*spjutii* (D.H.Norris & Vitt) F.Lara, Garilleti &
Goffinet¹⁸⁹*striata* (Hedw.) F.Lara, Garilleti & Goffinet**Limbella** (Müll.Hal.) Renauld & Cardot
(AMBLYSTEGIACEAE)*fryei* (R.S.Williams) Ochyra**Lindbergia** Kindb. (LESKEACEAE)*brachyptera* (Mitt.) Kindb.*mexicana* (Besch.) Cardot**Loeskeobryum** M.Fleisch. ex Broth. (HYLOCOMIACEAE)*brevirostre* (Brid.) M.Fleisch. ex Broth.**Lorentziella** Müll.Hal. ex Besch. (GIGASPERMACEAE)*imbricata* (Mitt.) Broth.¹⁷⁹ See Goffinet et al. (2004)¹⁸⁰ We follow Gardiner et al. (2005) who merged *Pseudoleskea* with *Lescuraea*.¹⁸¹ Described as *Pseudoleskea tribulosa* by Shevock & Buck (2009) and accommodated in *Lescuraea* under the new combination *Lescuraea tribulosa* (Shevock & W.R.Buck) W.R.Buck & Goffinet comb. nov. ≡ *Pseudoleskea tribulosa* Shevock & W.R.Buck, *The Bryologist* 112(1): 184. 2009. TYPE: U.S.A., California, Mariposa Co.: Yosemite National Park, Hoffmann Creek watershed S of Mt. Hoffmann and N of Tioga Road, 37°49'52"N, 119°30'50"W, 4 Aug. 2006, J.R. Shevock et al. 28987 (holotype, CAS; isotypes, DUKE, H, MO, NY, UC, US, YM).¹⁸² Previously this taxon was known as *L. brachypus* var. *andrewsianus* Crum & Anderson (Stech et al. 2011).¹⁸³ See Lara et al. (2016).¹⁸⁴ Based on a collection, *D. H. Vitt 24* (MO) from Idaho; but see Vigalondo et al. (2020) who considered this taxon a form of *L. affinis*.¹⁸⁵ See Vigalondo et al. (2020).¹⁸⁶ See Vigalondo et al. (2020).¹⁸⁷ See Vigalondo et al. (2020).¹⁸⁸ See Garilleti et al. (2006b).¹⁸⁹ See Norris & Vitt (1993).

- Luisierella** Thér. & P.de la Varde (TIMMIELLACEAE)
barbula (Schwägr.) Steere
- Lyellia** R.Br. (POLYTRICHACEAE)
aspera (I.Hagen & C.E.O.Jensen) Frye
- Macrocoma** (Hornsch. ex Müll.Hal.) Grout
 (ORTHOTRICHACEAE)
tenuis (Hook. & Grev.) Vitt subsp. *sullivantii*
 (Müll.Hal.) Vitt
- Macromitrium** Brid. (ORTHOTRICHACEAE)
viticulosum (Raddi) Brid.¹⁹⁰
- Meesia** Hedw., *nom. cons.* (MEESIACEAE)
longiseta Hedw.
triquetra (L. ex Jolycl.) Ångstr.
uliginosa Hedw.
- Merceyopsis** Broth. & Dixon (POTTIACEAE)
cataractae (Mitt.) Brinda, Ignatov & Fedosov¹⁹¹
- Metaneckera** Steere (NECKERACEAE)
menziesii (Drumm.) Steere
- Meteorium** Dozy & Molk., *nom. cons.* (METEORACEAE)
nigrescens (Hedw.) Dozy & Molk.¹⁹²
- Microbryum** Schimp.¹⁹³ (POTTIACEAE)
commutatum (Limpr.) Cl.Schneid., Th.Schneid.
 & Mahévas¹⁹⁴
conicum (Schleich. ex Schwägr.) Cl.Schneid.,
 Th.Schneid. & Mahévas¹⁹⁵
davallianum (Sm.) R.H.Zander¹⁹⁶
floerkeanum (F.Weber & D.Mohr) Schimp.
fosbergii (E.B.Bartram) Ros, O.Werner & Rams¹⁹⁷
muticum (Venturi) Cl.Schneid., Th.Schneid. &
 Mahévas¹⁹⁸
starckeanum (Hedw.) R.H.Zander
- Micromitrium** Austin (MICROMITRIACEAE)
megalosporum Austin
synoicum (James) Austin
tenerum (Bruch & Schimp.) Crosby
wrightii (Müll.Hal.) Crosby
- Mielichhoferia** Hornsch. (MIELICHHOFERACEAE)
elongata (Hoppe & Hornsch. ex Hook.) Hornsch.
mielichhoferiana (Funck) Loeske
shevockii (A.J.Shaw) A.J.Shaw¹⁹⁹

¹⁹⁰ See Valente et al. (2020).

¹⁹¹ See Brinda & Fedosov (2023).

¹⁹² See Buck (1998).

¹⁹³ See Zander (1993).

¹⁹⁴ See Mahévas et al. (2016).

¹⁹⁵ See Mahévas et al. (2016).

¹⁹⁶ See Zander (2007d).

¹⁹⁷ See Ros et al. (2005).

¹⁹⁸ See Mahévas et al. (2016).

- Mnium** Hedw., *nom. cons.* (MNIACEAE)
arizonicum J.J.Amann
blyttii Bruch & Schimp.
hornum Hedw.
lycopodioides Schwägr.
marginatum (Dicks.) P.Beauv.
spinosum (Voit) Schwägr.
spinulosum Bruch & Schimp.
stellare Hedw.
thomsonii Schimp.
- Molendoa** Lindb. (POTTIACEAE)
handelii (Schiffn.) Brinda & R.H.Zander²⁰⁰
hornschuchiana (Hook.) Lindb. ex Limpr.
sendtneriana (Bruch & Schimp.) Limpr.
- Myrinia** Schimp., *nom. cons.* (MYRINIACEAE)
pulvinata (Wahlenb.) Schimp.
- Myurella** Schimp. (PLAGIOTHECIACEAE)
julacea (Schwägr.) Schimp.
sibirica (Müll.Hal.) Reimers
tenerrima (Brid.) Lindb.
- Myuroclada** Besch. (BRACHYTHECIACEAE)
maximowiczii (G.G.Borshch.) Steere & W.B.Schofield
- Neckera** Hedw., *nom. cons.* (NECKERACEAE)
californica Hook. & Arn.²⁰¹
douglasii Hook.
pennata Hedw.
- Neckeropsis** Reichardt (NECKERACEAE)
disticha (Hedw.) Kindb.
undulata (Hedw.) Reichardt
- Neodictyella** (Nog.) W.R.Buck (METERORIACEAE)
pendula (Sull.) W.R.Buck²⁰²
- Neomacounia** Ireland (NECKERACEAE)
nitida (Lindb.) Ireland
- Nogopterium** Crosby & W.R.Buck (LEUCODONTACEAE)
gracile (Hedw.) Crosby & W.R.Buck²⁰³
- Nyholmia** Holmen & E.Warncke²⁰⁴ (ORTHOTRICHACEAE)
gymnostoma (Bruch ex Brid.) Holmen & E.Warncke
obtusifolia (Brid.) Holmen & E.Warncke
- Octoblepharum** Hedw. (OCTOBLEPHARACEAE)
albidum Hedw.
- Oedipodium** Schwägr. (OEDIPODIACEAE)
griffithianum (Dicks.) Schwägr.

¹⁹⁹ See Shaw (2009).

²⁰⁰ See Brinda & Zander (2020).

²⁰¹ See Olsson et al. (2011).

²⁰² See Buck (1994).

²⁰³ See Crosby & Buck (2011).

²⁰⁴ See Sawicki et al. (2010).

Oligotrichum DC., *nom. cons.* (POLYTRICHACEAE)

- aligerum* Mitt.
falcatum Steere
hercynicum (Hedw.) DC.
parallellum (Mitt.) Kindb.

Oncophorus (Brid.) Brid. (DICRANACEAE)

- integerrimus* Hedenäs²⁰⁵
virens (Hedw.) Brid.

Oreas Brid., *nom. cons.* (DICRANACEAE)

- martiana* (Hoppe & Hornsch.) Brid.

Orthodontium Schwägr. (ORTHODONTIACEAE)

- gracile* (Wilson *ex* Hook.) Schwägr. *ex* Bruch & Schimp.
pellucens (Hook.) Bruch & Schimp.

Ortholimnobia Dixon (PLAGIOTHECIACEAE)

- handelii* (Broth.) Schröck & J.T.Wynns²⁰⁶

Orthothecium Schimp., *nom. cons.* (PLAGIOTHECIACEAE)

- acuminatum* Bryhn
chryseon (Schwägr.) Schimp.
intricatum (Hartm.) Schimp.
rufescens (Dicks. *ex* Brid.) Schimp.
strictum Lorentz

Orthotrichum Hedw. (ORTHOTRICHACEAE)

- alpestre* Hornsch. *ex* Bruch & Schimp.
anodon F.Lara, Garilleti & Mazimpaka²⁰⁷
anomalum Hedw.
bartramii R.S.Williams
consimile Mitt.
consobrinum Cardot²⁰⁸
cucullatum F.Lara, R.Medina & Garilleti²⁰⁹
cupulatum Hoffm. *ex* Brid.
cylindrocarpum Lesq.²¹⁰
diaphanum Schrad. *ex* Brid.
euryphyllum Venturi
flowersii Vitt
franciscanum F.Lara, R.Medina & Garilleti²¹¹
hallii Sull. & Lesq.
mazimpakanum Garilleti & F.Lara²¹²
norrisii F.Lara, R.Medina & Garilleti²¹³

²⁰⁵ See Afonina (2020).

²⁰⁶ See Wynns & Schröck (2018).

²⁰⁷ See Garilleti et al. (2006a).

²⁰⁸ See Flagmeier et al. (2021).

²⁰⁹ See Medina et al. (2013).

²¹⁰ See Medina et al. (2019b).

²¹¹ See Medina et al. (2013).

²¹² See Garilleti et al. (2011).

²¹³ See Medina et al. (2008).

- ohioense* Sull. & Lesq.
pallens Bruch *ex* Brid. var. *pallens*

var. *crenulatum* Vitt

var. *johnseniae* Vitt

pellucidum Lindb.

pulchellum Brunt.

pumilum Sw.

pusillum Mitt.

rivulare Turner

shevockii Lewinsky-Haapasaari & D.H.Norris²¹⁴

stellatum Brid.

stramineum Hornsch. *ex* Brid.

strangulatum P.Beauv.

underwoodii F.Lara, Garilleti & Mazimpaka²¹⁵

Oxyrrhynchium (Schimp.) Warnst., *nom. cons.*²¹⁶

(BRACHYTHECIACEAE)

hians (Hedw.) Loeske

Ozobryum G.L.Sm.Merr. (POTTIACEAE)

ogalalense G.L.Sm.Merr.²¹⁷

Palamocladium Müll.Hal. (BRACHYTHECIACEAE)

leskeoides (Hook.) E.Britton

Paludella Ehrh. *ex* Brid. (MEESIACEAE)

squarrosa (Hedw.) Brid.

Palustriella Ochyra (AMBLYSTEGIACEAE)

commutata (Hedw.) Ochyra

falcata (Brid.) Hedenäs

Paraleucobryum (Lindb. *ex* Limpr.) Péterfi (DICRANACEAE)

enerve (Thed.) Loeske

longifolium (Ehrh. *ex* Hedw.) Péterfi

sauteri (Bruch & Schimp.) Loeske²¹⁸

Pelekium Mitt., *nom. cons.*²¹⁹ (THUIDIACEAE)

involvens (Hedw.) Touw

minutulum (Hedw.) Touw

pygmaeum (Schimp.) Touw

schistocalyx (Müll.Hal.) Touw

Philonotis Brid. (BARTRAMIACEAE)

americana Dism.²²⁰

breuteloides Shevock & Aguero²²¹

capillaris Lindb.

cernua (Wilson) D.G.Griffin & W.R.Buck²²²

²¹⁴ See Lewinsky-Haapasaari (1998).

²¹⁵ See Garilleti et al. (2001).

²¹⁶ See Ignatov & Huttunen (2002).

²¹⁷ See Merrill (1992).

²¹⁸ See Haring (1961) and Allred et al. (2024).

²¹⁹ See Touw (2001).

²²⁰ See Koponen et al. (2012).

²²¹ See Shevock & Aguero (2023).

fontana (Hedw.) Brid.
longiseta (Michx.) E.Britton
marchica (Hedw.) Brid.
minuta (Taylor) A.Jaeger²²³
sphaericarpa (Hedw.) Brid.
tomentella Molendo
uncinata (Schwägr.) Brid.
yezoana Besch. & Cardot

Physcomitrium (Brid.) Brid. (FUNARIACEAE)

californicum E.Britton²²⁴
collenchymatum Gier
hookeri Hampe
immersum Sull.
patens (Hedw.) Mitt.²²⁵
pygmaeum James²²⁶
pyriforme (Hedw.) Brid.
readeri Müll.Hal.²²⁷
serratum (Wilson & Hook.) Müll.Hal.²²⁸

Pilosium (Müll.Hal.) M.Fleisch. (STEREOPHYLLACEAE)

chlorophyllum (Hornsch.) Müll.Hal.²²⁹

Pirella Cardot (PTEROBRYACEAE)

cymbifolia (Sull.) Cardot
pohlii (Schwägr.) Cardot

Plagiobryum Lindb. (BRYACEAE)

demissum (Hook.) Lindb.
zieri (Dicks. ex Hedw.) Lindb.

Plagiomnium T.J.Kop., *nom. cons.* (MNIACEAE)

ciliare (Müll.Hal.) T.J.Kop.
curvatulum (Lindb.) Schljakov²³⁰
cuspidatum (Hedw.) T.J.Kop.
drummondii (Bruch & Schimp.) T.J.Kop.
ellipticum (Brid.) T.J.Kop.
floridanum R.E.Wyatt & Odrzyk.²³¹
insigne (Mitt.) T.J.Kop.
medium (Bruch & Schimp.) T.J.Kop.
rhynchophorum (Harv.) T.J.Kop.²³²

²²² See Griffin & Buck (1989).

²²³ See Jiménez & Toren (2018).

²²⁴ See Norris & Shevock (2004a,b).

²²⁵ See Medina et al. (2019a).

²²⁶ See Medina et al. (2022).

²²⁷ See Medina et al. (2019a).

²²⁸ See Medina et al. (2019a).

²²⁹ Newly reported from North America based on a collection, *A.J. Shaw 8153*, on 14 Sep. 1996, from North Carolina (DUKE, NY).

²³⁰ See Wyatt et al. (1993).

²³¹ See Wyatt & Odrzykoski (2012).

²³² See Wyatt et al. (2021a).

rostratum (Schrad.) T.J.Kop.
undulatum (Hedw.) T.J.Kop.²³³
venustum (Mitt.) T.J.Kop.

Plagiopus Brid. (BARTRAMIACEAE)

oederianus (Sw.) H.A.Crum & L.E.Anderson
serratus Brid.²³⁴

Plagiothecium Schimp. (PLAGIOTHECIACEAE)

berggrenianum Frisvoll
cavifolium (Brid.) Z.Iwats.
curvifolium Schleiph. ex Limpr.²³⁵
denticulatum (Hedw.) Schimp. var. *denticulatum*
 var. *obtusifolium* (Turner) Moore²³⁶
laetum Schimp.
latebricola Schimp.
nemorale (Mitt.) A.Jaeger²³⁷
pacificum J.T.Wynns²³⁸
schofieldii Wolski & W.R.Buck²³⁹
succulentum (Wilson) Lindb.²⁴⁰
sylvaticum (Brid.) Schimp.²⁴¹
talbotii Wolski & W.R.Buck²⁴²
undulatum (Hedw.) Schimp.

Platydictya Berk. (PLAGIOTHECIACEAE)

jungermannioides (Brid.) H.A.Crum

Platygyrium Schimp., *nom. cons.* (PYLAISIADELPHACEAE)

fuscoluteum Cardot
repens (Brid.) Schimp.

Platyhypnum Loeske²⁴³ (AMBLYSTEGIACEAE)

alpestre (Hedw.) Ochyra
alpinum (Lindb.) Loeske
bestii (Renauld & Bryhn) Ochyra
cochlearifolium (Venturi) Ochyra
duriusculum (De Not.) Ochyra
molle (Hedw.) Loeske
norvegicum (Schimp.) Ochyra
smithii (Sw.) Ochyra

Platylomella A.L.Andrews (AMBLYSTEGIACEAE)

lescurii (Sull.) A.L.Andrews

²³³ See Joya & McIntosh (2012).

²³⁴ See Habeeb (1950) and Hedenäs (2020) as *P. alpinus*.

²³⁵ See Wolski et al. (2022c).

²³⁶ See Wolski et al. (2022a).

²³⁷ See Wolski et al. (2020).

²³⁸ See Wynns et al. (2018).

²³⁹ See Wolski et al. (2021).

²⁴⁰ See Wolski (2020).

²⁴¹ See Wolski et al. (2024).

²⁴² See Wolski et al. (2022b).

²⁴³ We follow the treatment by Ochyra (2013), supported by Kučera et al. (2019).

- Plaubelia** Brid.²⁴⁴ (POTTIACEAE)
sprengelii (Schwägr.) R.H.Zander
stomatodonta (Cardot) R.H.Zander
- Plenogenma** Plášek, Sawicki & Ochyra (ORTHOTRICHACEAE)
phyllantha (Brid.) Sawicki, Plášek & Ochyra²⁴⁵
- Pleuridium** Rabenh., *nom. cons.* (DITRICHACEAE)
acuminatum Lindb.
mexicanum Cardot²⁴⁶
ravenelii Austin
subulatum (Hedw.) Rabenh.
sullivantii Austin
- Pleurochaete** Lindb. (POTTIACEAE)
luteola (Besch.) Thér.²⁴⁷
- Pleuroziopsis** Kindb. ex E.Britton (CLIMACIACEAE)
ruthenica (Weinm.) Kindb. ex E.Britton
- Pleurozium** Mitt., *nom. cons.* (HYLOCOMIACEAE)
schreberi (Willd. ex Brid.) Mitt.
- Pogonatum** P.Beauv. (POLYTRICHACEAE)
brachyphyllum (Michx.) P.Beauv.
contortum (Menzies ex Brid.) Lesq.
dentatum (Menzies ex Brid.) Brid.
pensilvanicum (Bartram ex Hedw.) Paris
urnigerum (Hedw.) P.Beauv.
- Pohlia** Hedw. (MIELICHHOFERACEAE)
andalusica (Höhn.) Broth.
andrewsii A.J.Shaw
annotina (Hedw.) Lindb.
atropurpurea (Wahlenb.) H.Lindb.
beringiensis A.J.Shaw
bolanderi (Lesq.) Broth.
brevinervis Lindb. & Arnell
bulbifera (Warnst.) Warnst.
camptotrachela (Renauld & Cardot) Broth.
cardotii (Renauld) Broth.
columbica (Kindb.) A.L.Andrews
cruda (Hedw.) Lindb.
crudoides (Sull. & Lesq.) Broth.
drummondii (Müll.Hal.) A.L.Andrews
elongata Hedw.
erecta Lindb.
filum (Schimp.) Mårtensson
flexuosa Harv.²⁴⁸
greenii Brid.²⁴⁹
- lescuriana* (Sull.) Grout
longibracteata Broth.
longicolla (Hedw.) Lindb.
ludwigii (Spreng. ex Schwägr.) Broth.
melanodon (Brid.) A.J.Shaw
nutans (Hedw.) Lindb.
obtusifolia (Vill. ex Brid.) L.F.Koch
pacifica A.J.Shaw
proliger (Kindb. ex Breidl.) Lindb. ex Arnell
rabunbaldensis A.J.Shaw²⁵⁰
robertsonii Shevock & A.J.Shaw²⁵¹
sphagnicola (Bruch & Schimp.) Broth.²⁵²
tundrae A.J.Shaw
vexans (Limpr.) H.Lindb.
wahlenbergii (F.Weber & D.Mohr) A.L.Andrews
- Polytrichastrum** G.L.Sm.²⁵³ (POLYTRICHACEAE)
alpinum (Hedw.) G.L.Sm.
fragile (Bryhn) Schljakov
lyallii (Mitt.) G.L. Sm
papillatum G.L.Sm.
septentrionale (Sw. ex Brid.) E.I.Ivanova, N.E.
Bell & Ignatov
sexangulare (Flörke ex Brid.) G.L.Sm.
sphaerothecium (Besch.) J.-P.Frahm
- Polytrichum** Hedw.²⁵⁴ (POLYTRICHACEAE)
appalachianum L.E.Anderson
commune Hedw.
densifolium Wilson ex Mitt.²⁵⁵
formosum Hedw.
hyperboreum R.Br.
jensenii I.Hagen
juniperinum Hedw.
longisetum Sw. ex Brid.
ohioense Renauld & Cardot
pallidisetum Funck
perigoniale Michx.²⁵⁶
piliferum Hedw.
strictum Menzies ex Brid.
swartzii Hartm.

²⁴⁴ See Zander (2021).²⁴⁵ See Plášek et al. (2015).²⁴⁶ See Yip et al. (2007).²⁴⁷ See Eckel (2007) and Wyatt & Stoneburner (2018).²⁴⁸ See Shaw & Toren (2009).²⁴⁹ See Guerra (2021).²⁵⁰ See Shevock & Shaw (2005).²⁵¹ See Shevock & Shaw (2005).²⁵² Considered conspecific with *P. nutans* by Shaw (2014) but retained as distinct by Hodgetts et al. (2020).²⁵³ See Bell & Hyvönen (2010).²⁵⁴ See Bell & Hyvönen (2010).²⁵⁵ See Ivanova & Ignatov (2017).²⁵⁶ Considered a variety of *P. commune* by Merrill (2007) but a distinct species by Kariyawasam (2021).

- Pseudanomodon*** (Limpr.) Ignatov & Fedosov²⁵⁷
(NECKERACEAE)
attenuatus (Hedw.) Ignatov & Fedosov
- Pseudephemerum*** (Lindb.) I.Hagen (DITRICHACEAE)
nitidum (Hedw.) Loeske²⁵⁸
- Pseudisothecium*** Grout²⁵⁹ (LEMBOPHYLLACEAE)
cardotii (Kindb.) Ignatova, Fedosov & Ignatov
crisatum (Hampe) Ignatova, Fedosov & Ignatov
myosuroides (Brid.) Grout
stoloniferum (Brid.) Grout
- Pseudoamblystegium*** Vanderp. & Hedenäs
(AMBLYSTEGIACEAE)
subtile (Hedw.) Vanderp. & Hedenäs²⁶⁰
- Pseudobraunia*** (Lesq. & James) Broth. (HEDWIGIACEAE)
californica (Lesq.) Broth.
- Pseudobryum*** (Kindb.) T.J.Kop. (MNIACEAE)
cinclidioides (Huebener) T.J.Kop.
- Pseudocampylium*** Vanderp. & Hedenäs²⁶¹
(AMBLYSTEGIACEAE)
radicale (P.Beauv.) Vanderp. & Hedenäs
- Pseudocrossidium*** R.S.Williams (POTTIACEAE)
arenicola (Dusén) M.J.Cano²⁶²
hornschuchianum (Schultz) R.H.Zander
obtusulum (Lindb.) H.A.Crum & L.E.Anderson
replicatum (Taylor) R.H.Zander
- Pseudocryphaea*** E.Britton ex Broth. (RUTENBERGIACEAE)
domingensis (Spreng.) W.R.Buck
- Pseudoditrichum*** Steere & Z.Iwats. (PSEUDODITRICHACEAE)
mirabile Steere & Z.Iwats.
- Pseudohygrohypnum*** Kanda²⁶³ (PYLAIACEAE)
appalachianum Brinda, Fedosov & Ignatova²⁶⁴
eugyrium (Schimp.) Kanda
fauriei (Cardot) Jan Kučera & Ignatov
fertile (Sendtn.) Jan Kučera & Ignatov
neglectum Fedosova & Ignatova
subeugyrium (Renauld & Cardot) Ignatov & Ignatova
- Pseudoleskeella*** Kindb. (LESKEACEAE)
nervosa (Brid.) Nyholm

²⁵⁷ See Ignatov et al. (2019a).

²⁵⁸ See Yip (2002).

²⁵⁹ See Ignatova et al. (2019).

²⁶⁰ See Vanderpoorten & Hedenäs (2009).

²⁶¹ See Vanderpoorten & Hedenäs (2009).

²⁶² See Cano et al. (2016).

²⁶³ See Fedosov et al. (2022).

²⁶⁴ See Fedosov et al. (2022).

rupestris (Berggr.) Hedenäs & L.Söderstr.²⁶⁵
serpentinensis P.S.Wilson & D.H.Norris
tectorum (Funck ex Brid.) Kindb.

- Pseudoscleropodium*** (Limpr.) M.Fleisch.
(BRACHYTHECIACEAE)
purum (Hedw.) M.Fleisch.
- Pseudostereodon*** (Broth.) M.Fleisch.²⁶⁶ (PYLAIACEAE)
procerrimus (Molendo) M.Fleisch.
- Pseudotaxiphyllum*** Z.Iwats. (PLAGIOTHECIACEAE)
elegans (Brid.) Z.Iwats.
subfalcatum (Austin) X.Q.Li, Q.Zuo & Y.F.Wang²⁶⁷
- Psilopilum*** Brid. (POLYTRICHACEAE)
cavifolium (Wilson) I.Hagen
laevigatum (Wahlenb.) Lindb.
- Pterigynandrum*** Hedw. (PTERIGYNANDRACEAE)
filiforme Hedw.
- Pterygoneurum*** Jur., *nom. cons.* (POTTIACEAE)
×kieneri (Habeeb) W.R.Buck & Goffinet²⁶⁸
×kozlovii Laz.²⁶⁹
lamellatum (Lindb.) Jur.
ovatum (Hedw.) Dixon
subsessile (Brid.) Jur.
- Ptilium*** De Not. (PYLAIACEAE)
crista-castrensis (Hedw.) De Not.
- Ptychomitrium*** Fürnr., *nom. cons.* (PTYCHOMITRIACEAE)
drummondii (Wilson) Sull.
gardneri Lesq.
incurvum (Schwägr.) Spruce
serratum Bruch & Schimp.
sinense (Mitt.) A.Jaeger
- Pulviger*** Plášek, Sawicki & Ochrya²⁷⁰ (ORTHOTRICHACEAE)
howei (Renauld & Cardot) F.Lara, Draper & Garilleti²⁷¹

²⁶⁵ See Hedenäs & Söderström (1991).

²⁶⁶ See Câmara et al. (2018).

²⁶⁷ See Li et al. (2015).

²⁶⁸ *Pterygoneurum* *×kieneri* (Habeeb) W.R.Buck & Goffinet comb. et stat. nov. \equiv *P. subsessile* var. *kieneri* Habeeb, *Rhodora* 51: 128. 1949. TYPE: U.S.A., Nebraska, Chase Co.: 8 mi W of Champion, 31 Jul. 1941, W. Kiener 10627 p.p. (NEB). See Guerra et al. (1994), who demonstrated that this taxon is a hybrid between *P. ovatum* and *Phascum cuspidatum*. Recognizing this taxon at the species level is treating it like the intergeneric hybrid *Entosthodon* *×hungaricus* (Boros) Loeske (Beike et al. 2014; Patel et al. 2023). This may be the same taxon as *P. ×kozlovii*, but is recognized until additional data are available.

²⁶⁹ See Guerra et al. (1994) for demonstration of the hybrid nature of this taxon.

²⁷⁰ See Plášek et al. (2015).

- lyellii* (Hook. & Taylor) Plášek, Sawicki & Ochyra
papillosa (Hampe) F.Lara, Draper & Garilleti²⁷²
pringlei (Müll.Hal.) F.Lara, Draper & Garilleti²⁷³
Pylaisiadelphina Cardot²⁷⁴ (PYLAISIADELPHACEAE)
canadensis (W.B.Schofield) W.R.Buck & Goffinet²⁷⁵
henonii (Duby) W.R.Buck
recurvans (Michx.) W.R.Buck
roellii (Renauld & Cardot) W.R.Buck
tenuirostris (Bruch & Schimp. ex Sull.) W.R.Buck
Pylaisia Schimp., *nom. cons.* (PYLAISIAACEAE)
intricata (Hedw.) Schimp.
polyantha (Hedw.) Schimp.
selwynii Kindb.²⁷⁶
steerei (Ando & Higuchi) Ignatov
Pyramidula Brid. (FUNARIAEAE)
tetragona (Brid.) Brid.
Pyrrhobryum Mitt. (RHIZOGONIAEAE)
spiniforme (Hedw.) Mitt.
Racomitrium Brid. (GRIMMIAEAE)
aciculare (Hedw.) Brid.
aduncooides Bedn.-Ochyra²⁷⁷
afoninae Frisvoll²⁷⁸
affine (Schleich. ex F.Weber & D.Mohr) Lindb.
brevipes Kindb.
brevisetum Lindb.²⁷⁹
canescens (Hedw.) Brid. subsp. *canescens*
subsp. *latifolium* (C.E.O.Jensen) Frisvoll
corrugatum (Bedn.-Ochyra) Larraín & J.Muñoz²⁸⁰
depressum Lesq.
elongatum Ehrh. ex Frisvoll
ericoides (Brid.) Brid.

²⁷¹ See Lara et al. (2020).

²⁷² See Lara et al. (2020).

²⁷³ See Lara et al. (2020).

²⁷⁴ See Buck (1984).

²⁷⁵ Described as *Brotherella canadensis* by Schofield (2006) and accommodated in *Pylaisiadelphina* under the new combination *Pylaisiadelphina canadensis* (W.B.Schofield) W.R.Buck & Goffinet comb. nov. ≡ *Brotherella canadensis* W.B.Schofield, J. Hattori Bot. Lab. 100: 355. 2006. TYPE: Canada, British Columbia: just NE of Webster Island, Effingham Inlet, Barclay Cound, Vancouver Island, 49°05'N, 125°09'W, W.B. Schofield 43474 (holotype, UBC).

²⁷⁶ See Ignatova et al. (2020b).

²⁷⁷ See Bednarek-Ochyra (1999).

²⁷⁸ See Ochyra & Bednarek-Ochyra (2007) under *Bucklandiella*.

²⁷⁹ Currently only known from one unverified collection, *Lyford* (MSC-B-0008248) from Oregon.

²⁸⁰ Described by Bednarek-Ochyra (2004) as *Codriophorus*.

- fasciculare* (Schrad. ex Hedw.) Brid.
heterostichum (Hedw.) Brid.
lanuginosum (Hedw.) Brid.
lawtoniae Ireland
macounii Kindb. subsp. *macounii*
subsp. *alpinum* (E.Lawton) Frisvoll
microcarpon (Hedw.) Brid. fo. *microcarpon*
fo. *afoninae* Frisvoll²⁸¹
molle Cardot²⁸²
muticum (Kindb.) Frisvoll
norrisii Bedn.-Ochyra & Ochyra²⁸³
obesum Frisvoll
occidentale (Renauld & Cardot) Renauld & Cardot
pacificum Ireland & J.R.Spence
panschii (Müll.Hal.) Kindb.
pygmaeum Frisvoll
ryszardii Bedn.-Ochyra²⁸⁴
sudeticum (Funck) Bruch & Schimp.
varium (Mitt.) A.Jaeger
venustum Frisvoll
Racopilum P.Beauv. (RACOPILACEAE)
tomentosum (Hedw.) Brid.
Rauarella Reimers (THUIDIACEAE)
praelonga (Schimp. ex Besch.) Wijk & Margad.
scita (P.Beauv.) Reimers
Rectithecium Hedenäs & Huttunen²⁸⁵ (PLAGIOTHECIAEAE)
piliferum (Sw.) Hedenäs & Huttunen
Redfearnia J.T.Wynns²⁸⁶ (PLAGIOTHECIAEAE)
homomallifolia (Redf.) J.T.Wynns
Rhabdoweisia Bruch & Schimp. (RHABDOWEISIAEAE)
crenulata (Mitt.) H.Jameson
crispata (Dicks.) Lindb.
Rhachithecium Broth. ex Le Jol. (RHACHITHECIAEAE)
perpusillum (Thwaites & Mitt.) Broth.
Rhexophyllum Herzog (POTTIACEAE)
subnigrum (Mitt.) Hilp.
Rhizogemma Bonfim Santos, Siebel & Fedosov
(RHIZOGEMMACEAE)
staphylina (H.Whitehouse) Bonfim Santos, Siebel
& Fedosov²⁸⁷

²⁸¹ See Ochyra in Ellis et al. (2018).

²⁸² See Bednarek-Ochyra & Ochyra (2007) under *Codriophorus*.

²⁸³ See Bednarek-Ochyra & Ochyra (2000).

²⁸⁴ See Bednarek-Ochyra (2000).

²⁸⁵ See Huttunen et al. (2013).

²⁸⁶ See Wynns (2020).

²⁸⁷ See Whitehouse (2001).

Rhizomnium T.J.Kop. (MNIACEAE)

- andrewsianum* (Steere) T.J.Kop.
appalachianum T.J.Kop.
chlorophyllosum (Kindb.) T.J.Kop.²⁸⁸
glabrescens (Kindb.) T.J.Kop.
gracile T.J.Kop.
magnifolium (Horik.) T.J.Kop.
nudum (R.S.Williams) T.J.Kop.
pseudopunctatum (Bruch & Schimp.) T.J.Kop.
punctatum (Hedw.) T.J.Kop.

Rhodobryum (Schimp.) Limpr., *nom. cons.* (BRYACEAE)

- ontariense* (Kindb.) Paris
roseum (Hedw.) Limpr.

Rhynchostegium Schimp., *nom. cons.* (BRACHYTHECIACEAE)

- aquaticum* Spruce²⁸⁹
serrulatum (Hedw.) Austin

Rhytidiadelphus (Limpr.) Warnst. (HYLOCOMIACEAE)

- japonicus* (Reimers) T.J.Kop.
loreus (Hedw.) Warnst.
squarrosus (Hedw.) Warnst.
subpinnatus (Lindb.) T.J.Kop.

Rhytidiopsis Broth. (HYLOCOMIACEAE)

- robusta* (Hook.) Broth.

Rhytidium (Sull.) Austin ex C.F.Parker (RHYTIDIACEAE)

- rugosum* (Hedw.) Austin ex C.F.Parker

Roaldia P.E.A.S.Câmara & Carv.-Silva²⁹⁰ (PYLAISIACEAE)

- revoluta* (Mitt.) P.E.A.S.Câmara & Carv.-Silva

Roellobryon Ochyra²⁹¹ (ROELLOBRYACEAE)

- roellii* (Broth.) Ochyra

Ruficaulis Bonfim Santos & Fedosov²⁹² (RUFICAULACEAE)

- rufescens* (Dicks.) Bonfim Santos & Fedosov

Saelania Lindb. (SAELANIACEAE)

- glaucescens* (Hedw.) Broth.

Sanionia Loeske (AMBLYSTEGIACEAE)

- nivalis* Hedenäs
orthothecioides (Lindb.) Loeske
symmetrica (Renauld & Cardot) Wheldon
uncinata (Hedw.) Loeske

Schistidium Bruch & Schimp., *nom. cons.* (GRIMMIACEAE)

- agassizii* Sull. & Lesq.
ambiguum Sull.²⁹³
andreaeopsis (Müll.Hal.) Laz.

apocarpum (Hedw.) Bruch & Schimp.

atrichum (Müll.Hal. & Kindb.) W.A.Weber

atrofuscum (Schimp.) Limpr.

boreale Poelt

canadense (Dupr.) Ignatova & H.H.Blom

cinclidodonteum (Müll.Hal.) B.Bremer

confertum (Funck) Bruch & Schimp.

crassipilum H.H.Blom

crassithecium H.H.Blom ex B.H.Allen²⁹⁴

cryptocarpum Mogensen & H.H.Blom

dupretii (Thér.) W.A.Weber

echinatum Ignatova & H.H. Blom²⁹⁵

flaccidum (De Not.) Ochyra

flexipile (Lindb. ex Broth.) G.Roth

frigidum H.H.Blom

frisvollianum H.H.Blom

grandirete H.H.Blom

heterophyllum (Kindb.) T.T.McIntosh

holmenianum Steere & Brassard

lancifolium (Kindb.) H.H.Blom²⁹⁶

liliputanum (Müll.Hal.) Deguchi

maritimum (Turner ex R.R.Scott) Bruch & Schimp.

occidentale (E.Lawton) S.P.Churchill

papillosum Culm.

poeltii H.H.Blom

pulchrum H.H.Blom

relictum T.T.McIntosh, H.H.Blom & Ignatova²⁹⁷

rivulare (Brid.) Podp.

robustum (Nees & Hornsch.) H.H.Blom

sinensiapocarpum (Müll.Hal.) Ochyra²⁹⁸

splendens T.T.McIntosh, H.H.Blom, D.R.Toren
& Shevock²⁹⁹

squarrosum T.T.McIntosh, H.H.Blom, D.R.
Toren & Shevock³⁰⁰

strictum (Turner) Loeske ex Mårtensson

subjulaceum H.H.Blom

tenerum (J.E.Zetterst.) Nyholm

teretinerve (Limpr.) Limpr.³⁰¹

trichodon (Brid.) Poelt

venetum H.H.Blom

²⁸⁸ See Koponen & Sun (2017) and Wyatt et al. (2023).

²⁸⁹ See Wynns (2006) and Huttunen & Ignatov (2010).

²⁹⁰ See Câmara et al. (2018).

²⁹¹ See Ochyra (2011).

²⁹² See Fedosov et al. (2023b).

²⁹³ See Sullivant (1849) and Allred et al. (2024).

²⁹⁴ See Allen (2005).

²⁹⁵ See Ignatova et al. (2009).

²⁹⁶ Treated as conspecific with *Schistidium apocarpum* by McIntosh (2007) but here resurrected following Ignatova & Blom (2017).

²⁹⁷ See McIntosh et al. (2017).

²⁹⁸ See Allred et al. (2024).

²⁹⁹ See McIntosh et al. (2015).

³⁰⁰ See McIntosh et al. (2015).

³⁰¹ See Kiebacher & Meier (2021).

Schistostega D.Mohr (SCHISTOSTEGACEAE)
pennata (Hedw.) F.Weber & D.Mohr

Schlotheimia Brid. (ORTHOTRICHACEAE)

lancifolia E.B.Bartram

rugifolia (Hook.) Schwägr.

Schoenobryum Dozy & Molk. (CRYPHAEACEAE)

concauifolium (Griff.) Gangulee³⁰²

Schwetschkeopsis Broth. (ANOMODONTACEAE)

fabronia (Schwägr.) Broth.

Sciuro-hypnum (Hampe) Hampe³⁰³ (BRACHYTHECIACEAE)

curtum (Lindb.) Ignatov

glaciale (Schimp.) Ignatov & Huttunen

hylotapetum (N.L.Higinb. & B.L.Higinb.) Ignatov

& Huttunen

latifolium (Kindb.) Ignatov & Huttunen

oedipodium (Mitt.) Ignatov & Huttunen

ornellanum (Molendo) Ignatov & Huttunen

plumosum (Hedw.) Ignatov & Huttunen

populeum (Hedw.) Ignatov & Huttunen

reflexum (Starke) Ignatov & Huttunen

starkei (Brid.) Ignatov & Huttunen

uncinifolium (Broth. & Paris) Ochyra & Żarnowiec

Scleropodium Schimp. (BRACHYTHECIACEAE)

californicum (Lesq.) Kindb.

cespitans (Müll.Hal.) L.F.Koch

julaceum E.Lawton

obtusifolium (Mitt.) Kindb.

occidentale B.E.Carter³⁰⁴

touretii (Brid.) L.F.Koch

Scopelophila (Mitt.) Lindb. (POTTIACEAE)

ligulata (Spruce) Spruce³⁰⁵

Scorpidium (Schimp.) Limpr. (CALLIERGONACEAE)

cossonii (Schimp.) Hedenäs

revolvens (Sw.) Rubers

scorpioides (Hedw.) Limpr.

Scouleria Hook. (SCOULERIACEAE)

aquatica Hook.

marginata E.Britton

siskiyouensis Shevock & D.H.Norris³⁰⁶

Seligeria Bruch. & Schimp. (SELIGERIACEAE)

acutifolia Lindb.

brevifolia (Lindb.) Lindb. & Arnell

calcareo (Hedw.) Bruch & Schimp.

careyana Vitt & W.B.Schofield

donniana (Sm.) Müll.Hal.

oelandica C.E.O.Jensen & Medelius

pusilla (Hedw.) Bruch & Schimp.

tristichoides Kindb.

Sematophyllum Mitt. (SEMATOPHYLLACEAE)

adnatum (Michx.) E.Britton

demissum (Wilson) Mitt.

marylandicum (Müll.Hal.) E.Britton

Serpoleskea (Hampe ex Limpr.) Loeske³⁰⁷

(PYLAIADIACEAE)

confervoides (Brid.) Loeske

minutissima (Sull. & Lesq.) W.R.Buck & B.H.Allen³⁰⁸

Solmsiella Müll.Hal. (ERPODIACEAE)

biseriata (Austin) Steere³⁰⁹

Sphagnum L. (SPHAGNACEAE)

affine Renauld & Cardot

alaskense R.E.Andrus & Janssens

angermanicum Melin

angustifolium (C.E.O.Jensen ex Russow) C.E.O.Jensen

annulatum H.Lindb. ex Warnst.

aongstroemii C.Hartm.

arcticum Flatberg & Frisvoll

austinii Sull.

balticum (Russow) C.E.O.Jensen

beothuk R.E.Andrus

capillifolium (Ehrh.) Hedw.

carolinianum R.E.Andrus

centrale C.E.O.Jensen

compactum DC.

concinnum (Berggr.) Flatberg

contortum Schultz

cribrosum Lindb.

cuspidatum Ehrh. ex Hoffm.

cyclophyllum Sull. & Lesq.

diabolicum A.J.Shaw, Aguero & Nieto-Lugilde

divinum Flatberg & K.Hassel

fallax (H.Klinggr.) H.Klinggr.

fimbriatum Wilson

fitzgeraldii Renauld ex Lesq.

³⁰² See Buck (1994).

³⁰³ See Ignatov & Huttunen (2002).

³⁰⁴ See Carter (2012).

³⁰⁵ Kuznetsova et al. (2023) recognized the genus *Merceya* Schimp. to accommodate *Merceya ligulata* (Spruce)

Schimp., but this generic name is superfluous for the earlier *Scopelophila* (Mitt.) Lindb. (Brinda & Fedosov 2023).

³⁰⁶ See Shevock & Norris (2014).

³⁰⁷ See Vanderpoorten & Hedenäs (2009).

³⁰⁸ See Buck & Allen (2017).

³⁰⁹ See Pursell (2017).

flavicomans (Cardot) Warnst.
flexuosum Dozy & Molk.
fuscum (Schimp.) H.Klinggr.
girgensohnii Russow
imbricatum Hornsch. ex Russow
incundum Flatberg & K.Hassel
inexpectatum Flatberg
jensenii H.Lindb.
junghuhnianum Dozy & Molk.
kenaiense R.E.Andrus
lenense H.Lindb.
lescurii Sull.
lindbergii Schimp.
macrophyllum Bernh. ex Brid.
magniae A.J.Shaw, Aguero & Nieto-Lugilde
majus (Russow) C.E.O.Jensen
medium Limpr.
mendocinum Sull. & Lesq.
miyabeanum Warnst.³¹⁰
microcarpum Warnst.
mirum Flatberg & Thinggaard
mississippiense R.E.Andrus
missouricum Warnst. & Cardot³¹¹
molle Sull.
obtusum Warnst.
oregonense R.E.Andrus
orientale L.I.Savicz
pacificum Flatberg
palustre L.
papillosum Lindb.
perfoliatum L.I.Savicz
perichaetiale Hampe
platyphyllum (Lindb. ex Braithw.) Sull. ex Warnst.
portoricense Hampe
pulchrum (Lindb. ex Braithw.) Warnst.
pylaesii Brid.
quinquefarium (Lindb. ex Braithw.) Warnst.
recurvum P.Beauv.
riparium Ångstr.
rubellum Wilson
rubiginosum Flatberg
russowii Warnst.
splendens Maass
squarrosum Crome
steerei R.E.Andrus

strictum Sull.
subfulvum Sjörs
subnitens Russow & Warnst.
subsecundum Nees
talbotianum R.E.Andrus
tenellum (Brid.) Bory
tenerum Sull. & Lesq. ex Sull.
teres (Schimp.) Ångstr.
torreyanum Sull.
trinitense Müll.Hal.
tundrae Flatberg
warnstorffii Russow
wulfianum Girg.

Splachnobryum Müll.Hal. (POTTIACEAE)

obtusum (Brid.) Müll.Hal.

Splachnum Hedw. (SPLACHNACEAE)

ampullaceum Hedw.

luteum Hedw.

melanocaulon (Wahlenb.) Schwägr.³¹²

pennsylvanicum (Brid.) Grout ex H.A.Crum

rubrum Hedw.

sphaericum Hedw.

vasculosum Hedw.

Stegonia Venturi (POTTIACEAE)

hyalinotricha (Cardot & Thér.) R.H.Zander

latifolia (Schwägr.) Venturi ex Broth.

pilifera (Brid.) H.A.Crum & L.E.Anderson

Stereodon (Brid.) Brid.³¹³ (STEREODONTACEAE)

callichrous (Brid.) Brid.

hamulosus (Schimp.) Lindb.

holmenii (Ando) Ignatov & Ignatova³¹⁴

pratensis (W.D.J.Koch ex Spruce) E.Britton

subimponens (Lesq.) Broth.

Stereophyllum Mitt. (STEREOPHYLLACEAE)

radiculosum (Hook.) Spruce

Streblotrichum P.Beauv.³¹⁵ (POTTIACEAE)

convolutum (Hedw.) P.Beauv. var. *convolutum*

var. *eustegium* (Cardot & Thér.) W.R.Buck & Goffinet³¹⁶

³¹⁰ See Shaw et al. (2015).

³¹¹ See Shaw et al. (2012b).

³¹² See Worley & Iwatzuki (1970).

³¹³ See Kučera et al. (2019).

³¹⁴ Described by Ando as *Hypnum holmenii* (Ando 1994) and transferred to *Stereodon* by Ignatov & Ignatova (2004).

³¹⁵ See Kučera et al. (2013).

³¹⁶ Recognized by Zander (2007a) under *Barbula convoluta* and accommodated here in *Streblotrichum* under the new combination *Streblotrichum convolutum* var. *eustegium* (Cardot & Thér.) W.R.Buck & Goffinet comb. nov. ≡ *Barbula eustegia* Cardot & Thér., Bot. Gaz. 30: 17. 1900. TYPE:

var. *gallinula* (R.H.Zander) W.R.Buck & Goffinet³¹⁷

Symblepharis Mont. (RHABDOWEISACEAE)

raui (Austin) R.S.Williams ex Broth.³¹⁸

vaginata (Hook. ex Harv.) Wijk & Margad.

Syntrichia Brid.³¹⁹ (POTTIACEAE)

ammonsiana (H.A.Crum & L.E.Anderson) Ochyra

amphidiacea (Müll.Hal.) R.H.Zander³²⁰

bartramii (Steere) R.H.Zander

caninervis Mitt.

cascadensis W.A.Kramer

chisosa (Magill, Delgad. & L.R.Stark) R.H.Zander

coloradensis W.A.Kramer

fragilis (Taylor) Ochyra

hermannii W.A.Kramer

laevipila Brid.

laeviuscula (Kindb.) W.A.Kramer

latifolia (Bruch ex Hartm.) Huebener

leptotricha (Müll.Hal. & Kindb.) W.A.Kramer

lithophila (Dusén) Ochyra & R.H.Zander³²¹

norvegica F.Weber var. *norvegica*

var. *cainii* (H.A.Crum & L.E.Anderson) W.A. Kramer

novomexicana W.A.Kramer

obtusissima (Müll.Hal.) R.H.Zander subsp. *obtusissima*

subsp. *stellaris* W.A.Kramer

subsp. *substellaris* W.A.Kramer

U.S.A., Idaho, Latah Co.: Cedar Creek, 1897, L. F. Henderson (holotype, P).

³¹⁷ Recognized by Zander (2007a) under *Barbula convoluta* and accommodated here in *Streblotrichum* under the new combination *Streblotrichum convolutum* var. *gallinula* (R.H.Zander) W.R.Buck & Goffinet comb. nov. \equiv *Barbula convoluta* Hedw. var. *gallinula* R.H.Zander, *Phytologia* 44(4): 195. 1979. TYPE: Canada, Northwest Territories, Distr. Mackenzie: Nananni National Park, South Nahanni River, Virginia Falls, 61°38'N, 125°42'W, 3 Sep. 1974, G. Scotter 22433 (holotype, NY; isotype, BUF).

³¹⁸ See Fedosov et al. (2021).

³¹⁹ The genus was typified by Zander (1989). We follow the treatment by Kramer (2023).

³²⁰ The species was resolved as sister to a species of *Streptopogon* Wilson ex Mitt. in Jáuregui-Lazo et al. (2023) consistent with inferences by Gallego et al. (2022). However, in both cases *Streptopogon* is nested within a larger *Syntrichia* clade. We therefore retain *amphidiacea* within *Syntrichia* rather than accepting its transfer to *Streptopogon*, pending further systematic resolution of the complex. However, Kramer (2023) argued for its exclusion from *Syntrichia* and for its recognition within *Tortula* on morphological grounds.

³²¹ See Brooks & Jáuregui-Lazo (2023).

ovata W.A.Kramer subsp. *dendropapillosa* W.A. Kramer

subsp. *subdendropapillosa* W.A.Kramer

pagorum (Milde) J.J.Amann

papillosa (Wilson ex Spruce) Spruce

princeps (De Not.) Mitt.

ruralis (Hedw.) F.Weber & D.Mohr subsp. *ruralis*

subsp. *hirsuta* (Venturi) Düll

subsp. *subpapillosissima* (Bizot & R.B.Pierrot

ex W.A.Kramer) W.A.Kramer

sinensis (Müll.Hal.) Ochyra

subintermedia (Renauld & Cardot) W.A.Kramer

var. *subintermedia*

var. *umatillae* W.A.Kramer

sucrosa Kellman³²²

virescens (De Not.) Ochyra subsp. *virescens*

subsp. *zailiiskialatauensis* W.A.Kramer

Syrrophodon Schwägr. (CALYMPERACEAE)

gaudichaudii Mont.

incompletus Schwägr. var. *incompletus*

ligulatus Mont.

parasiticus (Sw. ex Brid.) Besch.

prolifer Schwägr. var. *scaber* (Mitt.) W.D.Reese

texanus Sull.

Takakia S.Hatt. & Inoue³²³ (TAKAKIACEAE)

ceratophylla (Mitt.) Grolle

lepidozoides S.Hatt. & Inoue

Taxiphillum M.Fleisch. (TAXIPHYLLACEAE)

alternans (Cardot) Z.Iwats.

cuspidifolium (Cardot) Z.Iwats.

deplanatum (Bruch & Schimp. ex Sull.) M.Fleisch.

taxirameum (Mitt.) M.Fleisch.

Taxithelium Spruce ex Mitt. (PYLAIADIACEAE)

planum (Brid.) Mitt.

Tayloria Hook. (SPLACHNACEAE)

acuminata Hornsch.

froelichiana (Hedw.) Mitt. ex Broth.

hornschuchii (Grev. & Arn.) Broth.

lingulata (Dicks.) Lindb.

serrata (Hedw.) Bruch & Schimp.

splachnoides (Schleich. ex Schwägr.) Hook.

Tetraphis Hedw. (TETRAPHIDACEAE)

geniculata Girg. ex Milde

pellucida Hedw. var. *pellucida*

var. *trachypoda* (Kindb.) Harpel³²⁴

³²² See Kellman (2008).

³²³ See Smith & Davison (1993) and Newton et al. (2000).

³²⁴ See Harpel (2006).

Tetradontium Schwägr. (TETRAPHIDACEAE)*brownianum* (Dicks.) Schwägr.*ovatum* (Funck) Schwägr.*repandum* (Funck) Schwägr.**Thamnobryum** Nieuwl. (NECKERACEAE)*neckeroides* (Hook.) E.Lawton*subserratum* (Hook. ex Harv.) Nog. & Z.Iwats.³²⁵**Thelia** Sull. (THELIACEAE)*asprella* (Schimp.) Sull.*hirtella* (Hedw.) Sull.*lescurii* Sull.**Thuidium** Schimp. (THUIDIACEAE)*alleniorum* Austin³²⁶*assimile* (Mitt.) A.Jaeger*delicatulum* (Hedw.) Schimp.*recognitum* (Hedw.) Lindb.*tamariscinum* (Hedw.) Schimp.**Timmia** Hedw., *nom. cons.* (TIMMIACEAE)*austriaca* Hedw.*bavarica* Hessel.*comata* Lindb. & Arnell³²⁷*megapolitana* Hedw.*norvegica* J.E.Zetterst.*sibirica* Lindb. & Arnell**Timmiella** (DeNot.) Limpr. (TIMMIELLACEAE)*anomala* (Bruch & Schimp.) Limpr.*crassinervis* (Hampe) L.F.Koch**Tomentypnum** Loeske³²⁸ (AMBLYSTEGIACEAE)*falcifolium* (Renauld ex Nichols) Tuom.*involutum* (Limpr.) Hedenäs & Ignatov³²⁹*nitens* (Hedw.) Loeske**Tortella** (Müll.Hal.) Limpr., *nom. cons.* (POTTIACEAE)*alpicola* Dixon*arctica* (Arnell) Crundw. & Nyholm³³⁰× *cuspidatissima* (Cardot & Thér.) O.Werner,
Köckinger & Ros³³¹*densa* (Lorentz & Molendo) Crundw. & Nyholm³³²*flavovirens* (Bruch) Broth.*fragilis* (Drumm.) Limpr.*humilis* (Hedw.) Jenn.*inclinata* (R.Hedw.) Limpr.*nitida* (Lindb.) Broth. var. *fragilifolia* (Jur.)
Köckinger & Hedenäs³³³*rigens* Alberts.*spitsbergensis* (Bizot & Thér.) O.Werner, Köckinger
& Ros³³⁴*tortuosa* (Schrad. ex Hedw.) Limpr.**Tortula** Hedw., *nom. cons.* (POTTIACEAE)*acaulon* (With.) R.H.Zander*amplexa* (Lesq.) Steere*atrovirens* (Sm.) Lindb.*bolanderi* (Lesq.) M.Howe*brevipes* (Lesq.) Broth.*brevissima* Schiffn.³³⁵*californica* E.B.Bartram*caucasica* Lindb.*cernua* (Huebener) Lindb.*cuneifolia* (Dicks.) Turner var. *blissii* R.H.Zander*deciduidentata* (Sharp & Z.Iwats.) R.H.Zander*guepinii* (Bruch & Schimp.) Broth.*hoppeana* (Schultz) Ochyra*inermis* (Brid.) Mont.*israelis* Bizot & F.Bilewsky³³⁶*laureri* (Schultz) Lindb.*leucostoma* (R.Br.) Hook. & Grev.*lindbergii* Kindb. ex Broth.*mucronifolia* Schwägr.*muralis* Hedw.*nevadensis* (Cardot & Thér.) R.H.Zander*obtusifolia* (Schwägr.) Mathieu*plinthobia* (Sull. & Lesq.) Broth.*porteri* (James) R.H.Zander*protobryoides* R.H.Zander*subulata* Hedw.*systylia* (Schimp.) Lindb.*truncata* (Hedw.) Mitt.*vlassovii* (Laz.) Ros & Herrnst.³³⁷³²⁵ See Mastracci (2003).³²⁶ Described as *T. allenii* but corrected orthographically because the species was named for two Allens (Austin 1880): "It gives me great pleasure to be able to dedicate so fine a species to the Allens, father and son."³²⁷ See Hedenäs (2011).³²⁸ For justification for placement in Amblystegiaceae versus Brachytheciaceae see Hedenäs et al. (2020) and references therein.³²⁹ See Hedenäs et al. (2020).³³⁰ See Werner et al. (2014).³³¹ See Werner et al. (2014).³³² See Köckinger et al. (2018).³³³ See Köckinger & Hedenäs (2021).³³⁴ See Werner et al. (2014).³³⁵ See Kellman (2012).³³⁶ See GBIF.org (2024d).³³⁷ See Ros & Herrnstadt (2010).

- Trachycystis** Lindb. (MNIACEAE)
flagellaris (Sull. & Lesq.) Lindb.
- Trachyxiophium** W.R.Buck (PILOTRICHACEAE)
hypnaceum (Müll.Hal.) W.R.Buck
- Trematodon** Michx. (BRUCHIACEAE)
ambiguus (Hedw.) Hornsch.
asanoi Tuzibe
brevicollis Hornsch.
laetevirens Hakelier & J.-P.Frahm³³⁸
longicollis Michx.
montanus Belland & Brassard
- Trichodon** Schimp. (DITRICHACEAE)
cylindricus (Hedw.) Schimp.
oblongus Lindb.³³⁹
- Trichostomopsis** Cardot³⁴⁰ (POTTIACEAE)
australasiae (Hook. & Grev.) H.Rob.
umbrosa (Müll.Hal.) H.Rob.³⁴¹
- Trichostomum** Bruch, *nom. cons.* (POTTIACEAE)
brachydontium Bruch
crispulum Bruch
planifolium (Dixon) R.H.Zander
portoricense H.A.Crum & Steere
- Tripterocladium** (Müll.Hal.) A.Jaeger (LEMBOPHYLLACEAE)
leucocladulum (Müll.Hal.) A.Jaeger
- Triquetrella** Müll.Hal. (POTTIACEAE)
californica (Lesq.) Grout
- Trochophyllohypnum** Jan Kučera & Ignatov³⁴²
(PYLAIADDELPHACEAE)
circinale (Hook.) Jan Kučera & Ignatov
- Tuerckheimia** Broth. (POTTIACEAE)
svihlae (E.B.Bartram) R.H.Zander
- Ulota** D.Mohr (ORTHOTRICHACEAE)
barclayi Mitt.
coarctata (P.Beauv.) Hammar
crispa (Hedw.) Brid.
crispula Bruch³⁴³
curvifolia (Wahlenb.) Sw.
drummondii (Hook. & Grev.) Brid.
hutchinsiae (Sm.) Hammar var. *hutchinsiae*
var. *rufescens* (E.Britton) Dixon
intermedia Schimp.³⁴⁴
- longifolia* Dixon & Sakurai³⁴⁵
megalospora Venturi
obtusiuscula Müll.Hal. & Kindb.
reptans Mitt.³⁴⁶
- Venturiella** Müll.Hal. (ERPODIACEAE)
acrifolia (Pursell) Pursell³⁴⁷
sinensis (Venturi) Müll.Hal. subsp. *angustiannulata*
(D.G.Griffin & Sharp) Pursell
- Vesicularia** (Müll.Hal.) Müll.Hal., *nom. cons.*
(HYPNACEAE)
vesicularis (Schwägr.) Broth.
- Vinealobryum** R.H.Zander³⁴⁸ (POTTIACEAE)
brachyphyllum (Sull.) R.H.Zander
californicum (J.A.Jiménez, D.R.Toren & Shevock)
R.H.Zander³⁴⁹
eckeliae (R.H.Zander) R.H.Zander
insulanum (De Not.) R.H.Zander
nicholsonii (Culm.) R.H.Zander
norrisii (R.H.Zander) J.A.Jiménez & M.J.Cano³⁵⁰
rubiginosum (Mitt.) J.A.Jiménez & M.J.Cano³⁵¹
vineale (Brid.) R.H.Zander
- Voitia** Hornsch.³⁵² (SPLACHNACEAE)
angustata (Hedw.) W.R.Buck & Goffinet³⁵³
hyperborea Grev. & Arn.
mnioides (Hedw.) W.R.Buck & Goffinet³⁵⁴
nivalis Hornsch.
pallida (I.Hagen) W.R.Buck & Goffinet³⁵⁵

³³⁸ See Stehn & Kofranek (2014).³³⁹ See Fedosov et al. (2024).³⁴⁰ See Jiménez et al. (2005, 2022).³⁴¹ See Robinson (1970).³⁴² See Kučera et al. (2019).³⁴³ See Caparrós et al. (2016).³⁴⁴ See Caparrós et al. (2016).³⁴⁵ Based on western North American (Alaska, British Columbia and Washington) specimens held in NY annotated as such by R. Caparrós in 2012.³⁴⁶ The occurrence of this species, originally described from Japan, was considered doubtful by Vitt (2014), although he acknowledged that it may be present. Hence, we prefer to retain it for now.³⁴⁷ See Pursell (2017).³⁴⁸ See Zander (2013).³⁴⁹ See Jiménez et al. (2014).³⁵⁰ See Jiménez et al. (2022).³⁵¹ See Jiménez et al. (2022).³⁵² Phylogenetic inferences resolve *Voitia* within *Tetraplodon* (Lewis et al. 2014; 2017); *Voitia* being an earlier name (Hornschuch 1818), *Tetrapodon* (Bruch & Schimper 1844) is considered a synonym.³⁵³ *Voitia angustata* (Hedw.) W.R.Buck & Goffinet comb. nov. ≡ *Splachnum angustatum* Hedw., Sp. Musc. Frond. 51–52. 1801. Holotype, G00040316.³⁵⁴ *Voitia mnioides* (Hedw.) W.R.Buck & Goffinet comb. nov. ≡ *Splachnum mnioides* Hedw., Sp. Musc. Frond. 51. 1801. Holotype, G00040318.³⁵⁵ *Voitia pallida* (I.Hagen) W.R.Buck & Goffinet comb. nov. ≡ *Tetraplodon pallidus* I.Hagen, Kongel. Norske Vidensk.

paradoxa (R.Br.) W.R.Buck & Goffinet³⁵⁶
urceolata (Hedw.) W.R.Buck & Goffinet³⁵⁷

Warnstorfia Loeske (CALLIERGONACEAE)

badia (Hartm.) W.R.Buck & Goffinet³⁵⁸
exannulata (Schimp.) Loeske³⁵⁹
fluitans (Hedw.) Loeske
pseudosarmentosa (Cardot & Thér.) Tuom. & T.J.Kop.
pseudostraminea (Müll.Hal.) Tuom. & T.J.Kop.
sarmentosa (Wahlenb.) Tuom. & T.J.Kop.
straminea (Dicks. ex Brid.) W.R.Buck & Goffinet³⁶⁰
 var. *straminea*
 var. *patens* (Lindb.) W.R.Buck & Goffinet³⁶¹
trichophylla (Warnst.) Tuom. & T.J.Kop.
tundrae (Arnell) Loeske
wickesiae (Grout) W.R.Buck & Goffinet³⁶²

Weissia Hedw. (POTTIACEAE)

Selsk. Skr. (Trondheim) 1893:75–76. 1894. TYPE: Norway and Sweden, *I. Hagen* (s).

³⁵⁶ *Voitia paradoxa* (R.Br.) W.R.Buck & Goffinet comb. nov. ≡ *Splachnum paradoxum* R.Br., Chlor. Melvill. 44–45. 1823. TYPE: Canada: inter Fort Enterprise et mare arcticum, D. Richardson (BM).

³⁵⁷ *Voitia urceolata* (Hedw.) W.R.Buck & Goffinet comb. nov. ≡ *Splachnum urceolatum* Hedw., Sp. Musc. Frond. 52. 1801. Holotype, G00040320.

³⁵⁸ Previously treated as *Loeskypnum badium* (Anderson et al. 1990; Hedenäs 2014a) and here accommodated within *Warnstorfia* under the new combination *Warnstorfia badia* (Hartm.) W.R.Buck & Goffinet comb. nov. ≡ *Hypnum badium* Hartm., Handb. Skan. Fl. (ed. 5) 332. 1849. TYPE: Sweden: Hn. Lul. lpm. Ananas, 23 Jun. 1848, C.J. Hartman (holotype, UPS).

³⁵⁹ See Hedenäs (2006).

³⁶⁰ Previously treated as *Straminergon straminea* (Dicks. ex Brid.) Hedenäs (Hedenäs 2014a) and here accommodated within *Warnstorfia* under the new combination *Warnstorfia straminea* (Dicks. ex Brid.) W.R.Buck & Goffinet comb. nov. ≡ *Hypnum stramineum* Dicks. ex Brid., Muscol. Recent. 2(2): 172. 1801. TYPE: ‘3, stramineum’ on a sheet with ‘Hypnum’ species in herb. Dickson (holotype, BM).

³⁶¹ Previously recognized as *Calliergon stramineum* var. *patens* (Lindb.) G.Roth (Allen 2014) and here accommodated within *Warnstorfia* under the combination *Warnstorfia straminea* var. *patens* (Lindb.) W.R.Buck & Goffinet comb. nov. ≡ *Amblystegium stramineum* var. *patens* Lindb., Musci Scand. 34. 1879. TYPE: Finland: Helsingfors, S.O. Lindberg (holotype, H).

³⁶² Previously treated as *Loeskypnum wickesiae* (Anderson et al. 1990; Hedenäs 2014a) and here accommodated within *Warnstorfia* under the combination *Warnstorfia wickesiae* (Grout) W.R.Buck & Goffinet comb. nov. ≡ *Calliergon wickesiae* Grout, Moss Fl. N. Amer. 1(4): 250. 1939. TYPE: Canada, Labrador, 1938, Wickes (holotype, DUKE).

armata (Thér. & Trab.) Fedosov³⁶³
brachycarpa (Nees & Hornsch.) Jur.

controversa Hedw.

inoperculata (H.A.Crum) H.A.Crum, Steere & L.E.Anderson

jamaicensis (Mitt.) Grout

ligulifolia (E.B.Bartram) Grout

ludoviciana (Sull.) W.D.Reese & B.A.E.Lemmon

muhlenbergiana (Sw.) W.D.Reese & B.A.E.Lemmon

occidentalis (H.A.Crum) A.H.Stoneb.

phascopsis R.H.Zander

sharpii L.E.Anderson & B.A.E.Lemmon

Wijkia H.A.Crum (PYLAIADIACEAE)

carlottae (W.B.Schofield) H.A.Crum

Zanderella J.A.Jiménez & M.J.Cano³⁶⁴ (POTTIACEAE)

purpurea (Müll.Hal.) J.A.Jiménez & M.J.Cano

Zelometeorium Manuel (BRACHYTHECIACEAE)

patulum (Hedw.) Manuel

Zygodon Hook. & Taylor (ORTHOTRICHACEAE)

conoideus (Dicks.) Hook. & Taylor

dentatus (Limpr.) Breidl. ex Gams

gracilis Wilson

reinwardtii (Hornsch.) A.Braun

rupestris Schimp. ex Lorentz

viridissimus (Dicks.) Brid.

SYNONYMS AND EXCLUDED TAXA

Aloina aloides (W.D.J.Koch ex Schultz) Kindb. var. *ambigua* (Bruch & Schimp.) Limpr. ≡ *A. ambigua* (Bruch & Schimp.) Limpr.

Alsia californica (Hook. & Arn.) Sull. ≡ *Neckera californica* Hook. & Arn.

Amblystegium radicale (P.Beauv.) Schimp. ≡ *Pseudocampyllum radicale* (P.Beauv.) Vanderp. & Hedenäs

Amblystegium serpens (Hedw.) Schimp. var. *juratzkanum* (Schimp.) Rau & Herv. = *A. serpens* (Hedw.) Schimp.

Amblystegium varium (Hedw.) Lindb. ≡ *Hygroamblystegium varium* (Hedw.) Mönk.

Anacolia menziesii (Turner) Paris var. *baueri* (Hampe) Flowers ≡ *A. baueri* (Hampe) Paris

Andraea obovata Thed. = *A. alpina* Hedw.

³⁶³ See Herid et al. (2019).

³⁶⁴ See Jiménez et al. (2022).

- Andreaea rothii* F.Weber & D.Mohr var. *papillosa* Müll.Hal. = *A. rothii* F.Weber & D.Mohr subsp. *falcata* (Schimp.) Lindb.
- Anoetangium handelii* Schiffn. ≡ *Molendoa handelii* (Schiffn.) Brinda & R.H.Zander
- Anoetangium incurvans* (Schimp. ex Besch.) E.B. Bartram = *Molendoa sendtneriana* (Bruch & Schimp.) Limpr.
- Anoetangium tenuinerve* (Limpr.) Paris = *Molendoa sendtneriana* (Bruch & Schimp.) Limpr.
- Anomobryum apiculatum* (Schwägr.) D.Bell & Holyoak ≡ *Bryum apiculatum* Schwägr.
- Anomobryum filiforme* (Dicks.) Husn. = *A. julaceum* (Schrad. ex G.Gaertn., B.Mey. & Scherb.) Schimp.
- Anomobryum leptostomoides* Schimp. = *A. concinatum* (Spruce) A.Jaeger
- Anomodon attenuatus* (Hedw.) Huebener ≡ *Pseudanomodon attenuatus* (Hedw.) Ignatov & Fedosov
- Anomodon longifolius* (Schleich. ex Brid.) Hartm. ≡ *Anomodontella longifolia* (Schleich. ex Brid.) Ignatov & Fedosov
- Anomodon rostratus* (Hedw.) Schimp. ≡ *Claopodium rostratum* (Hedw.) Ignatov
- Anomodontopsis rugelii* (Müll.Hal.) Ignatov & Fedosov ≡ *Anomodon rugelii* (Müll.Hal.) Keissl.
- Antitrichia curtispindula* (Hedw.) Brid. var. *gigantea* Sull. & Lesq. ≡ *A. gigantea* (Sull. & Lesq.) Kindb.
- Aphanorhagma serratum* (Wilson & Hook.) Sull. ≡ *Physcomitrium serratum* (Wilson & Hook.) Müll.Hal.
- Ardeuma annotinum* (Mitt. ex Dixon) R.H.Zander & Brinda ≡ *Hymenostylium annotinum* Mitt. ex Dixon
- Ardeuma aurantiacum* (Mitt.) R.H.Zander & Hedd. ≡ *Hymenostylium aurantiacum* Mitt.
- Ardeuma insigne* (Dixon) R.H.Zander & Hedd. = *Hymenostylium annotinum* Mitt. ex Dixon
- Ardeuma recurvirostrum* (Hedw.) R.H.Zander & Hedd. ≡ *Hymenostylium recurvirostrum* (Hedw.) Dixon
- Arrhenopterum heterostichum* Hedw. ≡ *Aulacomnium heterostichum* (Hedw.) Bruch & Schimp.
- Astomum ludovicianum* (Sull.) Sull. ≡ *Weissia ludoviciana* (Sull.) W.D.Reese & B.A.E.Lemmon
- Astomum muhlenbergianum* (Sw.) Grout ≡ *Weissia muhlenbergiana* (Sw.) W.D.Reese & B.A.E.Lemmon
- Astomum occidentale* Flowers ≡ *Weissia occidentalis* (Flowers) A.H.Stoneb.
- Astomum phascoides* (Drumm.) Grout ≡ *Weissia phascopsis* R.H.Zander
- Atractylocarpus flagellaceus* (Müll.Hal.) R.S.Williams ≡ *Campylopodia flagellacea* (Müll.Hal.) J.-P. Frahm & Isov.
- Atrichum haussknechtii* Jur. & Milde = *A. flavisetum* Mitt.
- Atrichum oerstedianum* (Müll.Hal.) Mitt. - excluded
- Atrichum oerstedianum sensu auct. Amer. non* (Müll. Hal) Mitt. = *A. crispulum* Schimp. ex Besch.³⁶⁵
- Barbella pendula* (Sull.) M.Fleisch. ≡ *Neodictyella pendula* (Sull.) W.R.Buck
- Barbula agraria* Hedw. ≡ *Hyophiladelphus agrarius* (Hedw.) R.H.Zander
- Barbula amplexifolia* (Mitt.) A.Jaeger ≡ *Hydrogonium amplexifolium* (Mitt.) P.C.Chen
- Barbula bolleana* (Müll.Hal.) Broth. ≡ *Hydrogonium bolleanum* (Müll.Hal.) A.Jaeger
- Barbula cancellata* Müll.Hal. ≡ *Hydrogonium consanguineum* (Thwaites & Mitt.) Hilp. var. *cancellatum* (Müll.Hal.) Jan Kučera
- Barbula convoluta* Hedw. var. *convoluta* ≡ *Streblotrichum convolutum* (Hedw.) P.Beauv.
- Barbula convoluta* Hedw. var. *eustegia* (Cardot & Thér.) R.H.Zander ≡ *Streblotrichum convolutum* (Hedw.) P.Beauv. var. *eustegium* (Cardot & Thér.) W.R.Buck & Goffinet
- Barbula convoluta* Hedw. var. *gallinula* R.H.Zander ≡ *Streblotrichum convolutum* (Hedw.) P.Beauv. var. *gallinula* (R.H.Zander) W.R.Buck & Goffinet
- Barbula coreensis* (Cardot) K.Saito = *Hydrogonium amplexifolium* (Mitt.) P.C.Chen
- Barbula cruegeri* Sond. ex Müll.Hal. ≡ *Hydrogonium cruegeri* (Sond. ex Müll.Hal.) Jan Kučera
- Barbula ehrenbergii* (Lorentz) M.Fleisch. = *Hydrogonium bolleanum* (Müll.Hal.) A.Jaeger
- Barbula eustegia* Cardot & Thér. ≡ *Streblotrichum convolutum* (Hedw.) P.Beauv. var. *eustegium* (Cardot & Thér.) W.R.Buck & Goffinet
- Barbula indica* Brid. ex Broth. = *Hydrogonium orientale* (F.Weber) Jan Kučera
- Barbula indica* Brid. ex Broth. var. *gregaria* (Mitt.) R.H.Zander ≡ *Hydrogonium gregarium* (Mitt.) Jan Kučera
- Bartramia glauca* Lorentz = *B. potosica* Mont.
- Bartramia microstoma* Mitt. - excluded
- Bartramia microstoma sensu auct. Amer. non* Mitt. = *B. brevifolia* Brid.³⁶⁶

³⁶⁵ See Merrill & Ireland (2007).

Bartramidula wilsonii Bruch & Schimp. = *Philonotis cernua* (Wilson) D.G.Griffin & W.R.Buck³⁶⁷
Brachymenium klotzschii (Schwägr.) Paris - excluded
Brachymenium klotzschii sensu auct. Amer. non (Schwägr.) Paris = *B. macrocarpum* Cardot³⁶⁸
Brachythecium biventrosus (Müll.Hal.) A.Jaeger = *B. acuminatum* (Hedw.) Austin
Brachythecium bolanderi (Lesq.) A.Jaeger ≡ *Koponeniella bolanderi* (Lesq.) Huttunen & Ignatov
Brachythecium calcareum Kindb. = *B. campestre* (Müll.Hal.) Schimp.
Brachythecium collinum (Schleich. ex Müll.Hal.) Schimp. ≡ *Brachytheciastrum collinum* (Schleich. ex Müll.Hal.) Ignatov & Huttunen
Brachythecium delicatulum Flowers ≡ *Brachytheciastrum delicatulum* (Flowers) Ignatov
Brachythecium digastrum Müll.Hal. & Kindb. = *B. laetum* (Brid.) Schimp.
Brachythecium edentatum R.S.Williams = *Drepanocladus aduncus* (Hedw.) Warnst.³⁶⁹
Brachythecium fendleri (Sull.) A.Jaeger ≡ *Brachytheciastrum fendleri* (Sull.) Ochrya & Żarnowiec
Brachythecium glaciale Schimp. ≡ *Sciuro-hypnum glaciale* (Schimp.) Ignatov & Huttunen
Brachythecium groenlandicum (C.E.O.Jensen) Schljakov = *B. tauriscorum* Molendo & Lorentz
Brachythecium holzingeri (Grout) Grout = *Sciuro-hypnum oedipodium* (Mitt.) Ignatov & Huttunen
Brachythecium hylotapetum B.L.Higinb. & N.L.Higinb. ≡ *Sciuro-hypnum hylotapetum* (B.L.Higinb. & N.L.Higinb.) Ignatov & Huttunen
Brachythecium latifolium Kindb. ≡ *Sciuro-hypnum latifolium* (Kindb.) Ignatov & Huttunen
Brachythecium leibergii Grout ≡ *Brachytheciastrum leibergii* (Grout) Ignatov & Huttunen
Brachythecium mildeanum (Schimp.) Schimp. ex Milde – excluded³⁷⁰
Brachythecium nelsonii Grout = *Sciuro-hypnum latifolium* (Kindb.) Ignatov & Huttunen
Brachythecium oedipodium (Mitt.) A.Jaeger ≡ *Sciuro-hypnum oedipodium* (Mitt.) Ignatov & Huttunen

Brachythecium oxycladon (Brid.) A.Jaeger = *B. acuminatum* (Hedw.) Austin
Brachythecium oxycladon sensu auct. Amer. non (Brid.) A.Jaeger = *B. laetum* (Brid.) Schimp.³⁷¹
Brachythecium plumosum (Hedw.) Schimp. ≡ *Sciuro-hypnum plumosum* (Hedw.) Ignatov & Huttunen
Brachythecium populeum (Hedw.) Schimp. ≡ *Sciuro-hypnum populeum* (Hedw.) Ignatov & Huttunen
Brachythecium reflexum (Starke) Schimp. ≡ *Sciuro-hypnum reflexum* (Starke) Ignatov & Huttunen
Brachythecium rotaeaeum De Not. = *B. capillaceum* (F.Weber & D.Mohr) Giacom.
Brachythecium starkei (Brid.) Schimp. ≡ *Sciuro-hypnum starkei* (Brid.) Ignatov & Huttunen
Brachythecium stereopoma (Spruce ex Mitt.) A.Jaeger = *B. ruderale* (Brid.) W.R.Buck
Brachythecium trachypodium (Brid.) Schimp. ≡ *Brachytheciastrum trachypodium* (Brid.) Ignatov & Huttunen
Brachythecium velutinum (Hedw.) Schimp. ≡ *Brachytheciastrum velutinum* (Hedw.) Ignatov & Huttunen
Brotherella canadensis W.B.Schofield ≡ *Pylaisiadelpha canadensis* (W.B.Schofield) W.R.Buck & Goffinet
Brotherella henonii (Duby) M.Fleisch. ≡ *Pylaisiadelpha henonii* (Duby) W.R.Buck
Brotherella recurvans (Michx.) M.Fleisch. ≡ *Pylaisiadelpha recurvans* (Michx.) W.R.Buck
Brotherella roellii (Renauld & Cardot) M.Fleisch. ≡ *Pylaisiadelpha roellii* (Renauld & Cardot) W.R.Buck
Bryhnia graminicolor (Brid.) Grout ≡ *Koponeniella graminicolor* (Brid.) Huttunen, Ignatov, Min Li & Y.F.Wang
Bryhnia hultenii E.B.Bartram ≡ *Brachythecium novae-angliae* (Sull. & Lesq.) Austin subsp. *hultenii* (E.B.Bartram) Huttunen
Bryhnia novae-angliae (Sull. & Lesq.) Grout ≡ *Brachythecium novae-angliae* (Sull. & Lesq.) Austin
Bryochenea sachalinensis (Lindb.) C.Gao & K.C.Chang ≡ *Echinophyllum sachalinense* (Lindb.) T.J.O'Brien
Bryoerythrophyllum recurvum (Griff.) K.Saito ≡ *Bellibarbula recurva* (Griff.) R.H.Zander

³⁶⁶ See Griffin (2014).

³⁶⁷ See Griffin & Buck (1989).

³⁶⁸ See Spence (2014).

³⁶⁹ See Żarnowiec (2001).

³⁷⁰ See Ignatov (2014).

³⁷¹ See Robinson & Ignatov (1997).

- Bryohaplocladium angustifolium* (Hampe & Müll. Hal.) R.Watan. & Z.Iwats. ≡ *Haplocladium angustifolium* (Hampe & Müll.Hal.) Broth.
- Bryohaplocladium microphyllum* (Sw. ex Hedw.) R. Watan. & Z.Iwats. ≡ *Haplocladium microphyllum* (Sw. ex Hedw.) Broth.
- Bryohaplocladium virginianum* (Brid.) R.Watan. & Z.Iwats. ≡ *Haplocladium virginianum* (Brid.) Broth.
- Bryum axel-blyttii* H.Philb. = *B. calophyllum* R.Br.³⁷²
- Bryum badium* (Brid.) Bruch ex Milde = *B. imbri-catulum* Müll.Hal.
- Bryum barnesii* J.B.Wood ex Schimp. = *B. dichotomum* Hedw.
- Bryum capillare* Hedw. var. *ferchelii* (Funck ex Brid.) Bruch & Schimp. = *B. elegans* Nees ex Brid.
- Bryum gemmascens* Kindb. = *B. sanguilentum* Renault & Cardot
- Bryum laevifilum* Syed = *B. moravicum* Podp.
- Bryum lisae* De Not. var. *cuspidatum* (Bruch & Schimp.) Margad. = *B. creberrimum* Taylor
- Bryum longisetum* Blandow ex Schwägr. var. *labradorensis* (H.Philib.) C.E.O.Jensen = *B. longisetum* Blandow ex Schwägr. var. *longisetum*
- Bryum porsildii* (I.Hagen) C.J.Cox & Hedd. = *Haplodontium macrocarpum* (Drumm.) J.R.Spence
- Bryum pyriferum* Crundw. & H.Whitehouse = *B. valparaisense* Thér.³⁷³
- Bryum renauldii* Röhl - excluded³⁷⁴
- Bryum stirtonii* Schimp. = *B. elegans* Nees ex Brid.
- Buckiella undulata* (Hedw.) Ireland ≡ *Plagiothecium undulatum* (Hedw.) Schimp.
- Bucklandiella affinis* (Schleich. ex F.Weber & D. Mohr) Bedn.-Ochyra & Ochyra ≡ *Racomitrium affine* (Schleich. ex F.Weber & D.Mohr) Lindb.
- Bucklandiella afoninae* (Frisvoll) Bedn.-Ochyra & Ochyra ≡ *Racomitrium afoninae* Frisvoll
- Bucklandiella brevipes* (Kindb.) Bedn.-Ochyra & Ochyra ≡ *Racomitrium brevipes* Kindb.
- Bucklandiella heterosticha* (Hedw.) Bedn.-Ochyra & Ochyra ≡ *Racomitrium heterostichum* (Hedw.) Brid.
- Bucklandiella lawtoniae* (Ireland) Bedn.-Ochyra & Ochyra ≡ *Racomitrium lawtoniae* Ireland
- Bucklandiella macounii* (Kindb.) Bedn.-Ochyra & Ochyra ≡ *Racomitrium macounii* Kindb.
- Bucklandiella macounii* (Kindb.) Bedn.-Ochyra & Ochyra subsp. *alpina* (E.Lawton) Bedn.-Ochyra & Ochyra ≡ *Racomitrium macounii* Kindb. subsp. *alpinum* (E.Lawton) Frisvoll
- Bucklandiella microcarpos* (Hedw.) Bedn.-Ochyra & Ochyra ≡ *Racomitrium microcarpon* (Hedw.) Brid.
- Bucklandiella obesa* (Frisvoll) Bedn.-Ochyra & Ochyra ≡ *Racomitrium obesum* Frisvoll
- Bucklandiella occidentalis* (Renauld & Cardot) Bedn.-Ochyra & Ochyra ≡ *Racomitrium occidentale* (Renauld & Cardot) Renault & Cardot
- Bucklandiella pacifica* (Ireland & J.R.Spence) Bedn.-Ochyra & Ochyra ≡ *Racomitrium pacificum* Ireland & J.R.Spence
- Bucklandiella sudetica* (Funck) Bedn.-Ochyra & Ochyra ≡ *Racomitrium sudeticum* (Funck) Bruch & Schimp.
- Bucklandiella venusta* (Frisvoll) Bedn.-Ochyra & Ochyra ≡ *Racomitrium venustum* Frisvoll
- Calliargon cordifolium* (Hedw.) Kindb. subsp. *orbicularicordatum* (Renauld & Cardot) Grout ≡ *C. orbicularicordatum* (Renauld & Cardot) Broth.
- Calliargon macounii* Kindb. = *C. richardsonii* (Mitt.) Kindb. ex G.Roth
- Calliargon obtusifolium* Karczm. = *C. richardsonii* (Mitt.) Kindb. ex G.Roth
- Calliargon stramineum* (Dicks. ex Brid.) Kindb. ≡ *Warnstorfia straminea* (Dicks. ex Brid.) W.R. Buck & Goffinet
- Calliargon stramineum* (Dicks. ex Brid.) Kindb. var. *patens* (Lindb.) G.Roth ≡ *Warnstorfia straminea* (Dicks. ex Brid.) W.R.Buck & Goffinet var. *patens* (Lindb.) W.R.Buck & Goffinet
- Calliargon trifarium* (F.Weber & D.Mohr) Kindb. ≡ *Drepanocladus trifarius* (F.Weber & D.Mohr) Broth.
- Campyliadelphus chrysophyllus* (Brid.) Kanda ≡ *Campylium chrysophyllum* (Brid.) Lange
- Campylium halleri* (Hedw.) Lindb. ≡ *Campylophyllum halleri* (Hedw.) M.Fleisch.
- Campylium hispidulum* (Brid.) Mitt. ≡ *Campylophyllopsis hispidula* (Brid.) Ochyra
- Campylium polygamum* (Schimp.) Lange & C.E.O. Jensen ≡ *Drepanocladus polygamus* (Schimp.) Hedenäs

³⁷² See Holyoak (2004).³⁷³ See Arts et al. (1995).³⁷⁴ See Toren et al. (2024).

- Campylium radicale* (P.Beauv.) Grout \equiv *Pseudocampylium radicale* (P.Beauv.) Vanderp. & Hedenäs
- Campylium stellatum* (Hedw.) Lange & C.E.O. Jensen var. *arcticum* (R.S.Williams) L.I.Savicz \equiv *Drepanocladus arcticus* (R.S.Williams) Hedenäs
- Campylium stellatum* (Hedw.) Lange & C.E.O. Jensen var. *protensum* (Brid.) Bryhn \equiv *C. protensum* (Brid.) Kindb.
- Campylophyllum hispidulum* (Brid.) Hedenäs \equiv *Campylophyllopsis hispidula* (Brid.) Ochyra
- Campylophyllum sommerfeltii* (Myrin) Hedenäs \equiv *Campylophyllopsis sommerfeltii* (Myrin) Ochyra
- Campylopus aureus* Bosch & Sande Lac. = *C. schmidii* (Müll.Hal.) A.Jaeger
- Campylopus japonicus* Broth. = *C. sinensis* (Müll.Hal.) J.-P.Frahm
- Campylopus paradoxus* Wilson = *C. flexuosus* (Hedw.) Brid.
- Campylopus schwarzii* Schimp. = *C. gracilis* (Mitt.) A.Jaeger
- Campylopus subporodictyon* (Broth.) B.H.Allen & Ireland \equiv *Atractylocarpus subporodictyon* (Broth.) Bonfim Santos & M.Stech
- Campylopus zygodonticarpus* (Müll.Hal.) Paris – excluded³⁷⁵
- Ceratodon purpureus* (Hedw.) Brid. subsp. *conicus* (Hampe ex Müll.Hal.) Dixon \equiv *C. ×conicus* (Hampe ex Müll.Hal.) Lindb.
- Ceratodon purpureus* (Hedw.) Brid. var. *rotundifolius* Berggr. = *C. heterophyllus* Kindb.
- Ceratodon purpureus* (Hedw.) Brid. subsp. *stenocarpus* (Bruch & Schimp.) Dixon \equiv *C. stenocarpus* Bruch & Schimp.
- Cirriphyllum brandegeei* (Austin) Grout \equiv *Brachythecium brandegeei* (Austin) H.Rob.
- Cirriphyllum cirrosum* (Schwägr.) Grout \equiv *Brachythecium cirrosum* (Schwägr.) Schimp.
- Codriophorus acicularis* (Hedw.) P.Beauv. \equiv *Racomitrium aciculare* (Hedw.) Brid.
- Codriophorus aduncooides* (Bedn.-Ochyra) Bedn.-Ochyra & Ochyra \equiv *Racomitrium aduncooides* Bedn.-Ochyra
- Codriophorus brevisetus* (Lindb.) Bedn.-Ochyra & Ochyra \equiv *Racomitrium brevisetum* Lindb.
- Codriophorus corrugatus* Bedn.-Ochyra \equiv *Racomitrium corrugatum* (Bedn.-Ochyra) Larraín & J.Muñoz
- Codriophorus depressus* (Lesq.) Bedn.-Ochyra & Ochyra \equiv *Racomitrium depressum* Lesq.
- Codriophorus fascicularis* (Schrad. ex Hedw.) Bedn.-Ochyra & Ochyra \equiv *Racomitrium fasciculare* (Schrad. ex Hedw.) Brid.
- Codriophorus mollis* (Cardot) Bedn.-Ochyra & Ochyra \equiv *Racomitrium molle* Cardot
- Codriophorus norrisii* (Bedn.-Ochyra & Ochyra) Bedn.-Ochyra & Ochyra \equiv *Racomitrium norrisii* Bedn.-Ochyra & Ochyra
- Codriophorus ryszardii* (Bedn.-Ochyra) Bedn.-Ochyra & Ochyra \equiv *Racomitrium ryszardii* Bedn.-Ochyra & Ochyra
- Codriophorus varius* (Mitt.) Bedn.-Ochyra & Ochyra \equiv *Racomitrium varium* (Mitt.) A.Jaeger
- Cratoneuron arcticum* Steere = *Hygroamblystegium varium* (Hedw.) Mönk.³⁷⁶
- Crossidium rosei* R.S.Williams - excluded³⁷⁷
- Crumia deciduidentata* Sharp & Z.Iwats. \equiv *Tortula deciduidentata* (Sharp & Z.Iwats.) R.H.Zander
- Ctenidium malacodes* Mitt. - excluded
- Ctenidium malacodes sensu auct. Amer. non* Mitt. = *C. subrectifolium* (Brid.) G.Pedano ex W.R.Buck & B.H.Allen³⁷⁸
- Ctenidium molluscum* (Hedw.) Mitt. - excluded
- Ctenidium molluscum sensu auct. Amer. non* (Hedw.) Mitt. = *C. subrectifolium* (Brid.) G.Pedano ex W.R.Buck & B.H.Allen³⁷⁹
- Cynodontium alpestre* (Wahlenb.) Milde \equiv *Cnestrum alpestre* (Wahlenb.) Nyholm
- Cynodontium glaucescens* (Lindb. & Arnell) Kindb. \equiv *Cnestrum glaucescens* (Lindb. & Arnell) Holmen ex Mogensen & Steere
- Cynodontium schisti* (F.Weber & D.Mohr) Lindb. \equiv *Cnestrum schisti* (F.Weber & D.Mohr) I.Hagen
- Cyrto-hypnum involvens* (Hedw.) W.R.Buck & H.A.Crum \equiv *Pelekium involvens* (Hedw.) Touw
- Cyrto-hypnum minutulum* (Hedw.) W.R.Buck & H.A.Crum \equiv *Pelekium minutulum* (Hedw.) Touw
- Cyrto-hypnum pygmaeum* (Schimp.) W.R.Buck & H.A.Crum \equiv *Pelekium pygmaeum* (Schimp.) Touw

³⁷⁵ See Frahm (2007).

³⁷⁶ See Ochyra (1989) for synonymy as *Pseudoleskea chilensis* (Lor.) Ochyra = *Hygroamblyamblystegium varium*.

³⁷⁷ See Delgadillo (1996).

³⁷⁸ See Buck & Allen (2004).

³⁷⁹ See Buck & Allen (2004).

- Cyrtohypnum schistocalyx* (Müll.Hal.) W.R.Buck & H.A.Crum ≡ *Pelekium schistocalyx* (Müll.Hal.) Touw
- Desmatodon cernuus* (Huebener) Bruch & Schimp. ≡ *Tortula cernua* (Huebener) Lindb.
- Desmatodon convolutus* (Brid.) Grout = *Tortula atrovirens* (Sm.) Lindb.
- Desmatodon guepinii* Bruch & Schimp. ≡ *Tortula guepinii* (Bruch & Schimp.) Broth.
- Desmatodon heimii* (Hedw.) Mitt. ≡ *Henediella heimii* (Hedw.) R.H.Zander
- Desmatodon heimii* (Hedw.) Mitt. var. *arcticus* (Lindb.) H.A.Crum = *Henediella heimii* (Hedw.) R.H.Zander
- Desmatodon latifolius* (Hedw.) Brid. = *Tortula hopeana* (Schultz) Ochyra³⁸⁰
- Desmatodon laureri* (Schultz) Bruch & Schimp. ≡ *Tortula laureri* (Schultz) Lindb.
- Desmatodon leucostoma* (R.Br.) Berggr. ≡ *Tortula leucostoma* (R.Br.) Hook. & Grev.
- Desmatodon obtusifolius* (Schwägr.) Schimp. ≡ *Tortula obtusifolia* (Schwägr.) Mathieu
- Desmatodon plinthobius* Sull. & Lesq. ≡ *Tortula plinthobia* (Sull. & Lesq.) Austin
- Desmatodon porteri* James ≡ *Tortula porteri* (James) Broth.
- Desmatodon randii* (Kenn.) Laz. = *Tortula cernua* (Huebener) Lindb.
- Desmatodon systylius* Schimp. ≡ *Tortula systylia* (Schimp.) Lindb.
- Dicranella crispa* (Hedw.) Schimp. ≡ *Dicranellopsis crispa* (Hedw.) Bonfim Santos, Siebel & Fedosov
- Dicranella grevilleana* (Brid.) Schimp. ≡ *Aongstroemia grevilleana* (Brid.) Müll.Hal.
- Dicranella howei* Renauld & Cardot ≡ *Calcidicranella howei* (Renauld & Cardot) Bonfim Santos, Fedosov & Jan Kučera
- Dicranella pacifica* W.B.Schofield ≡ *Calcidicranella pacifica* (W.B.Schofield) Jan Kučera & Fedosov
- Dicranella palustris* (Dicks.) Crundw. ex E.F.Warb. ≡ *Diobelonella palustris* (Dicks.) Ochyra
- Dicranella rufescens* (With.) Schimp. ≡ *Ruficaulis rufescens* (With.) Bonfim Santos & Fedosov
- Dicranella schreberiana* (Hedw.) Dixon ≡ *Aongstroemia schreberiana* (Hedw.) Bonfim Santos & Fedosov
- Dicranella schreberiana* (Hedw.) Dixon var. *robusta* (Schimp. ex Braithw.) H.A.Crum & L.E. Anderson = *Aongstroemia canadensis* (Mitt.) Siebel & Fedosov
- Dicranella staphylina* H.Whitehouse ≡ *Rhizogemma staphylina* (H.Whitehouse) Bonfim Santos, Siebel & Fedosov.
- Dicranella stickinensis* Grout = *Dicranellopsis subulata* (Hedw.) Bonfim Santos, Siebel & Fedosov
- Dicranella subulata* (Hedw.) Schimp. ≡ *Dicranellopsis subulata* (Hedw.) Bonfim Santos, Siebel & Fedosov
- Dicranella varia* (Hedw.) Schimp. ≡ *Calcidicranella varia* (Hedw.) Bonfim Santos, Fedosov & Jan Kučera
- Dicranodontium subporodictyon* Broth. ≡ *Atractyllocarpus subporodictyon* (Broth.) Bonfim Santos & M.Stech
- Dicranoweisia crispula* (Hedw.) Milde ≡ *Hymenoloma crispulum* (Hedw.) Ochyra
- Dicranum angustum* Lindb. = *D. spadiceum* J.E.Zetterst.
- Dicranum fuscescens* Turner var. *flexicaule* (Brid.) Wilson ≡ *D. flexicaule* Brid.
- Dicranum latifolium* J.J.Amann = *D. scoparium* Hedw.
- Didymodon asperifolius* (Mitt.) H.A.Crum, Steere & L.E.Anderson ≡ *Husnotiella asperifolia* (Mitt.) J. A.Jiménez & M.J.Cano
- Didymodon australasiae* (Hook. & Grev.) R.H. Zander ≡ *Trichostomopsis australasiae* (Hook. & Grev.) H.Rob.
- Didymodon bistratosus* Hébrard & R.B.Pierrot = *Zanderella purpurea* (Müll.Hal.) J.A.Jiménez & M.J.Cano
- Didymodon brachyphyllus* (Sull.) R.H.Zander ≡ *Vinealobryum brachyphyllum* (Sull.) R.H. Zander
- Didymodon californicus* J.A. Jiménez, D.R. Toren & Shevock ≡ *Vinealobryum californicum* (J.A. Jiménez, D.R. Toren & Shevock) R.H. Zander
- Didymodon constrictus* (Mitt.) K.Saito ≡ *Geheebia constricta* (Mitt.) R.H.Zander & Caners
- Didymodon eckeliae* R.H.Zander ≡ *Vinealobryum eckeliae* (R.H.Zander) R.H.Zander
- Didymodon fallax* (Hedw.) R.H.Zander ≡ *Geheebia fallax* (Hedw.) R.H.Zander
- Didymodon fallax* (Hedw.) R.H.Zander var. *reflexus* (Brid.) R.H.Zander = *Geheebia ferruginea* (Schimp. ex Besch.) R.H.Zander
- Didymodon ferrugineus* (Schimp. ex Besch.) M.O. Hill ≡ *Geheebia ferruginea* (Schimp. ex Besch.) R.H.Zander

³⁸⁰ See Ochyra (2004).

- Didymodon fragiliscuspis* Broth. ≡ *Husnotiella fragiliscuspis* (Broth.) J.A.Jiménez & M.J.Cano
- Didymodon giganteus* (Funck) Jur. ≡ *Geheebia gigantea* (Funck) Boulay
- Didymodon johansenii* (R.S.Williams) H.A.Crum ≡ *Husnotiella johansenii* (R.S.Williams) J.A.Jiménez & M.J.Cano
- Didymodon leskeoides* K.Saito ≡ *Geheebia leskeoides* (K.Saito) R.H.Zander
- Didymodon maximus* (Syed & Crundw.) M.O.Hill ≡ *Geheebia maxima* (Syed & Crundw.) R.H.Zander
- Didymodon michiganensis* (Steere) K.Saito = *D. maschalogenia* (Renauld & Cardot) Broth.
- Didymodon murrayae* Otnyukova = *Husnotiella fragiliscuspis* (Broth.) J.A.Jiménez & M.J.Cano
- Didymodon nevadensis* R.H.Zander ≡ *Gertrudiella nevadensis* (R.H.Zander) J.A.Jiménez & M.J.Cano
- Didymodon nicholsonii* Culm. ≡ *Vinealobryum nicholsonii* (Culm.) R.H.Zander
- Didymodon norrisii* R.H.Zander ≡ *Vinealobryum norrisii* (R.H.Zander) J.A.Jiménez & M.J.Cano
- Didymodon reedii* H.Rob. = *D. tectorum* (Müll.Hal.) K.Saito
- Didymodon revolutus* (Cardot) R.S.Williams ≡ *Husnotiella revoluta* Cardot
- Didymodon rigidulus* Hedw. var. *ditrichoides* (Broth.) R.H.Zander ≡ *D. ditrichoides* (Broth.) X.J.Li & S. He
- Didymodon rigidulus* Hedw. var. *gracilis* (Hook. & Grev.) R.H.Zander = *D. acutus* (Brid.) K.Saito
- Didymodon rigidulus* Hedw. var. *icmadophilus* (Schimp. ex Müll.Hal.) R.H.Zander ≡ *D. icmadophilus* (Schimp. ex Müll.Hal.) K.Saito
- Didymodon rigidulus* Hedw. var. *subulatus* (Thér. & E.B.Bartram) R.H.Zander ≡ *D. novae-hispaniae* J.A.Jiménez & M.J.Cano
- Didymodon tophaceus* (Brid.) Lisa ≡ *Geheebia tophacea* (Brid.) R.H.Zander
- Didymodon umbrosus* (Müll.Hal.) R.H.Zander ≡ *Trichostomopsis umbrosa* (Müll.Hal.) H.Rob.
- Didymodon vinealis* (Brid.) R.H.Zander ≡ *Vinealobryum vineale* (Brid.) R.H.Zander
- Didymodon vinealis* (Brid.) R.H.Zander var. *brachyphyllus* (Sull.) R.H.Zander ≡ *Vinealobryum brachyphyllum* (Sull.) R.H.Zander
- Didymodon vinealis* (Brid.) R.H.Zander var. *flaccidus* (Bruch & Schimp.) R.H.Zander = *Vinealobryum insulanum* (De Not.) R.H.Zander
- Didymodon vinealis* (Brid.) R.H.Zander var. *luridus* (Hornsch. ex Spreng.) R.H.Zander ≡ *Geheebia lurida* (Hornsch. ex Spreng.) J. A.Jiménez & M.J.Cano
- Didymodon vinealis* (Brid.) R.H.Zander var. *nicholsonii* (Culm.) R.H.Zander ≡ *Vinealobryum nicholsonii* (Culm.) R.H.Zander
- Didymodon vinealis* (Brid.) R.H.Zander var. *rubiginosus* (Mitt.) R.H.Zander ≡ *Vinealobryum rubiginosum* (Mitt.) J.A.Jiménez & M.J.Cano
- Dilutineuron brevisetum* (Lindb.) Bedn.-Ochyra, Sawicki, Ochyra, Szczecińska & Plášek ≡ *Racomitrium brevisetum* Lindb.
- Dilutineuron corrugatum* (Bedn.-Ochyra) Bedn.-Ochyra, Sawicki, Ochyra, Szczecińska & Plášek ≡ *Racomitrium corrugatum* (Bedn.-Ochyra) Larraín & J.Muñoz
- Dilutineuron fasciculare* (Schrad. ex Hedw.) Bedn.-Ochyra, Sawicki, Ochyra, Szczecińska & Plášek ≡ *Racomitrium fasciculare* (Schrad. ex Hedw.) Brid.
- Diphyscium cumberlandianum* Harvill = *D. mucronifolium* Mitt. ex Dozy & Molck.
- Distichium capillaceum* (Hedw.) Bruch & Schimp. var. *curvatum* Flowers = *D. capillaceum* (Hedw.) Bruch & Schimp.³⁸¹
- Ditrichum flexicaule* (Schwägr.) Brockm. ≡ *Flexitrichum flexicaule* (Schwägr.) Ignatov & Fedosov
- Ditrichum gracile* (Mitt.) Kuntze ≡ *Flexitrichum gracile* (Mitt.) Ignatov & Fedosov
- Ditrichum zonatum* (Brid.) Braithw. var. *scabrifolium* Dixon = *D. zonatum* (Brid.) Braithw.
- Drepanocladus aduncus* (Hedw.) Warnst. var. *kneiffii* (Schimp.) Grout = *D. aduncus* (Hedw.) Warnst.
- Drepanocladus aduncus* (Hedw.) Warnst. var. *polycarpus* (Blandow ex Voit) G.Roth = *D. aduncus* (Hedw.) Warnst.³⁸²
- Drepanocladus crassicostatus* Janssens = *D. capillifolius* (Warnst.) Warnst.
- Drepanocladus longifolius* (Wilson ex Mitt.) Broth. ex Paris - excluded, see *D. capillifolius* (Warnst.) Warnst.³⁸³
- Drepanocladus pseudosarmentosus* (Cardot & Thér.) Perss. ≡ *Warnstorfia pseudosarmentosa* (Cardot & Thér.) Tuom. & T.J.Kop.

³⁸¹ See Hedenäs (2021).³⁸² See Hedenäs (2008b).³⁸³ See Saluga et al. (2018).

- Drepanocladus sendtneri* (Schimp. ex H.Müll.) Warnst. – excluded³⁸⁴
- Drepanocladus simplicissimus* Warnst. = *D. aduncus* (Hedw.) Warnst.³⁸⁵
- Dryptodon incurvus* Schwägr. = *Grimmia elatior* Bruch ex Bals.-Criv. & De Not.
- Encalypta affinis* R.Hedw. var. *macounii* (Austin) H. A.Crum & L.E.Anderson ≡ *E. affinis* R.Hedw. subsp. *macounii* (Austin) D.G.Horton
- Encalypta brevicolla* (Bruch & Schimp.) Ångstr. var. *crumiana* (D.G.Horton) H.A.Crum & L.E. Anderson ≡ *E. brevicolla* (Bruch & Schimp.) Ångstr. subsp. *crumiana* D.G.Horton
- Encalypta intermedia* Jur. = *E. pilifera* Funck
- Encalypta microstoma* Bals.-Criv. & De Not. - excluded³⁸⁶
- Encalypta sibirica* (Weinm.) Warnst. - excluded³⁸⁷
- Entodon compressus* (Hedw.) Müll.Hal. = *E. challengerii* (Paris) Cardot
- Epipterygium tozeri* (Grev.) Lindb. - excluded
- Epipterygium tozeri sensu auct. Amer. non* (Grev.) Lindb. = *E. biauratum* Hanusch³⁸⁸
- Erpodium acrifolium* Pursell ≡ *Venturiella acrifolia* (Pursell) Pursell
- Erpodium biseriatum* (Austin) Austin ≡ *Solmsiella biseriata* (Austin) Steere
- Eurhynchium hians* (Hedw.) Sande Lac. ≡ *Oxyrrhynchium hians* (Hedw.) Loeske
- Eurhynchium oreganum* (Sull.) A.Jaeger ≡ *Kindbergia oregana* (Sull.) Ochyra
- Eurhynchium praelongum* (Hedw.) Schimp. ≡ *Kindbergia praelonga* (Hedw.) Ochyra
- Eurhynchium praelongum* (Hedw.) Schimp. var. *stonesii* (Turner) Dixon = *Kindbergia praelonga* (Hedw.) Ochyra
- Eurhynchium pringlei* (Cardot) H.A.Crum & L.E. Anderson ≡ *Donrichardsia pringlei* (Cardot) Huttunen & Ignatov
- Eurhynchium pulchellum* (Hedw.) Jenn. ≡ *Eurhynchiastrum pulchellum* (Hedw.) Ignatov & Huttunen
- Eurhynchium pulchellum* (Hedw.) Jenn. var. *barnesii* (Renauld & Cardot) H.A.Crum, Steere & L.E. Anderson ≡ *Eurhynchiastrum pulchellum* (Hedw.) Ignatov & Huttunen var. *barnesii* (Renauld & Cardot) Ignatov
- Fissidens allenianus* Brugg.-Nann. & Pursell = *F. santa-clarensis* Thér.
- Fissidens clebschii* Steere = *F. taxifolius* Hedw.
- Fissidens garberi* Lesq. & James = *F. pallidinervis* Mitt.
- Fissidens hallii* Austin = *F. amoenus* Müll.Hal.
- Fissidens kochii* H.A.Crum & L.E.Anderson = *F. submarginatus* Bruch
- Fissidens limbatus* Sull. = *F. crispus* Mont.
- Fissidens microcladus* Thwaites & Mitt. – excluded³⁸⁹
- Fissidens milobakeri* L.F.Koch = *F. curvatus* Hornsch.
- Fissidens minutus* Thwaites & Mitt. = *F. pallidinervis* Mitt.
- Fissidens neonii* (E.B.Bartram) Grout = *F. scalaris* Mitt.
- Fissidens obtusifolius* Wilson var. *apiculatus* Grout = *F. sublimbatus* Grout
- Fissidens obtusifolius* Wilson var. *kansanus* Renauld & Cardot = *F. obtusifolius* Wilson
- Fissidens obtusifolius* Wilson var. *marginatus* Flowers = *F. obtusifolius* Wilson
- Fissidens papillosus* Sande Lac. = *F. serratus* Müll. Hal.
- Fissidens radicans* Mont. - excluded
- Fissidens ravenelii* Sull. = *F. elegans* Brid.
- Fissidens reesei* H.A.Crum & L.E.Anderson = *F. leptophyllus* Mont.
- Fontinalis antipyretica* Hedw. var. *gigantea* (Sull.) Sull. = *F. antipyretica* Hedw.
- Fontinalis antipyretica* Hedw. var. *oreganensis* Renauld & Cardot = *F. antipyretica* Hedw.
- Fontinalis filiformis* Sull. & Lesq. = *F. sullivantii* Lindb.
- Fontinalis flaccida* Renauld & Cardot = *F. sullivantii* Lindb.
- Fontinalis hypnoides* Hartm. var. *duriaei* (Schimp.) Husn. = *F. hypnoides* Hartm.
- Fontinalis langloisii* Cardot = *F. sullivantii* Lindb.
- Fontinalis macmillanii* Cardot = *F. hypnoides* Hartm.
- Fontinalis missourica* Cardot = *F. sullivantii* Lindb.
- Fontinalis novae-angliae* Sull. var. *cymbifolia* (Austin) W.H.Welch = *F. novae-angliae* Sull.

³⁸⁴ See Hedenäs (2014b).³⁸⁵ See Hedenäs (2008b).³⁸⁶ See Horton (1983).³⁸⁷ See Magill (2006).³⁸⁸ See Hanusch et al. (2020).³⁸⁹ See Pursell & Vital (1986), under *F. gardneri*, of which it is a synonym.

Fontinalis novae-angliae Sull. var. *lorenziae* Cardot =
F. novae-angliae Sull.
Frisvöllia varia (Mitt.) Sawicki, Szczecińska, Bedn.-
 Ochyra & Ochyra ≡ *Racomitrium varium* (Mitt.)
 A.Jaeger
Funaria americana Lindb. ≡ *Entosthodon ameri-*
canus (Lindb.) Fife
Funaria apiculatopilosa Cardot ≡ *Entosthodon api-*
culatopilosus (Cardot) Fife
Funaria groutiana Fife = *F. arctica* (Berggr.) Kindb.
Funaria muhlenbergii Turner ≡ *Entosthodon muh-*
lenbergii (Turner) Fife
Funaria serrata Brid. ≡ *Entosthodon serratus* (Brid.)
 Fife
Gemmabryum apiculatum (Schwägr.) J.R.Spence &
 H.P.Ramsay ≡ *Bryum apiculatum* Schwägr.
Gemmabryum badium (Brid.) J.R.Spence = *Bryum*
imbricatulum Müll.Hal.
Gemmabryum barnesii (J.B.Wood ex Schimp.) J.R.
 Spence = *Bryum dichotomum* Hedw.
Gemmabryum brassicoides J.R.Spence & Kellman ≡
Bryum brassicoides (J.R.Spence & Kellman) W.
 R.Buck & Goffinet
Gemmabryum caespiticium (Hedw.) J.R.Spence ≡
Bryum caespiticium Hedw.
Gemmabryum californicum (Sull.) J.R.Spence ≡
Bryum californicum Sull.
Gemmabryum coronatum (Schwägr.) J.R.Spence ≡
Bryum coronatum Schwägr.
Gemmabryum demaretianum (Arts) J.R.Spence ≡
Bryum demaretianum Arts
Gemmabryum dichotomum (Hedw.) J.R.Spence ≡
Bryum dichotomum Hedw.
Gemmabryum exile (Dozy & Molk.) J.R.Spence & H.
 P.Ramsay ≡ *Brachymenium exile* (Dozy &
 Molk.) Bosch & Sande Lac.
Gemmabryum gemmiferum (R.Wilczek & Demaret)
 J.R.Spence ≡ *Bryum gemmiferum* R.Wilczek &
 Demaret
Gemmabryum gemmilucens (R.Wilczek & Demaret)
 J.R.Spence ≡ *Bryum gemmilucens* R.Wilczek &
 Demaret
Gemmabryum klinggraeffii (Schimp.) J.R.Spence ≡
Bryum klinggraeffii Schimp.
Gemmabryum kunzei (Hornsch.) J.R.Spence ≡
Bryum kunzei Hornsch.
Gemmabryum radiculosum (Brid.) J.R.Spence & H.
 P.Ramsay ≡ *Bryum radiculosum* Brid.

Gemmabryum ruderale (Crundw. & Nyholm) J.R.
 Spence ≡ *Bryum ruderale* Crundw. & Nyholm
Gemmabryum subapiculatum (Hampe) J.R.Spence
 ≡ *Bryum subapiculatum* Hampe
Gemmabryum tenuisetum (Limpr.) J.R.Spence & H.
 P.Ramsay ≡ *Bryum tenuisetum* Limpr.
Gemmabryum valparaisense (Thér.) J.R.Spence ≡
Bryum valparaisense Thér.
Gemmabryum vinosum J.R.Spence & Kellman ≡
Bryum vinosum (J.R.Spence & Kellman) W.R.
 Buck & Goffinet
Gemmabryum violaceum (Crundw. & Nyholm) J.R.
 Spence ≡ *Bryum violaceum* Crundw. & Nyholm
Grimmia affinis Hornsch. = *G. longirostris* Hook.
Grimmia anomala Hampe ex Schimp. ≡ *Dryptodon*
anomalus (Hampe ex Schimp.) Loeske
Grimmia arcuatifolia Kindb. ≡ *Dryptodon arcuatifo-*
lius (Kindb.) Ochyra & Żarnowiec
Grimmia bernoullii Müll.Hal. = *G. ovalis* (Hedw.)
 Lindb.³⁹⁰
Grimmia brevirostris R.S.Williams = *G. hamulosa*
 Lesq.
Grimmia fuscolutea Hook. ≡ *Dryptodon fuscoluteus*
 (Hook.) Ochyra & Żarnowiec
Grimmia hartmanii Schimp. ≡ *Dryptodon hartmanii*
 (Schimp.) Limpr.
Grimmia hermannii H.A.Crum = *Dryptodon muel-*
lenbeckii (Schimp.) Loeske
Grimmia leibergii Paris ≡ *Dryptodon leibergii* (Paris)
 Ochyra & Żarnowiec
Grimmia lisae De Not. ≡ *Dryptodon lisae* (De Not.)
 Loeske
Grimmia muehlenbeckii Schimp. ≡ *Dryptodon*
muehlenbeckii (Schimp.) Loeske
Grimmia poecilostoma Cardot & Sebille = *G. crinito-*
leucophaea Cardot
Grimmia pulvinata (Hedw.) Sm. var. *africana* (Hedw.)
 Hook.f. & Wilson = *G. pulvinata* (Hedw.) Sm.
Grimmia ramondii (DC.) Margad. = *Dryptodon pat-*
ens (Dicks. ex Hedw.) Brid.
Grimmia tenerrima Renauld & Cardot = *G. sessitana*
 De Not.
Grimmia teretinervis Limpr. ≡ *Schistidium tereti-*
nerve (Limpr.) Limpr.

³⁹⁰ The species was reported from the United States by Weber et al. (2003) but then considered conspecific with *G. ovalis* by Muñoz (1999).

- Grimmia torngakiana* Brassard & Hedd. = *G. incurva* Schwägr.
- Grimmia torquata* Hook. ≡ *Dryptodon torquatus* (Hook.) Brid.
- Grimmia torquata* Hook. var. *flettii* (Holz.) Vaar. = *Dryptodon torquatus* (Hook.) Brid.³⁹¹
- Grimmia trichophylla* Grev. ≡ *Dryptodon trichophyllus* (Grev.) Brid.
- Gymnostomiella orcuttii* E.B.Bartram = *G. vernicosa* (Hook. ex Harv.) M.Fleisch.
- Gyroweisia reflexa* (Brid.) Schimp. - excluded³⁹²
- Haplohymenium triste* (Ces.) Kindb. ≡ *Anomodon tristis* (Ces.) Sull.
- Hedwigia ciliata* (Hedw.) Boucher fo. *detonsa* (M. Howe) G.N.Jones ≡ *H. detonsa* (M.Howe) W.R. Buck & D.H.Norris
- Hedwigia ciliata* (Hedw.) Boucher var. *leucophaea* Bruch & Schimp. = *H. emodica* Hampe ex Müll. Hal.
- Hedwigia nemoralis* Ignatova, Ignatov & Fedosov³⁹³ = *H. filiformis* (Michx.) P.Beauv.
- Helodium blandowii* (F.Weber & D.Mohr) Warnst. ≡ *H. blandovii* (F.Weber & D.Mohr) Warnst.
- Helodium blandovii* (F.Weber & D.Mohr) Warnst. var. *elodioides* (Renauld & Cardot) H.A.Crum, Steere & L.E.Anderson ≡ *H. elodioides* (Renauld & Cardot) B.H.Allen
- Herzogiella adscendens* (Lindb.) Z.Iwats. & W.B. Schofield ≡ *Aquilonium adscendens* (Lindb.) Hedenäs, Schlesak & D.Quandt
- Heterocladium dimorphum* (Brid.) Schimp. ≡ *Heteroclediella dimorpha* (Brid.) Ignatov & Fedosov
- Heterocladium procurrens* (Mitt.) A.Jaeger ≡ *Heteroclediella procurrens* (Mitt.) Ignatov & Fedosov
- Heterophyllum affine* (Hook.) M.Fleisch. = *H. nemorosum* (W.D.J.Koch ex Brid.) Kindb.
- Homalothecium pinnatifidum* (Sull. & Lesq.) E. Lawton = *H. aureum* (Spruce) H.Rob.
- Hygroamblystegium tenax* (Hedw.) Jenn. var. *spinifolium* (Schimp.) Jenn ≡ *Cratoneuron filicinum* (Hedw.) Spruce³⁹⁴
- Hygrohypnum alpestre* (Hedw.) Loeske ≡ *Platyhypnum alpestre* (Hedw.) Ochyra
- Hygrohypnum alpinum* (Lindb.) Loeske ≡ *Platyhypnum alpinum* (Lindb.) Loeske
- Hygrohypnum bestii* (Renauld & Bryhn) Holz. ≡ *Platyhypnum bestii* (Renauld & Bryhn) Ochyra
- Hygrohypnum cochleariifolium* (Venturi) Broth. ≡ *Platyhypnum cochleariifolium* (Venturi) Ochyra
- Hygrohypnum duriusculum* (De Not.) D.W.Jamieson ≡ *Platyhypnum duriusculum* (De Not.) Ochyra
- Hygrohypnum eugyrium* (Schimp.) Loeske ≡ *Pseudohygrohypnum eugyrium* (Schimp.) Kanda
- Hygrohypnum micans* (Mitt.) Broth. ≡ *Hageniella micans* (Mitt.) B.C.Tan & Y.Jia
- Hygrohypnum molle* (Hedw.) Loeske ≡ *Platyhypnum molle* (Hedw.) Loeske
- Hygrohypnum montanum* (Lindb.) Broth. ≡ *Campylophyllum montanum* (Lindb.) B.H. Allen
- Hygrohypnum norvegicum* (Schimp.) J.J.Amann ≡ *Platyhypnum norvegicum* (Schimp.) Ochyra
- Hygrohypnum ochraceum* (Turner ex Wilson) Loeske ≡ *Hygrohypnella ochracea* (Turner ex Wilson) Ignatov & Ignatova
- Hygrohypnum polare* (Lindb.) Loeske ≡ *Hygrohypnella polaris* (Lindb.) Ignatov & Ignatova
- Hygrohypnum smithii* (Sw.) Broth. ≡ *Platyhypnum smithii* (Sw.) Ochyra
- Hygrohypnum subeugyrium* (Renauld & Cardot) Broth. ≡ *Pseudohygrohypnum subeugyrium* (Renauld & Cardot) Ignatov & Ignatova
- Hymenostylium insigne* (Dixon) Podp. = *H. annotinum* Mitt. ex Dixon
- Hymenostylium recurvirostrum* (Hedw.) Dixon var. *insigne* (Dixon) E.B.Bartram = *H. annotinum* Mitt. ex Dixon
- Hypnum bambergeri* Schimp. ≡ *Campylium bambergeri* (Schimp.) Hedenäs, Schlesak & D.Quandt
- Hypnum callichroum* Brid. ≡ *Stereodon callichroum* (Brid.) Brid.
- Hypnum circinale* Hook. ≡ *Trochophyllohypnum circinale* (Hook.) Jan Kučera & Ignatov
- Hypnum cupressiforme* Hedw. var. *lacunosum* Brid. = *H. cupressiforme* Hedw. var. *cupressiforme*
- Hypnum cupressiforme* Hedw. var. *resupinatum* (Taylor) Schimp. ≡ *H. resupinatum* Taylor

³⁹¹ See Muñoz & Pando (2000). Note that Maier (2010) placed this var. in synonymy with *G. elongata*.

³⁹² First reported by Conard (1945) and confirmed by Manierre (1998) but excluded by Zander (2007c).

³⁹³ See Ignatova et al. (2016).

³⁹⁴ See Hedenäs (2003).

- Hypnum cupressiforme* Hedw. var. *subjulaceum* Molendo ≡ *H. subjulaceum* (Molendo) Hedenäs, Schlesak & D.Quandt
- Hypnum curvifolium* Hedw. ≡ *Calliergonella curvifolia* (Hedw.) B.H.Allen
- Hypnum dieckii* Renauld & Cardot ≡ *Calliergonellopsis dieckii* (Renauld & Cardot) Jan Kučera & Ignatov
- Hypnum fujiyamae* (Broth.) Paris ≡ *Callicladium fujiyamae* (Broth.) Jan Kučera & Ignatov
- Hypnum geminum* (Mitt.) Lesq. & James = *Isopterygiella pulchella* (Hedw.) Ignatov & Ignatova
- Hypnum hamulosum* Schimp. ≡ *Stereodon hamulosus* (Schimp.) Lindb.
- Hypnum holmenii* Ando ≡ *Stereodon holmenii* (Ando) Ignatov & Ignatova
- Hypnum imponens* Hedw. ≡ *Callicladium imponens* (Hedw.) Hedenäs, Schlesak & D.Quandt
- Hypnum lindbergii* Mitt. ≡ *Calliergonella lindbergii* (Mitt.) Hedenäs
- Hypnum mammillatum* (Brid.) Lindb. ≡ *H. andoi* A.J.E.Sm.
- Hypnum pallescens* (Hedw.) P. Beauv. ≡ *Jochenia pallescens* (Hedw.) Hedenäs, Schlesak & D. Quandt
- Hypnum pallescens* (Hedw.) P. Beauv. var. *protuberans* (Brid.) Austin ≡ *Jochenia protuberans* (Brid.) Jan Kučera & Ignatov
- Hypnum plicatum* (Lindb.) A.Jaeger ≡ *Aquilonium plicatum* (Lindb.) Hedenäs, Schlesak & D.Quandt
- Hypnum pratense* W.D.J.Koch ex Spruce ≡ *Stereodon pratensis* (W.D.J.Koch ex Spruce) E.Britton.
- Hypnum procerrimum* Molendo ≡ *Pseudostereodon procerrimus* (Molendo) M.Fleisch.
- Hypnum recurvatum* (Lindb. & Arnell) Kindb. = *Drepanium fastigiatum* (Brid.) Lange & C.E.O. Jensen
- Hypnum revolutum* (Mitt.) Lindb. ≡ *Roaldia revoluta* (Mitt.) P.E.A.S.Câmara & Carv.-Silva
- Hypnum subimponens* Lesq. ≡ *Stereodon subimponens* (Lesq.) Broth.
- Hypnum vaucheri* Lesq. ≡ *Buckia vaucheri* (Lesq.) D.Ríos, M.T.Gallego & J.Guerra
- Hypopterygium fauriei* Besch. = *H. flavolimbatum* Müll.Hal.
- Hypopterygium tamariscinum sensu auct. Amer. non* (Hedw.) Brid. = *H. tamarisci* (Sm.) Müll.Hal.
- Imbribryum alpinum* (Huds. ex With.) N.Pedersen ≡ *Bryum alpinum* Huds. ex With.
- Imbribryum gemmiparum* (De Not.) J.R.Spence ≡ *Bryum gemmiparum* De Not.
- Imbribryum microchaeton* (Hampe) J.R.Spence ≡ *Bryum microchaeton* Hampe
- Imbribryum mildeanum* (Jur.) J.R.Spence ≡ *Bryum mildeanum* Jur.
- Imbribryum miniatum* (Lesq.) J.R.Spence ≡ *Bryum miniatum* Lesq.
- Imbribryum muehlenbeckii* (Bruch & Schimp.) N. Pedersen ≡ *Bryum muehlenbeckii* Bruch & Schimp.
- Imbribryum torenii* J.R.Spence & Shevock ≡ *Bryum torenii* (J.R.Spence & Shevock) W.R.Buck & Goffinet
- Isopterygiopsis alpicola* (Lindb.) Hedenäs ≡ *Isopterygiella alpicola* (Lindb.) Ignatov & Ignatova
- Isopterygiopsis pulchella* (Hedw.) Z.Iwats. ≡ *Isopterygiella pulchella* (Hedw.) Ignatov & Ignatova
- Isothecium cardotii* Kindb. ≡ *Pseudisothecium cardotii* (Kindb.) Ignatova, Fedosov & Ignatov
- Isothecium cristatum* (Hampe) H.Rob. ≡ *Pseudisothecium cristatum* (Hampe) Ignatova, Fedosov & Ignatov
- Isothecium myosuroides* Brid. ≡ *Pseudisothecium myosuroides* (Brid.) Grout
- Isothecium stoloniferum* Brid. ≡ *Pseudisothecium stoloniferum* (Brid.) Grout
- Kiaeria blyttii* (Bruch & Schimp.) Broth. ≡ *Arctoa blyttii* (Bruch & Schimp.) Loeske
- Kiaeria falcata* (Hedw.) I.Hagen ≡ *Pseudoblindia falcata* (Hedw.) Fedosov, M.Stech & Ignatov
- Kiaeria glacialis* (Berggr.) I.Hagen ≡ *Arctoa glacialis* (Berggr.) Fedosov, Jan Kučera & M.Stech
- Kiaeria starkei* (F.Weber & D.Mohr) I.Hagen ≡ *Arctoa starkei* (F.Weber & D.Mohr) Loeske
- Leptodictyum humile* (P.Beauv.) Ochyra ≡ *Hygroamblystegium humile* (P.Beauv.) Vanderp., Hedenäs & Goffinet
- Leptodontium viticulosoides* (P.Beauv.) Wijk & Margad. var. *sulphureum* (Müll.Hal.) R.H.Zander = *L. excelsum* (Sull.) E.Britton
- Leptostomopsis nivea* (Besch.) J.R.Spence ≡ *Brachymenium niveum* Besch.
- Leptostomopsis systylia* (Müll.Hal.) J.R.Spence ≡ *Brachymenium systylium* (Müll.Hal.) A.Jaeger
- Lescuraea arizonae* (R.S.Williams) P.S.Wilson & D. H.Norris ≡ *Pseudoleskeella arizonae* (R.S.Williams) E.Lawton

- Leskea angustata* Taylor - excluded³⁹⁵
Leskeella nervosa (Brid.) Loeske ≡ *Pseudoleskeella nervosa* (Brid.) Nyholm
Leucodon andrewsianus (H.A.Crum & L.E.Anderson) W.D.Reese & L.E.Anderson = *L. sciuroides* (Hedw.) Schwägr.
Leucodon brachypus Brid. var. *andrewsianus* H.A.Crum & L.E.Anderson = *L. sciuroides* (Hedw.) Schwägr.
Limprichtia cossonii (Schimp.) L.E.Anderson, H.A.Crum & W.R.Buck ≡ *Scorpidium cossonii* (Schimp.) Hedenäs
Limprichtia revolvens (Sw.) Loeske ≡ *Scorpidium revolvens* (Sw.) Rubers
Loeskypnum badium (Hartm.) H.K.G.Paul ≡ *Warnstorfia badia* (Hartm.) W.R.Buck & Goffinet
Loeskypnum wickesiae (Grout) Tuom. ≡ *Warnstorfia wickesiae* (Grout) W.R.Buck & Goffinet
Macrocoma sullivantii (Müll.Hal.) Grout ≡ *M. tenuis* (Hook. & Grev.) Vitt subsp. *sullivantii* (Müll.Hal.) Vitt
Macromitrium richardii Schwägr. = *M. viticulosum* (Raddi) Brid.³⁹⁶
Meiotrichum lyallii (Mitt.) G.L.Sm.Merr. ≡ *Polytrichastrum lyallii* (Mitt.) G.L.Sm.³⁹⁷
Microbryum davallianum (Sm.) R.H.Zander var. *commutatum* (Limpr.) R.H.Zander ≡ *M. commutatum* (Limpr.) Cl.Schneid., Th.Schneid. & Mahévas
Microbryum davallianum (Sm.) R.H.Zander var. *conicum* (Schleich. ex Schwägr.) R.H.Zander ≡ *M. conicum* (Schleich. ex Schwägr.) Cl.Schneid., Th.Schneid. & Mahévas
Microbryum starckeanum (Hedw.) R.H.Zander var. *brachyodus* (Bruch & Schimp.) R.H.Zander = *M. muticum* (Venturi) Cl.Schneid., Th.Schneid. & Mahévas
Microbryum starckeanum (Hedw.) R.H.Zander var. *fosbergii* (E.B.Bartram) R.H.Zander ≡ *M. fosbergii* (E.B.Bartram) Ros, O.Werner & Rams
Microbryum vlassovii (Laz.) R.H.Zander ≡ *Tortula vlassovii* (Laz.) Ros & Herrnst.
Micromitrium austinii Austin = *M. tenerum* (Bruch & Schimp.) Crosby
Mielichhoferia macrocarpa (Drumm.) Bruch & Schimp. ≡ *Haplodontium macrocarpum* (Drumm.) J.R.Spence
Mielichhoferia tehamensis Showers ≡ *Haplodontium tehamense* (Showers) J.R.Spence
Monocryphaea ravenelii (Austin) P.Rao ≡ *Cryphaea ravenelii* Austin
Neckera besseri (Łobarz.) Jur. ≡ *Alleniella besseri* (Łobarz.) S.Olsson, Enroth & D.Quandt
Neckera complanata (Hedw.) Huebener ≡ *Alleniella complanata* (Hedw.) S.Olsson, Enroth & D.Quandt
Neohyophila sprengelii (Schwägr.) H.A.Crum ≡ *Plaubelia sprengelii* (Schwägr.) R.H.Zander
Neohyophila stomatodonta (Cardot) H.A.Crum ≡ *Plaubelia stomatodonta* (Cardot) R.H.Zander
Neotrichostomum crispulum (Bruch) R.H.Zander ≡ *Trichostomum crispulum* Bruch
Niphotrichum canescens (Hedw.) Bedn.-Ochyra & Ochyra ≡ *Racomitrium canescens* (Hedw.) Brid.
Niphotrichum canescens (Hedw.) Bedn.-Ochyra & Ochyra subsp. *latifolium* (C.E.O.Jensen) Bedn.-Ochyra & Ochyra ≡ *Racomitrium canescens* (Hedw.) Brid. subsp. *latifolium* (C.E.O.Jensen) Frisvoll
Niphotrichum elongatum (Ehrh. ex Frisvoll) Bedn.-Ochyra & Ochyra ≡ *Racomitrium elongatum* Ehrh. ex Frisvoll
Niphotrichum ericoides (Brid.) Bedn.-Ochyra & Ochyra ≡ *Racomitrium ericoides* (Brid.) Brid.
Niphotrichum muticum (Kindb.) Bedn.-Ochyra & Ochyra ≡ *Racomitrium muticum* (Kindb.) Frisvoll
Niphotrichum panschii (Müll.Hal.) Bedn.-Ochyra & Ochyra ≡ *Racomitrium panschii* (Müll.Hal.) Kindb.
Niphotrichum pygmaeum (Frisvoll) Bedn.-Ochyra & Ochyra ≡ *Racomitrium pygmaeum* Frisvoll
Oncophorus raii (Austin) Grout ≡ *Symblepharis raii* (Austin) R.S.Williams ex Broth.
Oncophorus wahlenbergii Brid. ≡ *Brideliella wahlenbergii* (Brid.) Fedosov, M.Stech & Ignatov
Orthothecium chryseon (Schwägr.) Schimp. var. *cochleariifolium* (Lindb.) Limpr. = *O. chryseon* (Schwägr.) Schimp.³⁹⁸
Orthothecium diminutivum (Grout) H.A.Crum, Steere & L.E.Anderson = *Isopterygiella pulchella* (Hedw.) Ignatov & Ignatova

³⁹⁵ See Allen (2018) sub *Leskeadelphus*.³⁹⁶ See Valente et al. (2020).³⁹⁷ See Bell & Hyvönen (2010).³⁹⁸ See Ignatov et al. (2020).

- Orthotrichum affine* Brid. ≡ *Lewinskya affinis* (Brid.) F.Lara, Garilleti & Goffinet var. *affinis*
- Orthotrichum affine* Brid. var. *bohemicum* Plášek & Sawicki ≡ *Lewinskya affinis* (Brid.) F.Lara, Garilleti & Goffinet var. *bohemica* (Plášek & Sawicki) Plášek
- Orthotrichum bolanderi* Sull. ≡ *Lewinskya bolanderi* (Sull.) F.Lara, Garilleti & Goffinet
- Orthotrichum elegans* Schwägr. ex Hook. & Grev. ≡ *Lewinskya elegans* (Schwägr. ex Hook. & Grev.) F.Lara, Garilleti & Goffinet
- Orthotrichum epapillosum* E.Lawton & F.J.Herm. = *O. euryphyllum* Venturi³⁹⁹
- Orthotrichum exiguum* Sull. ≡ *Leratia exigua* (Sull.) Goffinet
- Orthotrichum fenestratum* Cardot & Thér. ≡ *Lewinskya fenestrata* (Cardot & Thér.) F.Lara, Garilleti & Goffinet
- Orthotrichum gymnostomum* Bruch ex Brid. ≡ *Nyholmiella gymnostoma* (Bruch ex Brid.) Holmen & E.Warncke
- Orthotrichum holzingeri* Renaud & Cardot ≡ *Lewinskya holzingeri* (Renaud & Cardot) F.Lara, Garilleti & Goffinet
- Orthotrichum keeverae* H.A.Crum & L.E.Anderson ≡ *Lewinskya keeverae* (H.A.Crum & L.E.Anderson) F.Lara, Garilleti & Goffinet
- Orthotrichum kellmanii* D.H.Norris, Shevock & Goffinet = *Orthotrichum shevockii* Lewinsky-Haapasaari & D.H.Norris⁴⁰⁰
- Orthotrichum laevigatum* J.E.Zetterst. ≡ *Lewinskya laevigata* (J.E.Zetterst.) F.Lara, Garilleti & Goffinet
- Orthotrichum lyellii* Hook. & Taylor ≡ *Pulvigerella lyellii* (Hook. & Taylor) Plášek, Sawicki & Ochyra
- Orthotrichum lyellii* Hook. & Taylor var. *howei* Renaud & Cardot ≡ *Pulvigerella howei* (Renaud & Cardot) F.Lara, Draper & Garilleti
- Orthotrichum obtusifolium* Brid. ≡ *Nyholmiella obtusifolia* (Brid.) Holmen & E.Warncke
- Orthotrichum papillosum* Hampe ≡ *Pulvigerella papillosa* (Hampe) F.Lara, Draper & Garilleti
- Orthotrichum praemorsum* Venturi ≡ *Lewinskya praemorsa* (Venturi) F.Lara, Garilleti & Goffinet
- Orthotrichum pycnophyllum* Schimp. ≡ *Lewinskya pycnophylla* (Schimp.) F.Lara, Garilleti & Goffinet
- Orthotrichum pylaisii* Brid. ≡ *Lewinskya pylaisii* (Brid.) F.Lara, Garilleti & Goffinet
- Orthotrichum rupestre* Schleich. ex Schwägr. ≡ *Lewinskya rupestris* (Schleich. ex Schwägr.) F.Lara, Garilleti & Goffinet
- Orthotrichum shawii* Wilson ≡ *Lewinskya shawii* (Wilson) F.Lara, Garilleti & Goffinet
- Orthotrichum sordidum* Sull. & Lesq. ≡ *Lewinskya sordida* (Sull. & Lesq.) F.Lara, Garilleti & Goffinet
- Orthotrichum speciosum* Nees ≡ *Lewinskya speciosa* (Nees) F.Lara, Garilleti & Goffinet
- Orthotrichum spjutii* D.H.Norris & Vitt ≡ *Lewinskya spjutii* (D.H.Norris & Vitt) F.Lara, Garilleti & Goffinet
- Orthotrichum striatum* Hedw. ≡ *Lewinskya striata* (Hedw.) F.Lara, Garilleti & Goffinet
- Orthotrichum tenellum* Bruch ex Brid. - excluded
- Oxystegus daldinianus* (De Not.) Köckinger, O. Werner & Ros ≡ *Chionoloma daldinianum* (De Not.) M.Alonso, M.J.Cano & J.A.Jiménez
- Oxystegus maragniphyllus* R.H.Zander & Eckel ≡ *Chionoloma maragniphyllum* (R.H.Zander & Eckel) W.R.Buck & Goffinet
- Oxystegus spiralis* (Grout) H.A.Crum & L.E.Anderson = *Chionoloma stenocarpum* (Thér.) M.Alonso, M.J.Cano & J.A.Jiménez
- Oxystegus tenuirostris* (Hook. & Taylor) A.J.E.Sm. ≡ *Chionoloma tenuirostre* (Hook. & Taylor) M.Alonso, M.J.Cano & J.A.Jiménez
- Palustriella commutata* (Hedw.) Ochyra - excluded⁴⁰¹
- Papillaria nigrescens* (Sw. ex Hedw.) A.Jaeger ≡ *Meteorium nigrescens* (Sw. ex Hedw.) Dozy & Molk.
- Paraleptodontium recurvifolium* (Taylor) D.G.Long ≡ *Chionoloma recurvifolium* (Taylor) M.Alonso, M.J.Cano & J.A.Jiménez
- Phascum cuspidatum* Schreb. ex Hedw. = *Tortula acaulon* (With.) R.H.Zander
- Phascum floerkeanum* F.Weber & D.Mohr ≡ *Microbryum floerkeanum* (F.Weber & D.Mohr) Schimp.
- Phascum hyalinotrichum* Cardot & Thér. ≡ *Stegonia hyalinotricha* (Cardot & Thér.) R.H.Zander

³⁹⁹ See Lewinsky-Haapasaari & Norris (1998).

⁴⁰⁰ See Vigalondo et al. (2019).

⁴⁰¹ See Hedenäs (2014c).

- Phascum vlassovii* Laz. ≡ *Tortula vlassovii* (Laz.) Ros & Herrnst.
- Philonotis fontana* (Hedw.) Brid. var. *americana* (Dism.) Flowers ≡ *P. americana* Dism.
- Philonotis fontana* (Hedw.) Brid. var. *caespitosa* (Jur.) Schimp. = *P. fontana* (Hedw.) Brid.⁴⁰²
- Philonotis fontana* (Hedw.) Brid. var. *pumila* (Turner) Brid. = *P. tomentella* Molendo
- Philonotis glaucescens* (Hornsch.) Broth. = *P. uncinata* (Schwägr.) Brid.
- Philonotis gracillima* Ångstr. = *P. uncinata* (Schwägr.) Brid.
- Philonotis muhlenbergii* (Schwägr.) Brid. = *P. marchica* (Hedw.) Brid.
- Physcomitrella patens* (Hedw.) Bruch & Schimp. ≡ *Physcomitrium patens* (Hedw.) Mitt.
- Physcomitrella readeri* (Müll.Hal.) I.G.Stone & G.A. M.Scott ≡ *Physcomitrium readeri* Müll.Hal.
- Physcomitridium readeri* (Müll.Hal.) G.Roth ≡ *Physcomitrium readeri* Müll.Hal.
- Physcomitrium washingtoniense* H.A.Crum & L.E. Anderson = *P. pyriforme* (Hedw.) Brid.
- Plagiobryoides brachyneura* (Kindb.) J.R.Spence ≡ *Bryum brachyneuron* Kindb.
- Plagiobryoides cellularis* (Hook.) J.R.Spence ≡ *Bryum cellulare* Hook.
- Plagiobryoides incrassatolimbata* (Cardot) J.R.Spence ≡ *Bryum incrassatolimbatum* Cardot
- Plagiobryoides renauldii* (Röll) J.R.Spence ≡ *Bryum renauldii* Röll - excluded⁴⁰³
- Plagiobryoides vinosula* (Cardot) J.R.Spence ≡ *Brachymenium vinosulum* Cardot
- Plagiomnium affine* (Blandow) T.J.Kop. - excluded⁴⁰⁴
- Plagiomnium carolinianum* (L.E.Anderson) T.J. Kop. = *P. rhynchophorum* (Harv.) T.J.Kop.
- Plagiomnium medium* (Bruch & Schimp.) T.J.Kop. var. *curvatulum* (Lindb.) H.A.Crum & L.E. Anderson ≡ *P. curvatulum* (Lindb.) Schljakov
- Plagiopus alpinus* (Schwägr.) Hedenäs = *P. serratus* Brid.
- Plagiopus oederi* (Brid.) Limpr. var. *alpinus* (Schwägr.) Dalla Torre & Sarnth. = *P. serratus* Brid.
- Plagiothecium handelii* Broth. ≡ *Ortholimnobia handelii* (Broth.) Schröck & J.T.Wynns
- Plagiothecium platyphyllum* Mönk. = *P. sylvaticum* (Brid.) Schimp.
- Plagiobryum austriacum* (Köckinger, Holyoak & Suanjak) J.R.Spence & Brinda ≡ *Bryum austriacum* Köckinger, Holyoak & Suanjak
- Platydictya confervoides* (Brid.) H.A.Crum ≡ *Serpoleskea confervoides* (Brid.) Loeske
- Platydictya minutissima* (Sull. & Lesq.) H.A.Crum ≡ *Serpoleskea minutissima* (Sull. & Lesq.) W.R. Buck & B.H.Allen
- Platydictya subtilis* (Hedw.) H.A.Crum ≡ *Pseudoamblystegium subtile* (Hedw.) Vanderp. & Hedenäs
- Platyhypnidium pringlei* (Cardot) Broth. ≡ *Donrichardsia pringlei* (Cardot) Huttunen & Ignatov
- Platyhypnidium riparioides sensu auct. Amer. non* (Hedw.) Dixon = *Rhynchostegium aquaticum* Spruce
- Plaubelia sprengelii* (Schwägr.) R.H.Zander var. *stomatodonta* (Cardot) R.H.Zander ≡ *P. stomatodonta* (Cardot) R.H.Zander
- Pleuridium californicum* Grout = *P. acuminatum* Lindb.⁴⁰⁵
- Pleuridium palustre* (Bruch & Schimp.) Bruch & Schimp. ≡ *Cleistocarpidium palustre* (Bruch & Schimp.) Ochyra & Bedn.-Ochyra
- Pleurochaete squarrosa* (Brid.) Lindb. - excluded
- Pohlia bolanderi* (Lesq.) Broth. var. *seriata* A.J.Shaw = *P. bolanderi* (Lesq.) Broth.
- Pohlia elongata* Hedw. var. *greenii* (Brid.) A.J.Shaw = *P. greenii* Brid.
- Polytrichastrum alpinum* (Hedw.) G.L.Sm. var. *fragile* (Bryhn) D.G.Long ≡ *P. fragile* (Bryhn) Schljakov
- Polytrichastrum appalachianum* (L.E.Anderson) G.L. Sm.Merr. ≡ *Polytrichum appalachianum* L.E. Anderson
- Polytrichastrum formosum* (Hedw.) G.L.Sm. ≡ *Polytrichum formosum* Hedw.
- Polytrichastrum formosum* (Hedw.) G.L.Sm. var. *densifolium* (Wilson ex Mitt.) Z.Iwats. & Nog. ≡ *Polytrichum densifolium* Wilson ex Mitt.⁴⁰⁶
- Polytrichastrum longisetum* (Sw. ex Brid.) G.L.Sm. ≡ *Polytrichum longisetum* Sw. ex Brid.
- Polytrichastrum ohioense* (Renauld & Cardot) G.L. Sm. ≡ *Polytrichum ohioense* Renauld & Cardot

⁴⁰² See Hodgetts et al. (2020).⁴⁰³ See Toren et al. (2024).⁴⁰⁴ See McIntosh & Newmaster (2014).⁴⁰⁵ See Yip (2000).⁴⁰⁶ See Ivanova & Ignatov (2017).

- Polytrichastrum pallidisetum* (Funck) G.L.Sm. ≡ *Polytrichum pallidisetum* Funck
- Polytrichastrum sexangulare* (Flörke ex Brid.) G.L.Sm. var. *vulcanicum* (C.E.O.Jensen) G.L.Sm. Merr. = *P. sphaerothecium* (Besch.) J.-P.Frahm⁴⁰⁷
- Polytrichum commune* Hedw. var. *jensenii* (I.Hagen) Mönk. ex Frye ≡ *P. jensenii* I.Hagen
- Polytrichum commune* Hedw. var. *perigoniale* (Michx.) Hampe ≡ *P. perigoniale* Michx.
- Polytrichum formosum* Hedw. var. *densifolium* (Wilson ex Mitt.) Osada ≡ *P. densifolium* Wilson ex Mitt.
- Polytrichum lyallii* (Mitt.) Kindb. ≡ *Polytrichastrum lyallii* (Mitt.) G.L.Sm.
- Polytrichum papillatum* (G.L.Sm.) L.E.Anderson, H.A.Crum & W.R.Buck ≡ *Polytrichastrum papillatum* G.L.Sm.
- Polytrichum sphaerothecium* (Besch.) Müll.Hal. ≡ *Polytrichastrum sphaerothecium* (Besch.) J.-P.Frahm
- Porotrichum bigelovii* (Sull.) Kindb. ≡ *Dannorisia bigelovii* (Sull.) Enroth⁴⁰⁸
- Porotrichum vancouveriense* (Kindb.) H.A.Crum ≡ *Bryolawtonia vancouveriensis* (Kindb.) D.H.Norris & Enroth
- Pottia arizonica* Wareham = *Microbryum starckeana* (Hedw.) R.H.Zander
- Pottia arizonica* Wareham var. *mucronulata* Wareham = *Microbryum muticum* (Venturi) Cl.Schneid., Th.Schneid. & Mahévas
- Pottia bryoides* (Dicks.) Mitt. ≡ *Tortula protobryoides* R.H.Zander
- Pottia davalliana* (Sm.) C.E.O.Jensen ≡ *Microbryum davallianum* (Sm.) R.H.Zander
- Pottia intermedia* (Turner) Fürnr. = *Tortula caucasica* Lindb.
- Pottia lanceolata* (Hedw.) Müll.Hal. = *Tortula lindbergii* Kindb. ex Broth.
- Pottia nevadensis* Cardot & Thér. ≡ *Tortula nevadensis* (Cardot & Thér.) R.H.Zander
- Pottia starckeana* (Hedw.) Müll.Hal. ≡ *Microbryum starckeanum* (Hedw.) R.H.Zander
- Pottia truncata* (Hedw.) Bruch & Schimp. ≡ *Tortula truncata* (Hedw.) Mitt.
- Pottia wilsonii* (Hook.) Bruch & Schimp. ≡ *Tortula wilsonii* (Hook.) R.H.Zander - excluded⁴⁰⁹
- Pseudoblindia* Fedosov, M.Stech & Ignatov = *Kiaeria* I.Hagen⁴¹⁰
- Pseudoblindia falcata* (Hedw.) Fedosov, M.Stech & Ignatov ≡ *Kiaeria falcata* (Hedw.) I.Hagen
- Pseudocalliergon angustifolium* Hedenäs ≡ *Drepanocladus angustifolius* (Hedenäs) Hedenäs & Rosborg
- Pseudocalliergon brevifolium* (Lindb.) Hedenäs ≡ *Drepanocladus brevifolius* (Lindb.) Warnst.
- Pseudocalliergon trifarium* (F.Weber & D.Mohr) Loeske ≡ *Drepanocladus trifarius* (F.Weber & D.Mohr) Broth.
- Pseudocalliergon turgescens* (T.Jensen) Loeske ≡ *Drepanocladus turgescens* (T.Jensen) Broth.
- Pseudocrossidium aureum* (E.B.Bartram) R.H.Zander = *P. arenicola* (Dusén) M.J.Cano
- Pseudocrossidium crinitum* (Schultz) R.H.Zander - excluded⁴¹¹
- Pseudocrossidium revolutum* (Brid.) R.H.Zander - excluded⁴¹²
- Pseudocrossidium revolutum* (Brid.) R.H.Zander var. *obtusulum* (Lindb.) B.C.Tan, R.H.Zander & T.Taylor ≡ *P. obtusulum* (Lindb.) H.A.Crum & L.E.Anderson
- Pseudoleskea atricha* (Kindb.) Kindb. ≡ *Lescuraea atricha* (Kindb.) E.Lawton
- Pseudoleskea baileyi* Best & Grout ≡ *Lescuraea baileyi* (Best & Grout) E.Lawton
- Pseudoleskea incurvata* (Hedw.) Loeske ≡ *Lescuraea incurvata* (Hedw.) E.Lawton
- Pseudoleskea incurvata* (Hedw.) Loeske var. *gigantea* (E.Lawton) H.A.Crum, Steere & L.E.Anderson ≡ *Lescuraea incurvata* (Hedw.) E.Lawton var. *gigantea* E.Lawton
- Pseudoleskea incurvata* (Hedw.) Loeske var. *tenuiretis* (Culm.) H.A.Crum, Steere & L.E.Anderson ≡ *Lescuraea incurvata* (Hedw.) E.Lawton var. *tenuiretis* (Culm.) E.Lawton
- Pseudoleskea patens* (Lindb.) Kindb. ≡ *Lescuraea patens* (Lindb.) Arnell & C.E.O.Jensen
- Pseudoleskea radicata* (Mitt.) Macoun & Kindb. var. *radicata* ≡ *Lescuraea radicata* (Mitt.) Mönk. var. *radicata*

⁴⁰⁷ See Ivanova & Ignatov (2017).

⁴⁰⁸ See Enroth et al. (2019).

⁴⁰⁹ See Zander (2007f).

⁴¹⁰ See Brinda & Fedosov (2023).

⁴¹¹ See Cano et al. (2016).

⁴¹² See Zander (2007e).

- Pseudoleskea radicata* (Mitt.) Macoun & Kindb. var. *compacta* Best \equiv *Lescuraea radicata* (Mitt.) Mönk. var. *compacta* (Best) E.Lawton
- Pseudoleskea radicata* (Mitt.) Macoun & Kindb. var. *denudata* (Kindb.) Wijk & Margad. \equiv *Lescuraea radicata* (Mitt.) Mönk. var. *denudata* (Kindb.) E.Lawton
- Pseudoleskea radicata* (Mitt.) Macoun & Kindb. var. *pallida* (Best) H.A.Crum, Steere & L.E.Anderson \equiv *Lescuraea radicata* (Mitt.) Mönk. var. *pallida* (Best) E.Lawton
- Pseudoleskea saviana* (DeNot.) Latzel \equiv *Lescuraea saviana* (De Not.) E.Lawton
- Pseudoleskea stenophylla* Renauld & Cardot \equiv *Lescuraea stenophylla* (Renauld & Cardot) Kindb.
- Pseudoleskea tribulosa* Shevock & W.R.Buck \equiv *Lescuraea tribulosa* (Shevock & W.R.Buck) W.R. Buck & Goffinet
- Pseudoleskeella sibirica* (Arnell) P.S.Wilson & D.H. Norris = *P. rupestris* (Berggr.) Hedenäs & L. Söderstr.
- Pseudosymblepharis angustata* (Mitt.) Hilp. \equiv *Chionoloma angustatum* (Mitt.) M.Menzel
- Pseudotaxiphyllum distichaceum* (Mitt.) Z.Iwats. \equiv *Longiella distichacea* (Mitt.) J.T.Wynns - excluded⁴¹³
- Pseudotaxiphyllum distichaceum sensu auct. Amer. non* (Mitt.) Z.Iwats. = *P. subfalcatum* (Austin) X.Q.Li, Q.Zuo & Y.F.Wang
- Pseudotaxiphyllum homomallifolium* (Redf.) Ireland \equiv *Redfearnia homomallifolia* (Redf.) J.T.Wynns
- Pterogonium gracile* (Hedw.) Sm. \equiv *Nogopterium gracile* (Hedw.) Crosby & W.R.Buck
- Pterygoneurum californicum* H.A.Crum = *P. subsessile* (Brid.) Jur.
- Ptychomitrium leibergii* Best = *P. sinense* (Mitt.) A. Jaeger
- Ptychostomum acutiforme* (Limpr. ex Ryan) J.R. Spence \equiv *Bryum acutiforme* Limpr. ex Ryan.
- Ptychostomum archangelicum* (Bruch & Schimp.) J. R.Spence \equiv *Bryum archangelicum* Bruch & Schimp.
- Ptychostomum arcticum* (R.Br.) J.R.Spence \equiv *Bryum arcticum* R.Br.
- Ptychostomum austriacum* (Köckinger, Holyoak & Suanjak) D.Bell & Holyoak \equiv *Bryum austriacum* Köckinger, Holyoak & Suanjak
- Ptychostomum axel-blyttii* (H.Philb.) J.R.Spence = *Bryum calophyllum* R.Br.
- Ptychostomum badium* (Brid.) J.R.Spence = *Bryum imbricatum* Müll.Hal.
- Ptychostomum bimum* (Schreb.) J.R.Spence \equiv *Bryum bimum* (Schreb.) Turner
- Ptychostomum calophyllum* (R.Br.) J.R.Spence \equiv *Bryum calophyllum* R.Br.
- Ptychostomum cernuum* (Hedw.) Hornsch. = *Bryum uliginosum* (Brid.) Bruch & Schimp.
- Ptychostomum compactum* Hornsch. = *Bryum algovicum* Sendtn. ex Müll.Hal.
- Ptychostomum creberrimum* (Taylor) J.R.Spence & H.P.Ramsay \equiv *Bryum creberrimum* Taylor
- Ptychostomum cryophilum* (Mårtensson) J.R.Spence \equiv *Bryum cryophilum* Mårtensson
- Ptychostomum cyclophyllum* (Schwägr.) J.R.Spence \equiv *Bryum cyclophyllum* Schwägr.
- Ptychostomum inclinatum* (Sw. ex Brid.) J.R.Spence = *Bryum amblyodon* Müll.Hal.
- Ptychostomum intermedium* (Brid.) J.R.Spence \equiv *Bryum intermedium* (Brid.) Turton
- Ptychostomum knowltonii* (Barnes) J.R.Spence \equiv *Bryum knowltonii* Barnes
- Ptychostomum lonchocaulon* (Müll.Hal.) J.R.Spence \equiv *Bryum lonchocaulon* Müll.Hal.
- Ptychostomum longisetum* (Blandow ex Schwägr.) J.R. Spence \equiv *Bryum longisetum* Blandow ex Schwägr.
- Ptychostomum marratii* (Hook. & Wilson) J.R. Spence \equiv *Bryum marratii* Hook. & Wilson
- Ptychostomum meesioides* (Kindb.) J.R.Spence \equiv *Bryum meesioides* Kindb.
- Ptychostomum neodamense* (Itzigs.) J.R.Spence \equiv *Bryum neodamense* Itzigs.
- Ptychostomum nitidulum* (Lindb.) J.R.Spence \equiv *Bryum nitidulum* Lindb.
- Ptychostomum pacificum* J.R.Spence & Shevock \equiv *Bryum pacificum* (J.R.Spence & Shevock) W.R. Buck & Goffinet
- Ptychostomum pallens* (Sw.) J.R.Spence \equiv *Bryum pallens* Sw.
- Ptychostomum pallescens* (Schleich. ex Schwägr.) J.R. Spence \equiv *Bryum pallescens* Schleich. ex Schwägr.
- Ptychostomum pendulum* Hornsch. = *Bryum algovicum* Sendtn. ex Müll.Hal.
- Ptychostomum pseudotriquetrum* (Hedw.) J.R.Spence & H.P.Ramsay ex Holyoak & N.Pedersen \equiv

⁴¹³ See Li et al. (2015) and Wynns (2020).

- Bryum pseudotriquetrum* (Hedw.) G.Gaertn., B. Meyer & Scherb.
Ptychostomum reedii (H.Rob.) J.R.Spence ≡ *Bryum reedii* H.Rob.
Ptychostomum rutilans (Brid.) J.R.Spence ≡ *Bryum rutilans* Brid.
Ptychostomum salinum (I.Hagen ex Limpr.) J.R. Spence ≡ *Bryum salinum* I.Hagen ex Limpr.
Ptychostomum schleicheri (Schwägr.) J.R.Spence ≡ *Bryum schleicheri* Schwägr.
Ptychostomum subneodamense (Kindb.) J.R.Spence ≡ *Bryum subneodamense* Kindb.
Ptychostomum turbinatum (Hedw.) J.R.Spence ≡ *Bryum turbinatum* (Hedw.) Turner
Ptychostomum warneum (Schwägr. ex Steud.) J.R. Spence ≡ *Bryum warneum* (Schwägr. ex Steud.) Brid.
Ptychostomum weigeli (Biehler) J.R.Spence ≡ *Bryum weigeli* Biehler
Ptychostomum wrightii (Sull.) J.R.Spence ≡ *Bryum wrightii* Sull.
Pylaisia condensata (Mitt.) A.Jaeger - excluded⁴¹⁴
Pylaisiella intricata (Hedw.) Grout ≡ *Pylaisia intricata* (Hedw.) Schimp.
Pylaisiella polyantha (Hedw.) Grout ≡ *Pylaisia polyantha* (Hedw.) Schimp.
Pylaisiella selwynii (Kindb.) H.A.Crum, Steere & L. E.Anderson ≡ *Pylaisia selwynii* Kindb.
Pylaisiella steerei Ando & Higuchi ≡ *Pylaisia steerei* (Ando & Higuchi) Ignatov
Racomitrium aquaticum (Brid. ex Schrad.) Brid. - excluded
Racomitrium aquaticum sensu auct. Amer. non (Brid. ex Schrad.) Brid. = *R. ryszardii* Bedn.-Ochyra⁴¹⁵
Rhynchostegium riparioides sensu auct. Amer. non (Hedw.) Cardot = *R. aquaticum* Spruce
Rhytidadelphus triquetrus (Hedw.) Warnst. ≡ *Hylocomiadelphus triquetrus* (Hedw.) Ochyra & Stebel
Roellia roellii (Broth.) A.L.Andrews ex H.A.Crum ≡ *Roellobryon roellii* (Broth.) Ochyra
Rosulabryum andersonii (H.A.Crum) J.R.Spence ≡ *Brachymenium andersonii* H.A.Crum

⁴¹⁴ See Ignatova et al. (2020b).

⁴¹⁵ See Bednarek-Ochyra (2000).

- Rosulabryum andicola* (Hook.) Ochyra ≡ *Bryum andicola* Hook.
Rosulabryum biforme (R.S.Williams) J.R.Spence ≡ *Bryum biforme* R.S.Williams
Rosulabryum bornholmense (Wink. & R.Ruthe) J.R. Spence ≡ *Bryum bornholmense* Wink. & R. Ruthe
Rosulabryum canariense (Brid.) Ochyra ≡ *Bryum canariense* Brid.
Rosulabryum capillare (Hedw.) J.R.Spence ≡ *Bryum capillare* Hedw.
Rosulabryum elegans (Nees ex Brid.) Ochyra ≡ *Bryum elegans* Nees ex Brid.
Rosulabryum erythroloma (Kindb.) J.R.Spence ≡ *Bryum erythroloma* (Kindb.) Syed
Rosulabryum flaccidum (Brid.) J.R.Spence ≡ *Bryum flaccidum* Brid.
Rosulabryum gemmascens (Kindb.) J.R.Spence = *Bryum sanguilentum* Renault & Cardot
Rosulabryum laevifilum (Syed) Ochyra = *Bryum moravicum* Podp.
Rosulabryum pseudocapillare (Besch.) Ochyra ≡ *Bryum pseudocapillare* Besch.
Rosulabryum rubens (Mitt.) J.R.Spence ≡ *Bryum rubens* Mitt.
Rosulabryum sanguilentum (Renauld & Cardot) J.R. Spence & Brinda ≡ *Bryum sanguilentum* Renault & Cardot
Rosulabryum torquescens (Bruch & Schimp.) J.R. Spence ≡ *Bryum torquescens* Bruch & Schimp.
Sanionia georgicouncinata (Müll.Hal.) Ochyra & Hedenäs - excluded
Sanionia georgicouncinata sensu auct. Amer. non (Müll.Hal.) Ochyra & Hedenäs = *S. nivalis* Hedenäs⁴¹⁶
Sanionia uncinata (Hedw.) Loeske var. *symmetrica* (Renauld & Cardot) H.A.Crum & L.E.Anderson ≡ *S. symmetrica* (Renauld & Cardot) Wheldon
Sarmentypnum exannulatum (Schimp.) Hedenäs ≡ *Warnstorfia exannulata* (Schimp.) Loeske
Sarmentypnum pseudosarmentosum (Cardot & Thér.) Hedenäs ≡ *Warnstorfia pseudosarmentosa* (Cardot & Thér.) Tuom. & T.J.Kop.
Sarmentypnum sarmentosum (Wahlenb.) Tuom. & T.J.Kop. ≡ *Warnstorfia sarmentosa* (Wahlenb.) Hedenäs

⁴¹⁶ See Hedenäs (2012).

- Sarmentypnum trichophyllum* (Warnst.) Hedenäs ≡ *Warnstorfia trichophylla* (Warnst.) Tuom. & T.J. Kop.
- Sarmentypnum tundrae* (Arnell) Hedenäs ≡ *Warnstorfia tundrae* (Arnell) Loeske
- Schistidium apocarpum* (Hedw.) Bruch & Schimp. subsp. *canadense* (Dupr.) H.H.Blom ex B.H. Allen & Pursell ≡ *S. canadense* (Dupr.) Ignatova & H.H.Blom
- Schofieldiella micans* (Mitt.) W.R.Buck ≡ *Hageniella micans* (Mitt.) B.C.Tan & Y.Jia
- Scleropodium cespitans* (Müll.Hal.) L.F.Koch var. *sublaeve* (Renauld & Cardot) Wijk & Margad. = *S. cespitans* (Müll.Hal.) L.F.Koch⁴¹⁷
- Scleropodium touretii* (Brid.) L.F.Koch var. *colpophyllum* (Sull.) E.Lawton ex H.A.Crum = *S. touretii* (Brid.) L.F.Koch
- Scopelophila cataractae* (Mitt.) Broth. ≡ *Merceyopsis cataractae* (Mitt.) Brinda, Ignatov & Fedosov
- Sematophyllum subpinnatum* (Brid.) E.Britton ≡ *Brittonodoxa subpinnata* (Brid.) W.R.Buck, P.E. A.S.Câmara & Carv.-Silva
- Sphagnum andrusii* (H.A.Crum) Flatberg = *S. miyabeum* Warnst.
- Sphagnum atlanticum* R.E.Andrus = *S. torreyanum* Sull.⁴¹⁸
- Sphagnum bartlettianum* Warnst. = *S. rubellum* Wilson⁴¹⁹
- Sphagnum bergianum* R.E.Andrus = *S. subfulvum* Sjörs⁴²⁰
- Sphagnum brevifolium* (Lindb.) Röhl = *S. fallax* (H. Klinggr.) H.Klinggr.⁴²¹
- Sphagnum crispum* R.E.Andrus = *S. subsecundum* Nees
- Sphagnum denticulatum* Brid. - excluded
- Sphagnum fimbriatum* Wilson subsp. *concinnum* (Berggr.) Flatberg & Frisvoll ≡ *S. concinnum* (Berggr.) Flatberg⁴²²
- Sphagnum henryense* Warnst. = *S. palustre* L.⁴²³
- Sphagnum inundatum* Russow - excluded⁴²⁴
- Sphagnum isoviitae* Flatberg = *S. fallax* (H.Klinggr.) H.Klinggr.⁴²⁵
- Sphagnum macrophyllum* Brid. var. *burinense* Maass = *S. macrophyllum* Brid.
- Sphagnum macrophyllum* Brid. var. *floridanum* Austin = *S. cribrosum* Lindb.
- Sphagnum majus* (Russow) C.E.O.Jensen ssp. *norvegicum* Flatberg = *S. majus* (Russow) C.E.O.Jensen
- Sphagnum mcqueenii* R.E.Andrus = *S. torreyanum* Sull.
- Sphagnum nitidum* Warnst. = *S. subfulvum* Sjörs
- Sphagnum rubroflexuosum* R.E.Andrus = *S. flexuosum* Dozy & Molk.⁴²⁶
- Sphagnum schofieldii* H.A.Crum = *S. quinquefarium* (Lindb. ex Braithw.) Warnst.
- Sphagnum sitchense* R.E.Andrus = *S. capillifolium* (Ehrh.) Hedw.⁴²⁷
- Sphagnum subobesum* Warnst. = *S. miyabeum* Warnst.⁴²⁸
- Sphagnum subsecundum* Nees var. *andrusii* H.A. Crum = *S. miyabeum* Warnst.
- Sphagnum subtile* (Russow) Warnst. = *S. capillifolium* (Ehrh.) Hedw.⁴²⁹
- Sphagnum viride* Flatberg = *S. cuspidatum* Ehrh. ex Hoffm.⁴³⁰
- Sphagnum wilfii* H.A.Crum = *S. rubellum* Wilson⁴³¹
- Splachnum luteum* Hedw. var. *melanocaulon* Wahlenb. ≡ *Splachnum melanocaulon* (Wahlenb.) Schwägr.
- Steerecleus serrulatus* (Hedw.) H.Rob. ≡ *Rhynchostegium serrulatum* (Hedw.) Austin
- Straminergon stramineum* (Dicks. ex Brid.) Hedenäs ≡ *Warnstorfia straminea* (Dicks. ex Brid.) W.R. Buck & Goffinet
- Streptopogon amphidiaceus* (Müll.Hal.) M.T.Gallego & M.J.Cano ≡ *Syntrichia amphidiacea* (Müll. Hal.) R.H.Zander
- Syntrichia cainii* (H.A.Crum & L.E.Anderson) R.H. Zander ≡ *S. norvegica* F.Weber var. *cainii* (H.A. Crum & L.E.Anderson) W.A.Kramer
- Syntrichia papillosissima* (Copp.) Loeske = *S. ruralis* (Hedw.) F.Weber & D.Mohr subsp. *hirsuta* (Venturi) Düll

⁴¹⁷ See Carter (2014).⁴¹⁸ See Shaw et al. (2009).⁴¹⁹ See Imwattana et al. (2024).⁴²⁰ See Shaw et al. (2009).⁴²¹ See Duffy et al. (2020).⁴²² See Shaw et al. (2012a).⁴²³ See Karlin et al. (2010).⁴²⁴ See Shaw et al. (2012b).⁴²⁵ See Duffy et al. (2020).⁴²⁶ See Duffy et al. (2020).⁴²⁷ See Imwattana et al. (2024).⁴²⁸ See Shaw et al. (2015).⁴²⁹ See Imwattana et al. (2024).⁴³⁰ See Robinson et al. (2024).⁴³¹ See Imwattana et al. (2024).

- Tetraplodon angustatus* (Hedw.) Bruch & Schimp. ≡
Voitia angustata (Hedw.) W.R.Buck & Goffinet
- Tetraplodon mnioides* (Hedw.) Bruch & Schimp. ≡
Voitia mnioides (Hedw.) W.R.Buck & Goffinet
- Tetraplodon pallidus* I.Hagen ≡ *Voitia pallida* (I.
Hagen) W.R.Buck & Goffinet
- Tetraplodon paradoxus* (R.Br.) I.Hagen ≡ *Voitia
paradoxa* (R.Br.) W.R.Buck & Goffinet
- Tetraplodon urceolatus* (Hedw.) Bruch & Schimp. ≡
Voitia urceolata (Hedw.) W.R.Buck & Goffinet
- Tetrodontium brownianum* (Dicks.) Schwägr. var.
ovatum (Funck) Wijk & Margad. ≡ *T. ovatum*
(Funck) Schwägr.
- Thamnobryum alleghaniense* (Müll.Hal.) Nieuwl. =
T. subserratum (Hook. ex Harv.) Nog. & Z.
Iwats.
- Thuidium delicatulum* (Hedw.) Schimp. var. *radicans*
(Kindb.) H.A.Crum, Steere & L.E.Anderson = *T.
assimile* (Mitt.) A.Jaeger
- Thuidium philibertii* Limpr. = *T. assimile* (Mitt.) A.
Jaeger
- Timmia megapolitana* Hedw. var. *bavarica* (Hessl.)
Brid. ≡ *T. bavarica* Hessl.
- Timmia norvegica* J.E.Zetterst. var. *excurrens* Bryhn =
T. comata Lindb. & Arnell
- Torrentaria riparioides* (Hedw.) Ochyra – excluded
- Tortella inclinata* (R.Hedw.) Limpr. var. *densa* (Lorentz
& Molendo) Limpr. ≡ *T. densa* (Lorentz &
Molendo) Crundw. & Nyholm
- Tortella tortelloides* (S.W.Greene) H.Rob. = *T. alpi-
cola* Dixon
- Tortella tortuosa* (Hedw.) Limpr. var. *arctica*
(Arnell) Broth. ≡ *T. arctica* (Arnell) Crundw.
& Nyholm
- Tortella tortuosa* (Hedw.) Limpr. var. *fragilifolia*
(Jur.) Limpr. ≡ *T. nidita* (Lindb.) Broth. var.
fragilifolia (Jur.) Köckinger & Hedenäs
- Tortula ammonsiana* H.A.Crum & L.E.Anderson ≡
Syntrichia ammonsiana (H.A.Crum & L.E.Anderson)
Ochyra
- Tortula amphidiacea* (Müll.Hal.) Broth. ≡ *Syntrichia
amphidiacea* (Müll.Hal.) R.H.Zander
- Tortula bartramii* Steere ≡ *Syntrichia bartramii*
(Steere) R.H.Zander
- Tortula cainii* H.A.Crum & L.E.Anderson ≡ *Syntrichia
norvegica* F.Weber var. *cainii* (H.A.Crum & L.E.
Anderson) W.A.Kramer
- Tortula caninervis* (Mitt.) Broth. ≡ *Syntrichia cani-
nervis* Mitt.
- Tortula chisosa* Magill, Delgad. & L.R.Stark ≡
Syntrichia chisosa (Magill, Delgad. & L.R.Stark)
R.H.Zander
- Tortula fragilis* Taylor ≡ *Syntrichia fragilis* (Taylor)
Ochyra
- Tortula laevipila* (Brid.) Schwägr. ≡ *Syntrichia laevi-
pila* Brid.
- Tortula laevipila* (Brid.) Schwägr. var. *meridionalis*
(Schimp.) Wijk & Margad. = *Syntrichia laevipila*
Brid.⁴³²
- Tortula lanceola* R.H.Zander = *T. lindbergii* Kindb.
ex Broth.
- Tortula latifolia* Bruch ex Hartm. ≡ *Syntrichia latifo-
lia* (Bruch ex Hartm.) Huebener
- Tortula modica* R.H.Zander = *T. caucasica* Lindb.
- Tortula norvegica* (F.Weber) Wahlenb. ex Lindb. ≡
Syntrichia norvegica F.Weber
- Tortula obtusissima* (Müll.Hal.) Mitt. ≡ *Syntrichia
obtusissima* (Müll.Hal.) R.H.Zander
- Tortula pagorum* (Milde) De Not. ≡ *Syntrichia
pagorum* (Milde) J.J.Amann
- Tortula papillosa* Wilson ex Spruce ≡ *Syntrichia
papillosa* (Wilson ex Spruce) Spruce
- Tortula princeps* De Not. ≡ *Syntrichia princeps* (De
Not.) Mitt.
- Tortula rhizophylla* (Sakurai) Z.Iwats. & K.Saito =
Chenia leptophylla (Müll.Hal.) R.H.Zander
- Tortula ruralis* (Hedw.) G.Gaertn., B.Mey. & Scherb.
≡ *Syntrichia ruralis* (Hedw.) F.Weber & D.Mohr
- Tortula scotteri* R.H.Zander & Steere = *Hilpertia
velenovskiyi* (Schiffn.) R.H.Zander
- Tortula sinensis* (Müll.Hal.) Broth. ≡ *Syntrichia
sinensis* (Müll.Hal.) Ochyra
- Tortula stanfordensis* Steere ≡ *Hennediella stanfor-
densis* (Steere) Blockeel
- Tortula virescens* (De Not.) De Not. ≡ *Syntrichia
virescens* (De Not.) Boros
- Trachybryum megaptilum* (Sull.) W.B.Schofield ≡
Homalothecium megaptilum (Sull.) H.Rob.
- Trachyxiphium heteroicum* (Cardot) W.R.Buck = *T.
hypnaceum* (Müll.Hal.) W.R.Buck
- Trematodon boasii* W.B.Schofield = *T. asanoi* Tuzibe
- Trichodon cylindricus* (Hedw.) Schimp. var. *oblongus*
(Lindb.) Podp. ≡ *Trichodon oblongus* Lindb.
- Trichostomopsis brevifolia* E.B.Bartram = *T. austral-
asiae* (Grev. & Hook.) H.Rob.

⁴³² See Gallego et al. (2004).

Trichostomum arcticum Kaal. = *Tortella spitsbergen-
sis* (Bizot & Thér.) O.Werner, Köckinger & Ros
Trichostomum brittonianum R.H.Zander - excluded⁴³³
Trichostomum molariforme R.H.Zander = *T. portori-
cense* H.A.Crum & Steere
Trichostomum recurvifolium (Taylor) R.H.Zander ≡
Chionoloma recurvifolium (Taylor) M.Alonso,
M.J.Cano & J.A.Jiménez
Trichostomum spirale Grout = *Chionoloma stenocar-
pum* (Thér.) M.Alonso, M.J.Cano & J.A.Jiménez
Trichostomum tenuirostre (Hook. & Taylor) Lindb.
≡ *Chionoloma tenuirostre* (Hook. & Taylor) M.
Alonso, M.J.Cano & J.A.Jiménez
Trichostomum tenuirostre (Hook. & Taylor) Lindb.
var. *gemmaiparum* (Schimp.) R.H.Zander =
Chionoloma orthodontum (Müll.Hal.) M.Alonso,
M.J.Cano & J.A.Jiménez⁴³⁴
Tuerckheimia angustifolia (K.Saito) R.H.Zander =
T. svihlae (E.B.Bartram) R.H.Zander
Ulota japonica (Sull. & Lesq.) Mitt. - excluded⁴³⁵
Ulota phyllantha Brid. ≡ *Plenogemma phyllantha*
(Brid.) Sawicki, Plášek & Ochyra
Venturiella sinensis (Venturi) Müll.Hal. var. *angus-
tiannulata* D.G.Griffin & Sharp ≡ *V. sinensis*
(Venturi) Müll.Hal. subsp. *angustiannulata* (D.
G.Griffin & Sharp) Pursell
Vinealobryum murrayae (Otnyukova) R.H.Zander =
Husnotiella fragilicuspis (Broth.) J.A.Jiménez &
M.J.Cano
Weissia andersoniana R.H.Zander = *W. ligulifolia*
(E.B.Bartram) Grout
Weissia condensa (Voit) Lindb. - excluded⁴³⁶
Weissia flavescens (E.Britton) W.D.Reese = *Trichostomum
brittonianum* R.H.Zander - excluded
Weissia glauca E.B.Bartram = *W. ligulifolia* (E.B.
Bartram) Grout
Weissia hedwigii H.A.Crum = *W. brachycarpa* (Nees
& Hornsch.) Jur.
Zygodon apiculatus Redf. = *Z. rupestris* Schimp. ex
Lorentz
Zygodon menziesii (Schwägr.) Arn. ≡ *Codonoblepharon
menziesii* Schwägr.
Zygodon viridissimus (Dicks.) Brid. var. *dentatus*
Limpr. ≡ *Z. dentatus* (Limpr.) Breidl. ex Gams

⁴³³ See Zander (2007g).

⁴³⁴ See Alonso-García et al. (2019).

⁴³⁵ See Vitt (2014).

⁴³⁶ See Zander (1993, 2007h).

Zygodon viridissimus (Dicks.) Brid. var. *rupestris*
Lindb. ex C.Hartm. ≡ *Z. rupestris* Schimp. ex
Lorentz

ACKNOWLEDGMENTS

We hesitantly thank colleagues because we seldom completely followed their suggestions! Nevertheless, we are grateful for their ideas and collaboration. Initially, Bruce Allen was going to work on this project but decided to withdraw because he would have had to make too many compromises. You can still see his influence throughout this checklist. Lars Hedenäs offered suggestions for the Amblystegiaceae s.l. We hope he won't be too disappointed that we were more conservative in recognition of genera. Jon Shaw gave us his suggestions for the *Sphagnum* list. Jim Shevock read over the list to help make sure we didn't omit any taxa recently reported for North America, most of which are from the West. We made abundant use of TROPICOS and bryonames.org (Brinda & Atwood 2024), both for nomenclature and for tracking down references. We appreciate the comments sent to us by John Atwood, John Brinda, María Cano, Vladimir Fedosov, Joe Rohrer, Dale Vitt and Richard Zander. The review by David Toren helped us to avoid missing recent North American additions. We apologize to those we may have contacted but have forgotten; this project has gone on for way too long.

LITERATURE CITED

- Afonina, O. M. 2020. *Oncophorus integerrimus* Hedenäs. In: Ellis, L. T. et al., New national and regional bryophyte records, 62. *Journal of Bryology* 42: 201.
- Afonina, O. M. & A. Breen. 2009. *Dicranum dispersum* (Dicranaceae) and *Sciuro-hypnum ornellanum* (Brachytheciaceae), new to North America. *The Bryologist* 112: 268–272.
- Allen, B. H. 1991. A new species of *Fontinalis* (Musci: Fontinalaceae) in North America. *The Bryologist* 94: 201–202.
- Allen, B. H. 2002. Moss flora of Central America, Part 2. Encalypta-ceae–Orthotrichaceae. *Monographs in Systematic Botany from the Missouri Botanical Garden* 90: i–viii, 1–699.
- Allen, B. H. 2005. Maine mosses: Sphagnaceae–Timmiaceae. *Memoirs of the New York Botanical Garden* 93: i–xii, 1–420.
- Allen, B. H. 2010. Hedwigiaceae. In: Moss flora of Central America. Part 3. Anomodontaceae–Symphyodontaceae. *Monographs in Systematic Botany from the Missouri Botanical Garden* 117: 16–30.
- Allen, B. H. 2014. Maine mosses: Drummondia-ceae–Polytrichaceae. *Memoirs of the New York Botanical Garden* 111: i–xiii, 1–607.
- Allen, B. H. 2018. Moss flora of Central America, Part 4. Fabronia-ceae–Polytrichaceae. *Monographs in Systematic Botany from the Missouri Botanical Garden* 132: i–x, 1–830.
- Allen, B. H. 2022. *Bryoerythrophyllum wallichii* new to North America. *Evansia* 39: 66–68.
- Allen, B. H. & R. E. Magill. 2004. *Leptodictyum wallacei* (Musci: Amblystegiaceae), a new species from Texas, USA. *Novon* 14: 377–379.
- Allred, K. W., R. Kleinman & K. S. Blisard. 2024. Flora Neomexicana IV: Bryophytes (Mosses & Liverworts). Lulu Press, Raleigh, NC. 559 pp.
- Alonso-García, M., J. A. Jiménez & M. J. Cano. 2019. Taxonomic revision of *Chionoloma* (Pottiaceae, Bryophyta). *Annals of the Missouri Botanical Garden* 104: 563–632.

- Alonso-García, M., J. A. Jiménez, S. Nylinder, L. Hedenäs & M. J. Cano. 2016. Disentangling generic limits in *Chionoloma*, *Oxystegus*, *Pachyneurospis* and *Pseudosymblepharis* (Bryophyta: Pottiaceae): an inquiry into their phylogenetic relationships. *Taxon* 65: 3–18.
- Anderson, L. E. 1990. A checklist of *Sphagnum* in North America north of Mexico. *The Bryologist* 93: 500–501.
- Anderson, L. E., H. A. Crum & W. R. Buck. 1990. List of the mosses of North America north of Mexico. *The Bryologist* 93: 448–499.
- Ando, H. 1994. A revision of *Hypnum aemulans* Limpr., with records of a new species, *H. holmenii* (sect. *Hamulosa*) from northern North America, Greenland and Finland. *Hikobia* 11: 363–370.
- Arts, T. 1998. A revision of the moss genus *Gymnostomiella* Fleisch. *Journal of Bryology* 20: 411–427.
- Arts, T., A. C. Crundwell & H. L. K. Whitehouse. 1995. *Bryum valparaisense* Thér. an earlier name for *B. pyriferum* Crundw. & Whiteh. *Journal of Bryology* 18: 797–801.
- Atwood, J. J. & J. C. Brinda. 2020. *Rhaphidostegium planifolium* is a synonym of *Calliergonella lindbergii* var. *lindbergii*, and a note on the variety *americana*. *Missouriensis* 38: 13–18.
- Austin, C. F. 1880. Bryological notes. *Bulletin of the Torrey Botanical Club* 7(2): 15–16.
- Bechteler, J., G. F. Peñaloza-Bojacá, D. Bell, G. Burleigh, S. F. McDaniel, E. C. Davis, E. B. Sessa, A. C. Bippus, D. C. Cargill, S. Chantanaorrapint, I. Draper, L. Endara, L. L. Forrest, R. Garilleti, S. W. Graham, S. Huttunen, J. A. Jáuregui-Lazo, F. Lara, J. Larraín, L. R. Lewis, D. G. Long, D. Quandt, K. S. Renzaglia, A. Schäfer-Verwimp, G. E. Lee, A. M. Sierra, M. von Konrat, C. E. Zartman, M. R. Pereira, B. Goffinet & J. C. Villarreal A. 2023. Comprehensive phylogenomic time tree of bryophytes reveals deep relationships and uncovers gene incongruences in the last 500 million years of diversification. *American Journal of Botany* 110(11): e16249 [1–20].
- Bednarek-Ochyra, H. 1995. The genus *Racomitrium* (Musci, Grimmiaceae) in Poland: taxonomy, ecology and phytogeography. *Fragmenta Floristica et Geobotanica, Series Polonica* 2: 1–307.
- Bednarek-Ochyra, H. 1999. *Racomitrium aduncooides* (Musci, Grimmiaceae), a remarkable new species from eastern North America. *Fragmenta Floristica et Geobotanica* 44: 277–285.
- Bednarek-Ochyra, H. 2000. *Racomitrium rysardii* (Musci, Grimmiaceae), a new hydrophilous species from the Pacific Northwest with comments on *Racomitrium aquaticum* in North America. *Cryptogamie Bryologie* 21: 275–284.
- Bednarek-Ochyra, H. 2004. *Codriophorus corrugatus* (Bryopsida, Grimmiaceae), a new species from East Asia and southern Alaska. *The Bryologist* 107: 377–384.
- Bednarek-Ochyra, H. & R. Ochyra. 2000. *Racomitrium norrisii* sp. nova (Grimmiaceae, Bryopsida), endemic of western North America. *Annales Botanici Fennici* 37: 235–241.
- Bednarek-Ochyra, H. & R. Ochyra. 2007. *Codriophorus*. In *Flora of North America* Editorial Committee (eds.), *Flora North America* 27: 295–305. Oxford University Press, New York.
- Bednarek-Ochyra, H., J. Sawicki, R. Ochyra, M. Szczecińska & V. Plášek. 2015. *Dilutineuron*, a new moss genus of the subfamily Racomitrioidae (Grimmiaceae, Bryophyta). *Acta Musei Silesiae, Scientiae Naturales* 64: 163–168.
- Beike, A. K., M. von Stackelberg, M. Schallenberg-Rüdinger, S. T. Hanke, M. Follo, D. Quandt, S. F. McDaniel, R. Reski, B. C. Tan & S. A. Rensing. 2014. Molecular evidence for convergent evolution and allopolyploid speciation within the *Physcomitrium-Physcomitrella* species complex. *BMC Evolutionary Biology* 14: 158 [1–19].
- Bell, N. E. & J. Hyvönen. 2010. A phylogenetic circumscription of *Polytrichastrum* (Polytrichaceae): reassessment of sporophyte morphology supports molecular phylogeny. *American Journal of Botany* 97: 566–578.
- Bonfim Santos, M. & M. Stech. 2017. Testing hypotheses on suprageneric relationships and morphological evolution in the Leucobryaceae (Bryophyta). *Plant Systematics and Evolution* 303: 1383–1397.
- Brinda, J. C. & J. J. Atwood (eds.). 2024. *The Bryophyte Nomenclator*. <https://www.bryonames.org/>.
- Brinda, J. C. & V. E. Fedosov. 2023. Nomenclatural corrections in the Dicranidae (Bryophyta). *Arctoa* 32: 261–263.
- Brinda, J. C., D. R. Toren & J. R. Shevock. 2016. *Campylostelium* (Ptychomitriaceae) in the southwestern United States: *Campylostelium laegerae* sp. nov. and *C. pitardii* new to the Americas. *Madroño* 63: 353–358.
- Brinda, J. C. & R. H. Zander. 2020. A new combination in the genus *Molendoa* (Pottiaceae, Bryophyta). *Phytotaxa* 454: 300.
- Brooks, J. R. & J. A. Jáuregui-Lazo. 2023. *Syntrichia lithophila*, a fantastic new amphitropical disjunct from Oregon, United States. *The Bryologist* 126: 221–225.
- Bruch, P. & W. P. Schimper. 1844. *Oedipodium, Tayloria, Dissodon, Tetraplodon, Splachnum, Orthodontium, Mieliichhoferia*. In: P. Bruch, W. P. Schimper & Th. Gumbel, *Bryologia Europaea, seu, genera muscorum Europaeorum monographice illustrata*, Fasc. 23–24. E. Schweizerbart, Stuttgart.
- Buck, W. R. 1984. *Pylaisiadelpha* replaces *Brotherella* (Sematophyllaceae). *Yushania* 1(2): 11–13.
- Buck, W. R. 1994. Three additions to the United States moss flora. *The Bryologist* 97: 89–90.
- Buck, W. R. 1998. Pleurocarpous mosses of the West Indies. *Memoirs of the New York Botanical Garden* 82: 1–400.
- Buck, W. R. & B. H. Allen. 2004. A new name for the eastern North American species of *Ctenidium*. *Evansia* 21: 38–39.
- Buck, W. R. & B. H. Allen. 2017. *Serpoleskea minutissima* comb. nov., a micro-rheophyte in the Pylaisiadelphaceae. *Bryophyte Diversity and Evolution* 39: 69–74.
- Buck, W. R. & D. H. Norris. 1996. *Hedwigia stellata* and *H. detonsa* (Hedwigiaceae) in North America. *Nova Hedwigia* 62: 361–370.
- Câmara, P. E. A. S., M. Carvalho-Silva, D. K. Henriques, J. Guerra, M. T. Gallego, D. R. Poveda & M. Stech. 2018. Pylaisiaceae Schimp. (Bryophyta) revisited. *Journal of Bryology* 40: 251–264.
- Cano, M. J. 2008. Taxonomic revision of *Hennediella* Paris (Pottiaceae, Bryophyta). *Bryophytorum Bibliotheca* 64: 1–142.
- Cano, M. J., J. F. Jiménez, M. Alonso-García & J. A. Jiménez. 2016 [2015]. Untangling *Pseudocrossidium crinitum* s.l. (Pottiaceae, Bryophyta) through molecular and morphometric analysis. *Nova Hedwigia* 102: 89–106.
- Cano, M. J., J. A. Jiménez, M. Martínez, & J. Guerra. 2022. *Hymenostylium chapadense* M.J.Cano & J.A.Jiménez (Pottiaceae), a new species from Brazil and its phylogenetic position based on molecular data. *Journal of Bryology* 44: 51–61.
- Caparrós, R., F. Lara, I. Draper, V. Mazimpaka & R. Garilleti. 2016. Integrative taxonomy sheds light on an old problem: the *Ulota crispa* complex (Orthotrichaceae, Musci). *Botanical Journal of the Linnean Society* 180: 427–451.
- Carter, B. E. 2012. *Scleropodium occidentale* (Brachytheciaceae), a new moss species from western North America. *The Bryologist* 115: 222–230.

- Carter, B. E. 2014. A global revision of *Scleropodium* (Brachytheciaceae, Bryophyta). *Systematic Botany* 39: 681–697.
- Carvalho-Silva, M., M. Stech, L. H. Soares-Silva, W. R. Buck, N. J. Wickett, Y. Liu & P. E. A. S. Câmara. 2017. A molecular phylogeny of the Sematophyllaceae s.l. (Hypnales) based on plastid, mitochondrial and nuclear markers, and its taxonomic implications. *Taxon* 66: 811–831.
- Conard, H. S. 1945. *Gyroweisia reflexa* in North America. *The Bryologist* 48: 28.
- Cox, C. J., B. Goffinet, A. E. Newton, A. J. Shaw & T. A. Hedderson. 2000. Phylogenetic relationships among the diplolepidous-alternate mosses (Bryidae) inferred from nuclear and chloroplast DNA sequences. *The Bryologist* 103: 224–241.
- Cox, C. J. & T. A. Hedderson. 1999. Phylogenetic relationships among the ciliate arthrodontous mosses: evidence from chloroplast and nuclear DNA sequences. *Plant Systematics and Evolution* 215: 119–139.
- Crosby, M. R. & W. R. Buck. 2011. *Nogopterium*, a new name for the genus *Pterogonium* (Musci, Leucodontaceae). *Novon* 21: 424–425.
- Crosby, M. R., R. E. Magill, B. Allen & S. He. 1999. A Checklist of the Mosses (Revised). Missouri Botanical Garden, St. Louis.
- Crum, H. 2001. A *Brachythecium* of Michigan worthy of note. *Evansia* 18: 62.
- Crum, H., W. C. Steere & L. E. Anderson. 1965. A list of the mosses of North America. *The Bryologist* 68: 377–432.
- Crum, H., W. C. Steere & L. E. Anderson. 1973. A new list of mosses of North America north of Mexico. *The Bryologist* 76: 85–130.
- Delgadillo, M. C. 1996. Notes on the distribution of *Aloina* and *Crossidium* (Musci). *The Bryologist* 99: 416–417.
- De Luna, E. 2022. Taxonomy of *Hedwigia filiformis* (Bryopsida: Hedwigiaceae), with the clarification of the confounded identity of its synonym *Hedwigia integrifolia*. *Acta Botanica Mexicana* 129: e2080 [1–16].
- Duffy, A., B. Aguero, H. Stenøien, K. I. Flatberg, M. S. Ignatov, K. Hassel & A. J. Shaw. 2020. Phylogenetic structure in the *Sphagnum recurvum* complex (Bryophyta: Sphagnaceae) relative to taxonomy and geography. *American Journal of Botany* 107: 1283–1295.
- Eckel, P. M. 2007. *Pleurochaete*. In: *Flora of North America Editorial Committee* (eds.), *Flora of North America* 27: 496–498. Oxford University Press, New York.
- Ellis, L. T. (ed.), O. M. Afonina, R. L. Andriamiarisoa, G. Asthana, R. Bharti, P. Aymerich, B. Bambe, M. F. Boiko, M. Brugués, E. Ruiz, Ll. Sáez Goñalons, M. J. Cano, R. M. Ros, L. Číhal, J. Deme, J. Csiky, G. Dihoru, P. Dřevojan, T. Ezer, V. E. Fedosov, E. A. Ignatova, A. P. Seregin, C. A. Garcia, A. Martins, C. Sérgio, M. Sim-Sim, A. S. B. Rodrigues, S. R. Gradstein, C. Reeb, A. Irmah, M. Suleiman, T. Koponen, J. Kučera, M. Lebouvier, Y. LiQun, D. G. Long, A. I. Maksimov, T. A. Maksimova, J. Muñoz, M. Nobis, A. Nowak, R. Ochrya, S. V. O’Leary, F. Osorio, O. Yu. Pisarenko, V. Plášek, Z. Skoupá, A. Schäfer-Verwimp, N. Schnyder, J. R. Shevock, S. Ștefănuț, S. Mamtimin, B.-Y. Sun, S. J. Park, D. Ya. Tubanova, J. Váňa, G. J. Wolski, K.-Y. Yao, Y.-J. Yoon & E. Yücel. 2018. New national and regional bryophyte records, 56. *Journal of Bryology* 40: 271–296.
- Enroth, J., S. Olsson, S. Huttunen, V. Buchbender, R. Tangney, M. Stech, L. Hedenäs & D. Quandt. 2019. Orthostichellaceae fam. nov. and other novelties in pleurocarpous mosses revealed by phylogenetic analyses. *The Bryologist* 122: 219–245.
- Enroth, J., S. Olsson, S. Huttunen & D. Quandt. 2022. *Neckera*, *Forstroemia* and *Alleniella* (Neckeraceae, Bryophyta) redefined based on phylogenetic analyses. *The Bryologist* 125: 311–327.
- Fedosov, V. E. 2012. *Encalypta* sect. *Rhabdotheca* in Russia. *Arctoa* 21: 101–112.
- Fedosov, V. E., I. V. Czernyadjeva, O. M. Afonina & E. A. Ignatova. 2023a. On the distribution of *Ditrichum divaricatum* (Ditrichaceae, Bryophyta) in North Pacific region. *Botanica Pacifica* 12: 163–167.
- Fedosov, V. E., A. V. Fedorova, A. E. Fedosov & M. S. Ignatov. 2016b. Phylogenetic inference and peristome evolution in haplolepidous mosses, focusing on Pseudoditrichaceae and Ditrichaceae s.l. *Botanical Journal of the Linnean Society* 181: 139–155.
- Fedosov, V. E., A. V. Fedorova, E. A. Ignatova, J. C. Brinda & J. Kučera. 2024. Overlooked and misunderstood: a morpho-molecular revision of Ditrichaceae s.str. (Dicranidae, Bryophyta), with a focus on the Holarctic species of *Ditrichum*. Taxon in review.
- Fedosov, V. E., A. V. Fedorova, E. A. Ignatova & M. S. Ignatov. 2017. A revision of the genus *Seligeria* (Seligeriaceae, Bryophyta) in Russia inferred from molecular data. *Phytotaxa* 323: 27–50.
- Fedosov, V. E., A. V. Fedorova, E. A. Ignatova & J. Kučera. 2023b. New taxonomic arrangement of *Dicranella* s.l. and *Aongstroemia* s.l. (Dicranidae, Bryophyta). *Plants* 12(6): 1360 [1–38].
- Fedosov, V. E., A. V. Fedorova, J. Larraín, M. Bonfim Santos, M. Stech, J. Kučera, J. C. Brinda, D. Ya. Tubanova, M. von Konrat, E. A. Ignatova & M. S. Ignatov. 2021 [2020]. Unity in diversity: phylogenetics and taxonomy of Rhabdoweisiaceae (Dicranales, Bryophyta). *Botanical Journal of the Linnean Society* 195: 545–567.
- Fedosov, V. E., A. V. Fedorova, A. V. Troitsky, V. K. Bobrova & M. S. Ignatov. 2016a. On the systematic position of *Hymenoloma* (Bryophyta). *Arctoa* 25: 119–130.
- Fedosov, V. E., A. V. Shkurko, A. V. Fedorova, E. A. Ignatova, E. N. Solovyeva, J. C. Brinda, M. S. Ignatov & J. Kučera. 2022. Need for split: integrative taxonomy reveals unnoticed diversity in the subaquatic species of *Pseudohygrohypnum* (Pylaisiaceae, Bryophyta). *PeerJ* 10: e13260 [1–69].
- Flagmeier, M., I. Draper, B. Vigalondo, R. Garilleti & F. Lara. 2021. So close yet so far: the disjunct global range of *Orthotrichum consobrinum* (Orthotrichaceae) includes North America. *The Bryologist* 124: 403–413.
- Flora of North America Editorial Committee (eds.). 2007. Bryophyta, part 1. *Flora of North America North of Mexico* 27: i–xxi, 1–713. Oxford University Press, New York.
- Flora of North America Editorial Committee (eds.). 2014. Bryophyta, part 2. *Flora of North America North of Mexico* 28: i–xxi, 1–702. Oxford University Press, New York.
- Frahm, J.-P. 2007. *Campylopus*. In: *Flora of North America Editorial Committee* (eds.), *Flora of North America* 27: 366–376. Oxford University Press, New York.
- Fransén, S. 1995[1996]. A taxonomic revision of neotropical *Bartramia* section *Vaginella*. *Lindbergia* 20: 147–179.
- Gallego, M. T., M. J. Cano & J. Guerra. 2004. A taxonomic study of *Syntrichia laevipila* (Pottiaceae, Musci) complex. *Botanical Journal of the Linnean Society* 145: 219–230.
- Gallego, M. T., M. J. Cano, J. A. Jiménez & J. Guerra. 2022. Circumscription and phylogenetic position of two propagulose species of *Syntrichia* (Pottiaceae, Bryophyta) reveals minor realignments within the tribe Syntricheae. *Plants* 11(5): 626 [1–22].

- Gardiner, A. A., M. S. Ignatov, S. Huttunen & A. V. Troitsky. 2005. On resurrection of the families Pseudoleskeaceae Schimp. and Pylaisiaceae Schimp. (Musci, Hypnales). *Taxon* 54: 651–663.
- Garilleti, R., F. Lara & V. Mazimpaka. 2001. *Orthotrichum underwoodii*, sp. nov. (Orthotrichaceae, Bryopsida) from California. *The Bryologist* 104: 268–273.
- Garilleti, R., F. Lara & V. Mazimpaka. 2006a. *Orthotrichum anodon* (Orthotrichaceae), a new species from California, and its relationships to other *Orthotricha* with puckered mouths. *The Bryologist* 109: 188–196.
- Garilleti, R., F. Lara & V. Mazimpaka. 2006b. On the presence of *Orthotrichum shawii* (Orthotrichaceae) in California. *The Bryologist* 109: 510–515.
- Garilleti, R., J. R. Shevock, D. H. Norris & F. Lara. 2011. *Orthotrichum mazimpakanum* sp. nov. and *O. anodon* (Orthotrichaceae), two similar species from California. *The Bryologist* 114: 346–355.
- GBIF.org. 2024a. GBIF occurrence download for *Acaulon mediterraneum* on 22 October 2024. <https://doi.org/10.15468/dl.z8gftm>.
- GBIF.org. 2024b. GBIF occurrence download for *Hedwigia emodica* on 23 July 2024. <https://doi.org/10.15468/dl.jje9pa>.
- GBIF.org. 2024c. GBIF occurrence download for *Hedwigia nivalis* on 23 July 2024. <https://doi.org/10.15468/dl.bpsz2s>.
- GBIF.org. 2024d. GBIF occurrence download for *Tortula israelis* on 22 October 2024. <https://doi.org/10.15468/dl.dw7tf6>.
- Goffinet, B., W. R. Buck & A. J. Shaw. 2009a. Addenda to the classification of mosses. I. Andreaeophytina stat. nov. and Andreaebryophytina stat. nov. *The Bryologist* 112: 856–857.
- Goffinet, B., W. R. Buck & A. J. Shaw. 2009b[2008]. Morphology, anatomy, and classification of the Bryophyta. Pp. 55–138. In B. Goffinet & A. J. Shaw (eds.), *Bryophyte Biology* (ed. 2). Cambridge University Press, Cambridge, U.K.
- Goffinet, B. & A. J. Shaw. 2002. Independent origins of cleistocarpy in the Splachnaceae: analyses of cpDNA sequences and polyploidy of the Voitiioideae (Bryophyta). *Systematic Botany* 27: 203–208.
- Goffinet B., A. J. Shaw, C. J. Cox, N. J. Wickett & S. Boles. 2004. Phylogenetic inferences in the Orthotrichoideae (Orthotrichaceae: Bryophyta) based on variation in four loci from all genomes. *Monographs in Systematic Botany from the Missouri Botanical Garden* 98: 270–289.
- Goffinet, B. & D. H. Vitt. 1998. Revised generic classification of the Orthotrichaceae based on a molecular phylogeny and comparative morphology. Pages 143–159. In: J. W. Bates, N. W. Ashton & J. G. Duckett (eds.), *Bryology for the Twenty-first Century*. Maney Publishing and the British Bryological Society, Leeds, U.K.
- Granzow-de la Cerda, Í. 1997. Revision and phylogeny of *Anomodon* and *Herpetineuron* (Anomodontaceae, Musci). *Contributions from the University of Michigan Herbarium* 21: 205–275.
- Granzow-de la Cerda, Í. & R. P. G. Düll. 2009. *Anomodon longifolius* and *A. thraustus* (Anomodontaceae) new to North America, north of Mexico. *The Bryologist* 112: 606–610.
- Greven, H. C. 2010. *Grimmia texicana* sp. nov. (Grimmiaceae) from Texas and its separation from *Grimmia arizonae*. *The Bryologist* 113: 360–364.
- Griffin, III, D. G. 2014. Bartramiaceae. In: *Flora of North America* Editorial Committee (eds.), *Flora of North America* 28: 97–112. Oxford University Press, New York.
- Griffin, III, D. G. & W. R. Buck. 1989. Taxonomic and phylogenetic studies on the Bartramiaceae. *The Bryologist* 92: 368–380.
- Grout, A. J. 1940. List of mosses of North America north of Mexico. *The Bryologist* 43: 117–131.
- Guerra, J. 2021. Notes on the taxonomy, chorology and habitat of *Pohlia greenii* Brid. (Bryophyta) in the Iberian Peninsula. *Herzogia* 34: 197–202.
- Guerra, J., R. M. Ros & M. J. Cano. 1994. *Pterygoneurum subsessile* (Brid.) Jur. var. *kieneri* Hab. (Musci, Pottiaceae), a putative hybrid. *Nova Hedwigia* 59: 507–510.
- Habeeb, H. 1950. Nomenclatural and other notes on mosses. *Rhodorha* 52: 72–75.
- Hanusch, M., E. M. Ortiz, J. Patiño & H. Schaefer. 2020. Biogeography and integrative taxonomy of *Epipterygium* (Mniaceae, Bryophyta). *Taxon* 69: 1150–1171.
- Haring, I. M. 1961. A checklist of the mosses of the state of Arizona. *The Bryologist* 64: 222–240.
- Harpel, J. A. 2006. A new combination in *Tetraphis* (Bryophyta: Tetraphidaceae). *Sida* 22: 549–554.
- Hastings, R. I. 1999. Taxonomy and biogeography of the genus *Coscinodon* (Bryopsida, Grimmiaceae) in North America, including a new species. *The Bryologist* 102: 265–286.
- Hastings, R. I. 2008. *Grimmia torenii* sp. nov. (Grimmiaceae) from California and its separation from *G. ovalis* and *G. tergestina*. *The Bryologist* 111: 463–475.
- Hastings, R. I. & H. C. Greven. 2007. *Grimmia*. In: *In: Flora of North America* Editorial Committee (eds.), *Flora of North America* 27: 225–258. Oxford University Press, New York.
- He, S. 2005. A revision of the genus *Leptopterigynandrum* (Bryopsida, Leskeaceae). *Journal of the Hattori Botanical Laboratory* 97: 1–38.
- Hedenäs, L. 1990[1992]. Taxonomic and nomenclatural notes on the genera *Calliergonella* and *Breidleria*. *Lindbergia* 16: 161–168.
- Hedenäs, L. 1993. A generic revision of the *Warnstorfia-Calliergon* group. *Journal of Bryology* 17: 447–479.
- Hedenäs, L. 1997. A partial generic revision of *Campylium* (Musci). *The Bryologist* 100: 65–88.
- Hedenäs, L. 1998. An overview of the *Drepanocladus sendtneri* complex. *Journal of Bryology* 20: 83–102.
- Hedenäs, L. 2003. Amblystegiaceae (Musci). *Flora Neotropica Monograph* 89: 1–108.
- Hedenäs, L. 2006. Additional insights into the phylogeny of *Calliergon*, *Loeskypnum*, *Straminergon*, and *Warnstorfia* (Bryophyta: Calliergonaceae). *Journal of the Hattori Botanical Laboratory* 100: 125–134.
- Hedenäs, L. 2008a. Molecular variation and speciation in *Antitrichia curtispindula* s.l. (Leucodontaceae, Bryophyta). *Botanical Journal of the Linnean Society* 156: 341–354.
- Hedenäs, L. 2008b. Molecular variation in *Drepanocladus aduncus* s. l. does not support recognition of more than one species in Europe. *Journal of Bryology* 30: 108–120.
- Hedenäs, L. 2011. Relationships in the *Timmia norvegica* complex (Timmiaceae) based on nuclear and chloroplast sequence data. *The Bryologist* 114: 749–755.
- Hedenäs, L. 2012. Global phylogeography in *Sanionia uncinata* (Amblystegiaceae: Bryophyta). *Botanical Journal of the Linnean Society* 168: 19–42.
- Hedenäs, L. 2014a. Calliergonaceae. In: *Flora of North America* Editorial Committee (eds.), *Flora of North America* 28: 384–403. Oxford University Press, New York.
- Hedenäs, L. 2014b. *Drepanocladus*. In: *Flora of North America* Editorial Committee (eds.), *Flora of North America* 28: 292–297. Oxford University Press, New York.

- Hedenäs, L. 2014c. *Palustriella*. In: Flora of North America Editorial Committee (eds.), Flora of North America 28: 267–269. Oxford University Press, New York.
- Hedenäs, L. 2017. *Brachythecium tauriscorum* Molendo, a widely distributed Arctic-alpine species. *Journal of Bryology* 39: 201–203.
- Hedenäs, L. 2020. Cryptic and morphologically recognizable species diversity within Scandinavian *Plagiopus oederianus* (Bryophyta: Bartramiaceae). *Lindbergia* 43: 01130 [1–12].
- Hedenäs, L. 2021. Incipient speciation in Scandinavian *Distichium capillaceum* (Distichiaceae, Bryophyta). *Lindbergia* 42: 01144 [1–12].
- Hedenäs, L., O. I. Kuznetsova & M. S. Ignatov. 2020. A revision of the genus *Tomentypnum* (Amblystegiaceae) in northern Eurasia. *The Bryologist* 123: 377–395.
- Hedenäs, L., G. Oliván & P. Eldenäs. 2005. Phylogeny of the Calliergonaceae (Bryophyta) based on molecular and morphological data. *Plant Systematics and Evolution* 252: 49–61.
- Hedenäs, L. & L. Söderström. 1991[1992]. *Leskea rupestris*, an older name for *Pseudoleskeella sibirica*. *Lindbergia* 17: 64–66.
- Herid, W., S. He, & D.-P. Zhao. 2019. *Weissia armata*, a rare Mediterranean coastline and central Asian species new to North America. *Herzogia* 32: 252–255.
- Hernández-Maqueda, R., D. Quandt, O. Werner & J. Muñoz. 2008. Phylogeny and classification of the Grimmiaceae/Ptychomitriaceae complex (Bryophyta) inferred from cpDNA. *Molecular Phylogenetics and Evolution* 46: 863–877.
- Hodgetts, N. G., L. Söderström, T. L. Blockeel, S. Caspari, M. S. Ignatov, N. A. Konstantinova, N. Lockhart, B. Papp, C. Schröck, M. Sim-Sim & D. Bell. 2020. An annotated checklist of bryophytes of Europe, Macaronesia and Cyprus. *Journal of Bryology* 42: 1–116.
- Holyoak, D. T. 2004. Taxonomic notes on some European species of *Bryum* (Bryopsida: Bryaceae). *Journal of Bryology* 26: 247–264.
- Holyoak, D. T. & N. Pedersen. 2007. Conflicting molecular and morphological evidence of evolution within the Bryaceae (Bryopsida) and its implications for generic taxonomy. *Journal of Bryology* 29: 111–124.
- Hornschuch, C. F. 1818. De *Voitia* et *Systylis* Novis Muscorum Frondosorum Generibus Commentatio. 22 pp. Marcobritae, typis Knenleinianis, Erlangae.
- Hornschuch, C. F. 1824. *Ptychostomum* nomum muscorum frondosorum genus. Constituit et descripsit. *Sylloge Plantarum Novarum itemque Minus Cognitarum* 1(3): 62–64.
- Horton, D. G. 1983. A revision of the Encalyptaceae (Musci), with particular reference to the North American taxa. Part II. *Journal of the Hattori Botanical Laboratory* 54: 353–532.
- Huang, W.-Z., C. Shen, H. Xu, L. Shu, S. Mamtimin, Y.-H. Wu & R.-L. Zhu. 2024. A synopsis of *Dicranum* Hedw. (Dicranaceae, Bryophyta) in China, with special references to four species newly reported and re-evaluation of *Dicranum psathyrum* Klazenga. *Plants* 13(13): 1759 [1–26].
- Huttunen, S., A. A. Gardiner & M. S. Ignatov. 2007. Advances in knowledge of the Brachytheciaceae (Bryophyta). In A. E. Newton & R. S. Tangney (eds.), *Pleurocarpous mosses: systematics and evolution*. Systematics Association Special Volume 71: 117–143.
- Huttunen, S. & M. S. Ignatov. 2010. Evolution and taxonomy of aquatic species in the genus *Rhynchostegium* (Brachytheciaceae, Bryophyta). *Taxon* 59: 791–808.
- Huttunen, S., M. S. Ignatov, D. Quandt & L. Hedenäs. 2013. Phylogenetic position and delimitation of the moss family Plagiotheciaceae in the order Hypnales. *Botanical Journal of the Linnean Society* 171: 330–353.
- Huttunen, S., O. Kuznetsova, M. Li, Y.-F. Wang & M. S. Ignatov. 2015. East Asian origin and Northern Hemisphere diversification in the *Brachythecium novae-angliae*-complex (Bryophyta). *Taxon* 64: 911–930.
- Ignatov, M. S. 2014. *Brachythecium*. In: Flora of North America Editorial Committee (eds.), Flora of North America 28: 414–428. Oxford University Press, New York.
- Ignatov, M. S., O. M. Afonina & E. A. Ignatova. 2006[2007]. Check-list of mosses of East Europe and North Asia. *Arctoa* 15: 1–130.
- Ignatov, M. S., O. M. Afonina, O. I. Kuznetsova & E. A. Ignatova. 2012. The genus *Leptopterigynandrum* (Taxiphyllaceae, Bryophyta) in Russia. *Arctoa* 21: 207–220.
- Ignatov, M. S., A. V. Federova & V. E. Fedosov. 2019a. On the taxonomy of Anomodontaceae and *Heterocladium* (Bryophyta). *Arctoa* 28: 75–102.
- Ignatov, M. S. & S. Huttunen. 2002[2003]. Brachytheciaceae (Bryophyta) — a family of sibling genera. *Arctoa* 11: 245–296.
- Ignatov, M. S. & E. A. Ignatova. 2004. Moss flora of the Middle European Russia. Volume 2: Fontinalaceae–Amblystegiaceae. *Arctoa* 11(Supplement 2): 609–960.
- Ignatov, M. S. & E. A. Ignatova. 2017. Tetrarhizales. In: M. S. Ignatov (ed.), Moss flora of Russia. Volume 2: Oedipodiales–Grimmiales. *Arctoa* 26(Supplement 1): 115–123.
- Ignatov, M. S., E. A. Ignatova, O. I. Kuznetsova & V. E. Fedosov. 2019b. On moss genera *Hylocomiadelphus* Ochyra & Stebel and *Rhytidiadelphus* (Limpr.) Warnst. *Acta Musei Silesiae, Scientiae Naturales* 68: 123–134.
- Ignatov, M. S., J. Kučera, L. Hedenäs, O. I. Kuznetsova & E. A. Ignatova. 2020. A revision of the genus *Orthothecium* (Plagiotheciaceae, Bryophyta) in northern Eurasia. *Arctoa* 29: 10–48.
- Ignatova, E. A. & H. H. Blom. 2017. *Schistidium*. In: M. S. Ignatov (ed.), Moss flora of Russia. Volume 2: Oedipodiales–Grimmiales. *Arctoa* 26(Supplement 1): 438–551.
- Ignatova, E. A., H. H. Blom, D. V. Goryunov & I. A. Millyutina. 2009. On the genus *Schistidium* (Grimmiaceae, Musci) in Russia. *Arctoa* 19: 195–233.
- Ignatova, E. A., I. V. Czernyadjeva, A. V. Fedorova & M. S. Ignatov. 2021. A morphological and molecular phylogenetic study of the genus *Calliergon* (Calliergonaceae, Bryophyta) in Russia. *Arctoa* 30: 8–24.
- Ignatova, E. A. & V. E. Fedosov. 2008. Species of *Dicranum* (Dicranaceae, Bryophyta) with fragile leaves in Russia. *Arctoa* 17: 63–83.
- Ignatova, E. A., V. E. Fedosov, A. V. Fedorova, U. N. Spirina & M. S. Ignatov. 2019. On moss family Lembophyllaceae in the Russian Far East. *Arctoa* 28: 179–202.
- Ignatova, E. A., A. V. Fedorova & M. S. Ignatov. 2020a. On the genera *Isopterygiopsis* and *Isopterygiella*, gen. nov. (Plagiotheciaceae) in Russia. *Arctoa* 29: 49–62.
- Ignatova, E. A., O. I. Kuznetsova, V. E. Fedosov & M. S. Ignatov. 2016. On the genus *Hedwigia* (Hedwigiaceae, Bryophyta) in Russia. *Arctoa* 25: 241–277.
- Ignatova, E. A., O. I. Kuznetsova, N. R. Shafigullina, V. E. Fedosov & M. S. Ignatov. 2020b. The genus *Pylaisia* (Pylaisiaceae, Bryophyta) in Russia. *Arctoa* 29: 135–178.
- Imwattana, K., B. Aguero, M. Nieto-Lugilde, A. Duffy, J. Jaramillo-Chico, K. Hassel, O. Afonina, P. Lamkowski & A. J. Shaw. 2024.

- Parallel patterns of genetic diversity and structure in circumboreal species of the *Sphagnum capillifolium* complex. *American Journal of Botany* 111(5): e16348 [1–16].
- Inoue, Y. & H. Tsubota. 2014. On the systematic position of the genus *Timmiella* (Dicranidae, Bryopsida) and its allied genera, with the description of a new family Timmiellaceae. *Phytotaxa* 181: 151–162.
- Ireland, Jr., R. R. 2004. *Dacryophyllum falcifolium*, a new North American genus and species (Musci: Hypnaceae) from California. *Novon* 14: 70–74.
- Ivanova, E. I. & M. S. Ignatov. 2017. Polytrichopsida. In: M. S. Ignatov (ed.), *Moss flora of Russia. Volume 2: Oedipodiales–Grimmiales*. *Arctoa* 26(Supplement 1): 40–114.
- Iwatsuki, Z., Y. Tateishi & T. Suzuki. 1999. *Campylostelium* in Japan. *Hikobia* 13: 79–85.
- Jáuregui-Lazo, J. A., J. C. Brinda, GoFlag Consortium & B. D. Mishler. 2023[2022]. The phylogeny of *Syntrichia*: an ecologically diverse clade of mosses with an origin in South America. *American Journal of Botany* 110(1): e16103 [1–19].
- Jiménez, J. A., M. J. Cano & J. Guerra. 2022. A multilocus phylogeny of the moss genus *Didymodon* and allied genera (Pottiaceae): generic delimitations and their implications for systematics. *Journal of Systematics and Evolution* 60: 281–304.
- Jiménez, J. A., R. M. Ros, M. J. Cano & J. Guerra. 2005. A new evaluation of the genus *Trichostomopsis* (Pottiaceae, Bryophyta). *Botanical Journal of the Linnean Society* 147: 117–127.
- Jiménez, S. & D. R. Toren. 2018. *Philonotis minuta* (Bartramiaceae, Bryophyta) is proposed as the correct name for *P. brevifolia*, and recorded for the first time in North America from California (U.S.A.). *Cryptogamie, Bryologie* 39: 389–396.
- Jiménez, J. A., D. R. Toren & J. R. Shevock. 2014. *Didymodon californicus* (Pottiaceae), a new species from California, U.S.A. *Phytotaxa* 158: 105–110.
- Joya, S. & T. T. McIntosh. 2012. The moss, *Plagiomnium undulatum* (Mitt.) T.Kop., new to North America, reported from Vancouver and Victoria, British Columbia. *Botanical Electronic News* 459.
- Kariyawasam, I. U. 2021. Phylogeny, Species Delimitation and Taxonomy in *Polytrichum* sect. *Polytrichum* (Polytrichaceae Schwägr.; Bryophyta). Doctoral dissertation, University of Edinburgh, Edinburgh, Scotland.
- Karlin, E. F., M. M. Giusti, R. A. Lake, S. B. Boles & A. J. Shaw. 2010. Microsatellite analysis of *Sphagnum centrale*, *S. henryense*, and *S. palustre* (Sphagnaceae). *The Bryologist* 113: 90–98.
- Kellman, K. M. 2008. *Syntrichia sucrosa* sp. nov. (Pottiaceae), a new species from California. *The Bryologist* 111: 676–684.
- Kellman, K. M. 2012. *Tortula brevissima* Schiffn., new to North America. *Evansia* 29: 50–53.
- Kellman, K. M. 2017. Noteworthy collections: California, *Gemmarium eremaeum*. *Madroño* 64: 60.
- Kiebacher, T. & M. Meier. 2021. The generic placement of *Grimmia teretinervis* Limpr. *Herzogia* 33: 544–547.
- Köckinger, H. & L. Hedenäs. 2021[2020]. Solving nomenclatural problems surrounding *Mollia thrausta* Stirt., *Barbula tortuosa* var. *fragilifolia* Jur. and *Tortella fasciculata* (Culm.) Culm. *Journal of Bryology* 43: 186–189.
- Köckinger, H., M. Lüth, O. Werner & R. M. Ros. 2018. *Tortella mediterranea* (Pottiaceae), a new species from southern Europe, its molecular affinities, and taxonomic notes on *T. nitida*. *The Bryologist* 121: 560–570.
- Koponen, A. 1982. The generic classification of the Splachnaceae. *Beihefte zur Nova Hedwigia* 71: 239–247.
- Koponen, A. 1990. Entomophily in the Splachnaceae. *Botanical Journal of the Linnean Society* 104: 115–127.
- Koponen, T. 1968. Generic revision of Mniaceae Mitt. (Bryophyta). *Annales Botanici Fennici* 5: 117–151.
- Koponen, T., E. A. Ignatova, O. I. Kuznetsova & M. S. Ignatov. 2012. *Philonotis* (Bartramiaceae, Bryophyta) in Russia. *Arctoa* 21: 21–62.
- Koponen, T. & Y. Sun. 2017[2016]. Preliminary study on phylogenetic position and delimitation of the ciliate arthroodontous genera of the moss family Mniaceae. *Journal of Bryology* 39: 23–38.
- Kosovich-Anderson, Y. I. & M. S. Ignatov. 2010[2011]. Three interesting Brachytheciaceae mosses from the Beartooth Plateau (Rocky Mountains, Wyoming, U.S.A.). *Arctoa* 19: 183–190.
- Kramer, W. A. 2023. The genus *Syntrichia* Brid. (Pottiaceae, Bryophyta) in the Holarctic with special consideration of North America. *Nova Hedwigia Beiheft* 154: 1–234.
- Kučera, J., J. Košnar & O. Werner. 2013. Partial generic revision of *Barbula* (Musci: Pottiaceae): re-establishment of *Hydrogonium* and *Streblotrichum*, and the new genus *Gymnobarbula*. *Taxon* 62: 21–39.
- Kučera, J., O. I. Kuznetsova, A. Manukjanová & M. S. Ignatov. 2019. A phylogenetic revision of the genus *Hypnum*: towards completion. *Taxon* 68: 628–660.
- Kuznetsova, O. I., A. V. Fedorova, E. D. Lapshina, M. A. Kolesnikova, V. E. Fedosov, M. S. Ignatov & E. A. Ignatova (eds.). 2023. Bryophyte molecular barcoding records. 8. *Arctoa* 32: 75–77.
- Lara, F., I. Draper, M. Flagmeier, J. A. Calleja, V. Mazimpaka & R. Garilleti. 2020. Let's make *Pulviger* great again: re-circumscription of a misunderstood group of Orthotrichaceae that diversified in North America. *Botanical Journal of the Linnean Society* 193: 180–206.
- Lara, F., R. Garilleti, B. Goffinet, I. Draper, R. Medina, B. Vigalondo & V. Mazimpaka. 2016. *Lewinskya*, a new genus to accommodate the phaneroporou and monoicous taxa of *Orthotrichum* (Bryophyta, Orthotrichaceae). *Cryptogamie, Bryologie* 37: 361–382.
- Larraín, J., D. Quandt, M. Stech & J. Muñoz. 2013. Lumping or splitting? The case of *Racomitrium* (Bryophytina: Grimmiaceae). *Taxon* 62: 1117–1132.
- Lewinsky-Haapasaari, J. 1998. *Orthotrichum shevockii* (Orthotrichaceae), a new moss species from the southern Sierra, California. *The Bryologist* 101: 435–438.
- Lewinsky-Haapasaari, J. & D. H. Norris. 1998. A re-evaluation of *Orthotrichum euryphyllum*. *The Bryologist* 101: 295–302.
- Lewis, L. R., E. M. Biersma, S. B. Carey, K. Holsinger, S. F. McDaniel, R. Rozzi & B. Goffinet. 2017. Resolving the Northern Hemisphere source region for the long-distance dispersal event that gave rise to the South American endemic dung moss *Tetraplodon fuegianus*. *American Journal of Botany* 104: 1651–1659.
- Lewis, L. R., R. Rozzi & B. Goffinet. 2014. Direct long-distance dispersal shapes a New World amphitropical disjunction in the dispersal-limited dung moss *Tetraplodon* (Bryopsida: Splachnaceae). *Journal of Biogeography* 41: 2385–2395.
- Li, M., J. Ma, M. S. Ignatov, B. C. Tan, S. Huttunen, J.-C. Zhao & Y.-F. Wang. 2015. Taxonomic re-assessment of *Kindbergia* (Brachytheciaceae, Bryophyta) in China, with a description of *Pseudokindbergia* gen. nov. *Cryptogamie, Bryologie* 36: 47–68.
- Li, X.-Q., Q. Zuo, M. Li, S. He, Y. Jia & Y.-F. Wang. 2015. Phylogenetic relationships and species delimitation among the taxa with propagula in *Pseudotaxiphyllum* (Plagiotheciaceae, Bryophyta). *Journal of Systematics and Evolution* 53: 179–190.

- Li, Y.-F., L. Luo, Y. Liu, Q. He, N.-N. Yu, N. Gaowa, Z.-Q. Yi, J.-J. Wang, W. Han, T. Peng, B.-C. Ho, X.-L. He, L. Zhang, Z.-D. Chen, Y. Jia & Q.-H. Wang. 2024. The Bryophyte Phylogeny Group: a revised familial classification system based on plastid phylogenomic data. *Journal of Systematics and Evolution* 62: 577–588.
- Lindberg, S. O. 1862[1863]. Om ett nytt slägte, *Epiterygium*, bland bladmosorna. Öfversigt af Kongl. Vetenskaps-Akademiens Förhandlingar 19: 599–609.
- Liu, Y., J. M. Budke & B. Goffinet. 2012. Phylogenetic inference rejects sporophyte based classification of the Funariaceae (Bryophyta): rapid radiation suggests rampant homoplasy in sporophyte evolution. *Molecular Phylogenetics and Evolution* 62: 130–145.
- Liu, Y., M. G. Johnson, C. J. Cox, R. Medina, N. Devos, A. Vanderpoorten, L. Hedenäs, N. E. Bell, J. R. Shevock, B. Aguero, D. Quandt, N. J. Wickett, A. J. Shaw & B. Goffinet. 2019. Resolution of the ordinal phylogeny of mosses using targeted exons from organellar and nuclear genomes. *Nature Communications* 10: 1485 [1–11].
- Magill, R. E. 2006. A new species of *Encalypta* (Encalyptaceae) from West Texas. *The Bryologist* 109: 398–400.
- Magombo, Z. L. K. 2003. Taxonomic revision of the moss family Diphysciaceae M.Fleisch. *Journal of the Hattori Botanical Laboratory* 94: 1–86.
- Mahévas, T., C. Schneider, T. Schneider, D. Cartier & T. Géhin. 2016. Contribution à la connaissance de la bryoflore du massif vosgien. Les Nouvelles Archives de la Flore Jurassienne et du Nord-Est de la France 14: 2–45.
- Maier, E. 2010. The genus *Grimmia* Hedw. (Grimmiaceae, Bryophyta) — a morphological-anatomical study. *Boissiera* 63: 1–377.
- Manierre, W. R. 1998. *Gyroweisia reflexa*, a moss new to the United States. *Evansia* 15: 30–31.
- Marino, P. C. 1988. Coexistence on divided habitats: mosses in the family Splachnaceae. *Annales Zoologici Fennici* 25: 89–98.
- Marino, P. C. 1991a. Competition between mosses (Splachnaceae) in patchy habitats. *The Journal of Ecology* 79: 1031–1046.
- Marino, P. C. 1991b. Dispersal and coexistence of mosses (Splachnaceae) in patchy habitats. *The Journal of Ecology* 79: 1047–1060.
- Marino, P. C. 2014. Splachnaceae. In: *Flora of North America Editorial Committee* (eds.), *Flora of North America* 28: 14–20. Oxford University Press, New York.
- Marino, P., R. Raguso & B. Goffinet. 2009. The ecology and evolution of fly dispersed dung mosses (Family Splachnaceae): manipulating insect behaviour through odour and visual cues. *Symbiosis* 47: 61–76.
- Mastracci, M. 2003. *Thamnobryum neckeroides* (Bryopsida: Neckeraceae): lectotypification, synonymies, diagnostic characters, habitat and distribution. *Journal of Bryology* 25: 115–120.
- Matsui, T. & Z. Iwatsuki. 1990. A taxonomic revision of the family Ditrichaceae (Musci) of Japan, Korea and Taiwan. *Journal of the Hattori Botanical Laboratory* 68: 317–366.
- McIntosh, T. T. 2007. *Schistidium*. In: *Flora of North America Editorial Committee* (eds.), *Flora of North America* 27: 207–225. Oxford University Press, New York.
- McIntosh, T. T., H. H. Blom, O. I. Kuznetsova & E. A. Ignatova. 2017. *Schistidium relictum* (Grimmiaceae, Bryophyta), a new moss species from northwest North America and Siberia. *Phytotaxa* 299: 234–242.
- McIntosh, T. T., H. H. Blom, D. R. Toren & J. R. Shevock. 2015. Two new species of *Schistidium* (Grimmiaceae, Bryophyta) from western North America. *Phytotaxa* 213: 57–64.
- McIntosh, T. T. & S. G. Newmaster. 2014. *Plagiomnium* T.J.Koponen. In: *Flora of North America Editorial Committee* (eds.), *Flora of North America* 28: 229–235. Oxford University Press, New York.
- McLaughlin, J. T. 2024. The bryophyte flora of Henry W. Coe State Park. *Madroño* 71: 12–42.
- McLaughlin, J. T. & B. E. Carter. 2024. Morphological and ecological evidence for the recognition of *Anacolia baueri* (Bartramiaceae) as distinct from *A. menziesii*. *The Bryologist* 127: 316–324.
- Medina, R., M. G. Johnson, Y. Liu, N. J. Wickett, A. J. Shaw & B. Goffinet. 2019a. Phylogenomic delineation of *Physcomitrium* (Bryophyta: Funariaceae) based on targeted sequencing of nuclear exons and their flanking regions rejects the retention of *Physcomitrella*, *Physcomitridium* and *Aphanorhagma*. *Journal of Systematics and Evolution* 57: 404–417.
- Medina, R., M. G. Johnson, Y. Liu, N. Wilding, T. A. J. Hedderson, N. J. Wickett & B. Goffinet. 2018[2017]. Evolutionary dynamism in bryophytes: phylogenomic inferences confirm rapid radiation in the moss family Funariaceae. *Molecular Phylogenetics and Evolution* 120: 240–247.
- Medina, R., M. G. Johnson, N. R. Patel, G. E. Tocci, D. R. Toren & B. Goffinet. 2022. Vindication of *Physcomitrium pygmaeum* (Funariaceae), an elusive and endangered moss from North America's Great Basin. *The Bryologist* 125: 528–540.
- Medina, R., F. Lara, B. Goffinet, R. Garilleti & V. Mazimpaka. 2013. Unnoticed diversity within the disjunct moss *Orthotrichum tenellum* s.l. validated by morphological and molecular approaches. *Taxon* 62: 1133–1152.
- Medina, R., F. Lara, B. Goffinet, R. Garilleti & V. Mazimpaka. 2019b. On the priority of *Orthotrichum cylindrocarpum* over *O. coulteri* and Lesquereux's early vindication of an autonomous American bryology. *Taxon* 68: 137–141.
- Medina, R., F. Lara, V. Mazimpaka & R. Garilleti. 2008. *Orthotrichum norrisii* (Orthotrichaceae), a new epiphytic Californian moss. *The Bryologist* 111: 670–675.
- Merrill, G. L. S. 1992. *Ozobryum ogalalense* (Pottiaceae), a new moss genus and species from the American Great Plains. *Novon* 2: 255–258.
- Merrill, G. L. S. 2007. *Polytrichum*. In: *Flora of North America Editorial Committee* (eds.), *Flora of North America* 27: 133–140. Oxford University Press, New York.
- Merrill, G. L. S. & R. R. Ireland, Jr. 2007. *Atrichum*. In: *Flora of North America Editorial Committee* (eds.), *Flora of North America* 27: 147–155. Oxford University Press, New York.
- Müller, F. 2014. *Bartramia aprica* — the correct name for the Mediterranean and western North American species historically recognized as “*Bartramia stricta*.” *Herzogia* 27: 211–214.
- Muñoz, J. 1999. A revision of *Grimmia* (Musci, Grimmiaceae) in the Americas. 1: Latin America. *Annals of the Missouri Botanical Garden* 86: 118–191.
- Muñoz, J. & F. Pando. 2000. A world synopsis of the genus *Grimmia* (Musci, Grimmiaceae). *Monographs in Systematic Botany from the Missouri Botanical Garden* 83: [i–viii] 1–133.
- Newton, A. E., C. J. Cox, J. G. Duckett, J. A. Wheeler, B. Goffinet, T. A. J. Hedderson & B. D. Mishler. 2000. Evolution of the major moss lineages: phylogenetic analyses based on multiple gene sequences and morphology. *The Bryologist* 103: 187–211.

- Norris, D. H. & J. Enroth. 1990. Description of *Bryolawtonia* Norris & Enroth (Thamnobryaceae), with observations on the genus *Bestia*. The Bryologist 93: 328–331.
- Norris, D. H. & J. R. Shevock. 2004a. Contributions toward a bryoflora of California: I. A specimen-based catalogue of mosses. Madroño 51: 1–131.
- Norris, D. H. & J. R. Shevock. 2004b. Contributions toward a bryoflora of California: II. A key to the mosses. Madroño 51: 133–269.
- Norris, D. H. & D. H. Vitt. 1993. *Orthotrichum spjutii* (Orthotrichaceae) a new species from the northern Sierra of California. Nova Hedwigia 56: 259–262.
- O'Brien, T. J. & D. G. Horton. 2000. *Bryochenea* (Musci: Thuidiaceae) is *Cyrto-hypnum* (Thuidiaceae), but *B. sachalinensis* is *Echinophyllum* (Thuidiaceae), a new genus from the Pacific Rim. The Bryologist 103: 509–517.
- Ochyra, R. 1989. Animadversions on the moss genus *Cratoneuron* (Sull.) Spruce. Journal of the Hattori Botanical Laboratory 67: 203–242.
- Ochyra, R. 2004. *Tortula hoppeana*, the correct name for *Desmatodon latifolius* in *Tortula* (Bryopsida, Pottiaceae). The Bryologist 107: 497–500.
- Ochyra, R. 2011. *Roellobryon* Ochyra, a new name for *Roellia* Kindb., *hom. illeg.* Polish Botanical Journal 56: 119–120.
- Ochyra, R. 2013. The new generic name *Torrentaria*, a nomenclatural consequence of the legitimacy of *Platyhypnum* and the illegitimacy of *Platyhypnidium* (Bryophyta). Nova Hedwigia 96: 205–212.
- Ochyra, R. & H. Bednarek-Ochyra. 1996. Nomenclatural notes on mosses: 13. *Cleistocarpidium*, genus novum (Ditrichaceae). Fragmenta Floristica et Geobotanica 41: 1034–1037.
- Ochyra, R. & H. Bednarek-Ochyra. 2007. *Bucklandiella*. In: Flora of North America Editorial Committee (eds.), Flora of North America 27: 267–285. Oxford University Press, New York.
- Ochyra, R. & A. Stebel. 2008. *Hylocomiadelphus*, a new genus of pleurocarpous mosses. Pages 303–308. In: A. Stebel & R. Ochyra (eds.), Bryophytes of the Polish Carpathians. Sorus, Poznań.
- Ochyra, R., J. Żarnowiec & H. Bednarek-Ochyra. 2003. Census catalogue of Polish mosses. Biodiversity of Poland 3: 1–372.
- Olsson, S., J. Enroth, V. Buchbender, L. Hedenäs, S. Huttunen & D. Quandt. 2011. *Neckera* and *Thamnobryum* (Neckeraceae, Bryopsida): paraphyletic assemblages. Taxon 60: 36–50.
- Orgaz, J. D., M. J. Cano & J. Guerra. 2013. Taxonomic revision of *Brachytheciastrum* (Brachytheciaceae, Bryophyta) from the Mediterranean region. Systematic Botany 38: 283–294.
- Patel, N. R., R. Medina, L. D. Williams, O. Lemieux, B. Goffinet & M. G. Johnson. 2023. Frequent allopolyploidy with distant progenitors in the moss genera *Physcomitrium* and *Entosthodon* (Funariaceae) identified via subgenome phasing of targeted nuclear genes. Evolution 77: 2561–2575.
- Pedersen, N. 2005. Validation of *Imbriobryum* (Bryaceae). The Bryologist 108: 449.
- Pedersen, N. & L. Hedenäs. 2003. Phylogenetic investigations of a well supported clade within the acrocarpous moss family Bryaceae: evidence from seven chloroplast DNA sequences and morphology. Plant Systematics and Evolution 240: 115–132.
- Pedersen, N. & L. Hedenäs. 2005. Taxonomic and nomenclatural implications of phylogenetic studies of the Bryaceae based on molecular data and morphology. The Bryologist 108: 123–128.
- Pedersen, N., D. T. Holyoak & A. E. Newton. 2007[2006]. Systematics and morphological evolution within the moss family Bryaceae: a comparison between parsimony and Bayesian methods for reconstruction of ancestral character states. Molecular Phylogenetics and Evolution 43: 891–907.
- Plášek, V., J. Sawicki, R. Ochyra, M. Szczecińska & T. Kulik. 2015. New taxonomical arrangement of the traditionally conceived genera *Orthotrichum* and *Ulota* (Orthotrichaceae, Bryophyta). Acta Musei Silesiae, Scientiae Naturales 64: 169–174.
- Pursell, R. A. 2017. A taxonomic revision of the Erpodiaceae (Bryophyta). Memoirs of the New York Botanical Garden 116: [i–viii] 1–106.
- Pursell, R. A. & D. M. Vital. 1986. Distributional adumbrations of *Fissidens* in the Neotropics. The Bryologist 89: 300–301.
- Rae, S. P. 2021. A preliminary specimen-based catalogue of the moss species found in Napa County, California. Madroño 68: 191–208.
- Redfearn, P. L. Jr. 1991. *Gymnostomiella* (Musci: Pottiaceae) in the Neotropics and eastern Asia. The Bryologist 94: 392–395.
- Reese, W. D. & B. E. Lemmon. 1965. A natural hybrid between *Weissia* and *Astomum* and notes on the nomenclature of the North American species of *Astomum*. The Bryologist 68: 27–288.
- Robinson, H. E. 1970. A revision of the moss genus, *Trichostomopsis*. Phytologia 20: 184–191.
- Robinson, H. E. & M. S. Ignatov. 1997. *Brachythecium laetum* B.S. G. the correct name for the species commonly known as *Brachythecium oxycladon*. The Bryologist 100: 359–361.
- Robinson, S. C., M. Nieto-Lugilde, A. M. Duffy, K. Martínez Muñoz, B. Agüero, A. Merced, K. Hassel, K. I. Flatberg & A. J. Shaw. 2024[2023]. The *Sphagnum cuspidatum* complex: phylogeny, species delimitation, and morphology. Biological Journal of the Linnean Society 141: 466–492.
- Ros, R. M. & I. Herrnstadt. 2010. New taxonomical data in the genus *Tortula* (Pottiaceae, Bryophyta). Tropical Bryology 31: 150–153.
- Ros, R. M., O. Werner & S. Rams Sánchez. 2005. New taxonomical data on the genus *Microbryum* Schimp. (Pottiaceae, Musci). Cryptogamie, Bryologie 26: 167–172.
- Saługa, M., R. Ochyra, J. Żarnowiec & M. Ronikier. 2018. Do Antarctic populations represent local or widespread phylogenetic and ecological lineages? Complicated fate of bipolar moss concepts with *Drepanocladus longifolius* as a case study. Organisms, Diversity & Evolution 18: 263–278.
- Sawicki, J., V. Plášek & M. Szczecińska. 2010. Molecular studies resolve *Nyholmiella* (Orthotrichaceae) as a separate genus. Journal of Systematics and Evolution 48: 183–194.
- Sawicki, J., M. Szczecińska, H. Bednarek-Ochyra & R. Ochyra. 2015. Mitochondrial phylogenomics supports splitting the traditionally conceived genus *Racomitrium* (Bryophyta: Grimmiaceae). Nova Hedwigia 100: 293–317.
- Schlesak, S., L. Hedenäs, M. Nebel & D. Quandt. 2018. Cleaning a taxonomic dustbin: placing the European *Hypnum* species in a phylogenetic context! Bryophyte Diversity & Evolution 40: 37–54.
- Schofield, W. B. 2006. *Brotherella canadensis* sp. nov., from the Pacific coast of North America. Journal of the Hattori Botanical Laboratory 100: 355–360.
- Schofield, W. B. 2014. *Hypnum*. In: Flora of North America Editorial Committee (eds.), Flora of North America 28: 532–549. Oxford University Press, New York.
- Shaw, A. J. 2009. *Mielichhoferia shevockii*, comb. nov. (Mielichhoferiaceae). The Bryologist 112: 605.

- Shaw, A. J. 2014. Mielichhoferiaceae. In: Flora of North America Editorial Committee (eds.), Flora of North America 28: 189–214. Oxford University Press, New York.
- Shaw, A. J., K. I. Flatberg, P. Szövényi, M. Ricca, M. G. Johnson, H. K. Stenøien & B. Shaw. 2012a. Systematics of the *Sphagnum fimbriatum* complex: phylogenetic relationships, morphological variation, and allopolyploidy. *Systematic Botany* 37: 15–30.
- Shaw, A. J., B. Shaw, M. G. Johnson, N. Devos & B. Carter. 2015. Genetic and phylogenetic structure of the Pacific Rim clade of *Sphagnum* subg. *Subsecunda*: haploid and allodiploid taxa. *Biological Journal of the Linnean Society* 116: 295–311.
- Shaw, A. J., B. Shaw, M. Ricca & K. I. Flatberg. 2012b. A phylogenetic monograph of the *Sphagnum subsecundum* complex (Sphagnaceae) in eastern North America. *The Bryologist* 115: 128–152.
- Shaw, A. J. & D. R. Toren. 2009. *Pohlia flexuosa* (Mielichhoferiaceae) in North America. *The Bryologist* 112: 178–183.
- Shaw, B., S. Terracciano & A. J. Shaw. 2009. A genetic analysis of two recently described peat moss species, *Sphagnum atlanticum* and *S. bergianum* (Sphagnaceae). *Systematic Botany* 34: 1–9.
- Shevock, J. R. 2000. Two mosses new for North America with Southern Hemisphere affinities. *Evansia* 17: 97–98.
- Shevock, J. R. & B. Aguero. 2023. *Philonotis breuteloides* (Bartramiaceae, Bryophyta), a new montane fen moss species endemic to California. *Madroño* 69: 349–358.
- Shevock, J. R. & W. R. Buck. 2009. A new species of *Pseudoleskea* (Leskeaceae) from the Sierra Nevada of California, U.S.A. *The Bryologist* 112: 184–187.
- Shevock, J. R. & D. H. Norris. 2014. *Scouleria siskiyouensis* (Scouleriaceae), a new rheophytic moss endemic to southern Oregon, USA. *Madroño* 61: 137–143.
- Shevock, J. R. & A. J. Shaw. 2005. *Pohlia robertsonii* and *P. rabunbaldensis* (Bryopsida, Mniaceae), two new species from the western and eastern United States. *The Bryologist* 108: 177–182.
- Shevock, J. R., D. R. Toren & D. H. Wagner. 2021. Bryoflora of the Russian Wilderness and adjacent slopes of the Salmon Mountains, Klamath National Forest, Siskiyou County, California. *Madroño* 68: 209–256.
- Smith, D. K. & P. G. Davison. 1993. Antheridia and sporophytes in *Takakia ceratophylla* (Mitt.) Grolle: evidence for reclassification among mosses. *Journal of the Hattori Botanical Laboratory* 73: 263–271.
- Snyder, E. A. 2023. A small addition to the flora of North America: *Ephemerum recurvifolium* (Dickson) Boulay (Bryopsida: Potentiales). *Evansia* 40: 40–45.
- Solano, I., D. Quandt, D. Zumel & J. Muñoz. 2023. *Grimmia shevockii* and *G. insolita*, two new species endemic to California. *Journal of Bryology* 45: 219–235.
- Spence, J. R. 1996. *Rosulabryum* genus novum (Bryaceae). *The Bryologist* 99: 221–225.
- Spence, J. R. 2005. New genera and combinations in Bryaceae (Bryales, Musci) for North America. *Phytologia* 87: 15–28.
- Spence, J. R. 2007. Four species of the Bryaceae new to the U.S.A. *Evansia* 24: 29–30.
- Spence, J. R. 2014. Bryaceae. In: Flora of North America Editorial Committee (eds.), Flora of North America 28: 117–185. Oxford University Press, New York.
- Spence, J. R., J. C. Brinda, S. M. Buckley & N. J. Trigoboff. 2022. New records of Bryaceae (Bryophyta) for the *Flora of North America* region, including a major range extension from Europe. *Evansia* 39: 87–96.
- Spence, J. R. & K. M. Kellman. 2015. New and interesting species of *Gemmabryum* J.R.Spence & H.P.Ramsay (Bryaceae, Bryopsida) from California and the West. *Madroño* 62: 124–135.
- Spence, J. R. & H. P. Ramsay. 2005. New genera and combinations in the Bryaceae (Bryales, Musci) for Australia. *Phytologia* 87: 61–72.
- Spence, J. R. & J. R. Shevock. 2012. *Ptychostomum pacificum* (Bryaceae), a new fen species from California, Oregon, and western Nevada, USA. *Madroño* 59: 156–162.
- Spence, J. R. & J. R. Shevock. 2015. *Imbriobryum torenii* (Bryaceae), a new species from western North America. *Madroño* 62: 68–72.
- Stech, M., O. Werner, J. M. González-Mancebo, J. Patiño, M. Sim-Sim, S. Fontinha, I. Hildebrandt & R. M. Ros. 2011. Phylogenetic inference in *Leucodon* Schwägr. subg. *Leucodon* (Leucodontaceae, Bryophyta) in the North Atlantic region. *Taxon* 60: 79–88.
- Stehn, S. E. & D. Kofranek. 2014. *Trematodon laetevirens* (Bruchiaceae) new to the United States, and other interesting bryophyte records from Wrangell-St. Elias National Park and Preserve, Alaska. *Evansia* 31: 93–98.
- Streiff, A. 2006. Phylogenetic study of *Grimmia* (Grimmiaceae) based on plastid DNA sequences (*trnL-trnF* and *rps4*) and on morphological characters. *The Bryologist* 109: 224–235.
- Sullivant, W. S. 1849. Contributions to the bryology and hepaticology of North America. Part II. *Memoirs of the American Academy of Arts and Sciences*, new series, 4: 169–176.
- Tan, B. C. 1979. A new classification for the genus *Physcomitrella*. *Journal of the Hattori Botanical Laboratory* 46: 327–336.
- Tan, B. C. & Y. Jia. 1999. A preliminary revision of Chinese Sematophyllaceae. *Journal of the Hattori Botanical Laboratory* 86: 1–70.
- Toren, D. R. 2008. *Funaria convexa* Spruce (Musci: Funariaceae) new to North America. *Evansia* 25: 94–96.
- Toren, D. R. 2015. A moss flora of Lake County, California. *Madroño* 62: 241–268.
- Toren, D. R. & K. Heise. 2009. *Bryum chryseum* Mitt. (Musci: Bryaceae) new to North America north of Mexico. *Evansia* 26: 98–101.
- Toren, D. R., K. M. Kellman & J. R. Shevock. 2016. *Archidium crassicutatum* (Archidiaceae), a new and long-overlooked species from California, U.S.A. *Madroño* 63: 348–352.
- Toren, D. R., J. R. Spence, J. R. Shevock & J. Larráin. 2024. *Bryum flabelliforme* (Bryaceae, Bryophyta), a North American species with unusual rhizoidal tubers and a new name for the austral *Splachnobryum valdiviae*. *Madroño* 71(4) in press.
- Touw, A. 2001. A review of the Thuidiaceae (Musci) and a realignment of taxa traditionally accommodated in *Thuidium* sensu amplo (*Thuidium* Schimp., *Thuidiopsis* (Broth.) M.Fleisch., and *Pelekium* Mitt.) including *Aequatoriella* gen. nov., and *Indothuidium* gen. nov. *Journal of the Hattori Botanical Laboratory* 90: 167–209.
- Tubeanova, D. Ya., D. V. Goryunov, E. A. Ignatova & M. S. Ignatov. 2010. On the taxonomy of *Dicranum acutifolium* and *D. fuscescens* complexes (Dicranaceae, Bryophyta) in Russia. *Arctoa* 19: 151–164.
- Turland, N. J., J. H. Wiersema, F. R. Barrie, W. Greuter, D. L. Hawksworth, P. S. Herendeen, S. Knapp, W.-H. Kusber, D.-Z. Li, K. Marhold, T. W. May, J. McNeill, A. M. Monro, J. Prado, M. J. Price & G. F. Smith (eds.). 2018. International Code of Nomenclature for algae, fungi, and plants (Shenzhen Code) adopted by the Nineteenth International Botanical Congress,

- Shenzhen, China, July, 2017. *Regnum Vegetabile* 159: i–xxxviii, 1–254.
- Valente, D. V., D. F. Peralta, R. X. A. Prudêncio & P. E. A. S. Câmara. 2020. Taxonomic notes and new synonyms on Brazilian *Macromitrium* Bridel (Bryophyta, Orthotrichaceae). *Phytotaxa* 454: 213–225.
- Vanderpoorten, A. & L. Hedenäs. 2009. New combinations in the Amblystegiaceae. *Journal of Bryology* 31: 129–132.
- Vanderpoorten, A., L. Hedenäs, C. J. Cox & A. J. Shaw. 2002. Phylogeny and morphological evolution of the Amblystegiaceae (Bryopsida). *Molecular Phylogenetics and Evolution* 23: 1–21.
- Vigalondo, B., I. Draper, V. Mazimpaka, J. A. Calleja, F. Lara & R. Garilleti. 2020. The *Lewinskya affinis* complex (Orthotrichaceae) revisited: species description and differentiation. *The Bryologist* 123: 455–482.
- Vigalondo, B., J. Patiño, I. Draper, V. Mazimpaka, J. R. Shevock, A. Losada-Lima, J. M. González-Mancebo, R. Garilleti & F. Lara. 2019. The long journey of *Orthotrichum shevockii* (Orthotrichaceae, Bryopsida): from California to Macaronesia. *PLoS ONE* 14(2): e0211017 [1–24].
- Vitt, D. H. 2014. *Ulota*. In: *Flora of North America Editorial Committee* (eds.), *Flora of North America* 28: 72–78. Oxford University Press, New York.
- Wang, C.-Y. & J.-C. Zhao. 2009. Phylogeny of *Ptychostomum* (Bryaceae, Musci) inferred from sequences of nuclear ribosomal DNA internal transcribed spacer (ITS) and chloroplast *rps4*. *Journal of Systematics and Evolution* 47: 311–320.
- Weber, W. A., R. C. Wittmann, & R. Worthington. 2003. *Grimmia bernoullii* Mueller Hal. in the United States. *Evansia* 20: 104–105.
- Werner, O., H. Köckinger, M. Magdy & R. M. Ros. 2014. On the systematic position of *Tortella arctica* and *Trichostomum arcticum* (Bryophyta, Pottiaceae). *Nova Hedwigia* 98: 273–293.
- Werner, O., S. Rams, J. Kučera, J. Larraín, O. M. Afonina, S. Pisa & R. M. Ros. 2013. New data on the moss genus *Hymenoloma* (Bryophyta), with special reference to *H. mulahaceni*. *Cryptogamie, Bryologie* 34: 13–30.
- Whitehouse, H. L. K. 2001[2000]. Bryophytes of arable fields in Québec and Slovakia, including new records of *Bryum demartianum* Arts. *Lindbergia* 26: 29–32.
- Wilson, K. L. 2021. Report of the General Committee: 22. *Taxon* 70: 1360–1362.
- Wilson, K. L. 2024. Report of the General Committee: 29. *Taxon* 73: 619–621.
- Wilson, P. S. & D. H. Norris. 1989. *Pseudoleskeella* in North America and Europe. *The Bryologist* 92: 387–396.
- Wilson, W. 1857. Enumeration of the mosses collected in India by Dr. J. D. Hooker, F.R.S., and Dr. Thomas Thomson, F.R.S., with their habitats, elevations, and the numbers under which they have been distributed; arranged by W. Mitten. *Hooker's Journal of Botany and Kew Garden Miscellany* 9: 289–300, 321–333, 363–470.
- Wolski, G. J. 2020. Reassessing the taxonomic diversity of *Plagiothecium* section *Orthophyllum* in the North American bryoflora. *Brittonia* 72: 337–350.
- Wolski, G. J., D. Bożyk & J. Proćków. 2022a. Revision of the original material of *Plagiothecium denticulatum* var. *obtusifolium* (Turn.) Moore and new synonyms for this taxon. *Plants* 11(19): 2446 [1–20].
- Wolski, G. J., A. Faltyn-Parzymyska & J. Proćków. 2020. Lectotypification of the name *Stereodon nemoralis* Mitt. (Plagiotheciaceae), a basionym of *Plagiothecium nemorale* (Mitt.) A.Jaeger. *PhytoKeys* 155: 141–153.
- Wolski, G. J., M. Latoszewski & W. R. Buck. 2024. Why *Plagiothecium sylvaticum* (Brid.) Schimp. (Plagiotheciaceae) has priority over *P. platyphyllum* Mönk.? *PhytoKeys* 241: 103–120.
- Wolski, G. J., P. Nowicka-Krawczyk & W. R. Buck. 2021. *Plagiothecium schofieldii*, a new species from the Aleutian Islands (Alaska, USA). *PhytoKeys* 184: 127–138.
- Wolski, G. J., P. Nowicka-Krawczyk & W. R. Buck. 2022b. *Plagiothecium talbotii*, a new species from the Aleutian Islands (Alaska, U.S.A.). *PhytoKeys* 194: 63–73.
- Wolski, G. J., P. Nowicka-Krawczyk & W. R. Buck. 2022c. Taxonomic revision of the *Plagiothecium curvifolium* complex. *PLoS ONE* 17(11): e0275665 [1–24].
- Worley, I. A. & Z. Iwatsuki. 1970. A checklist of the mosses of Alaska. *The Bryologist* 73: 59–71.
- Wyatt, R. E., E. S. J. Harris & I. J. Odrzykoski. 2021a. The pantropical moss *Plagiomnium rhynchophorum* (Mniaceae) disjunct to the Southern Appalachians and Hawaii. *The Bryologist* 124: 343–351.
- Wyatt, R. E. & I. J. Odrzykoski. 2012. *Plagiomnium floridanum* sp. nov. (Mniaceae), a new moss from the southeastern United States. *The Bryologist* 115: 527–535.
- Wyatt, R. E., I. J. Odrzykoski & A. H. Stoneburner. 1993. Isozyme evidence regarding the origins of the allopolyploid moss *Plagiomnium curvatulum*. *Lindbergia* 18: 49–58.
- Wyatt, R. E., I. J. Odrzykoski & A. H. Stoneburner. 2023. Isozyme evidence regarding relationships within *Rhizomnium* (Mniaceae). *The Bryologist* 126: 45–51.
- Wyatt, R. E. & A. H. Stoneburner. 2018. Morphological studies show that *Pleurochaete luteola* is distinct from “*P. squarrosa*” (Pottiaceae) in North America. *Hattoria* 9: 1–10.
- Wyatt, R. E. & A. H. Stoneburner. 2021. *Cinclidium minutifolium* (Mniaceae) in Alaska: confirmation and clarification. *The Bryologist* 124: 85–89.
- Wyatt, R. E., A. H. Stoneburner & I. J. Odrzykoski. 2021b. *Cinclidium alaskanum* sp. nov. (Mniaceae), a new moss from the North Slope of Alaska. *The Bryologist* 124: 1–8.
- Wynns, J. T. 2006. Taxonomic Studies in the Aquatic Moss Genus *Platyhypnidium* (Brachytheciaceae). M.Sc. thesis, Appalachian State University, Boone, North Carolina.
- Wynns, J. T. 2015. Molecular Phylogeny and Systematic Revision of the Pleurocarpus Moss Genus *Plagiothecium*. Doctoral Dissertation. University of Copenhagen, Copenhagen, Denmark.
- Wynns, J. T. 2020. New names and new combinations for hypnalean mosses. *The Bryologist* 123: 633–656.
- Wynns, J. T., K. Rysbjerg Munk & C. B. Asmussen Lange. 2018 [2017]. Molecular phylogeny of *Plagiothecium* and similar hypnalean mosses, with a revised sectional classification of *Plagiothecium*. *Cladistics* 34: 469–501.
- Wynns, J. T. & C. Schröck. 2018. Range extensions for the rare moss *Plagiothecium handelii*, and its transfer to the resurrected genus *Ortholimnobia*. *Lindbergia* 41(1): 01087 [1–7].
- Yip, K. L. 2000. A Revision of the Genus *Pleuridium* (Musci). Doctoral Dissertation. University of Cincinnati, Cincinnati, Ohio, U.S.A.
- Yip, K. L. 2002. *Pseudephemerum*, new to the United States and Mexico. *The Bryologist* 105: 256–258.

- Yip, K. L., P. S. Wilson, D. R. Toren & K. M. Kellman. 2007. *Pleuri-dium mexicanum* (Ditrichaceae) new to the USA from California. *The Bryologist* 110: 510–513.
- Zander, R. H. 1989. Seven new genera in Pottiaceae (Musci) and a lectotype for *Syntrichia*. *Phytologia* 65: 424–436.
- Zander, R. H. 1993. Genera of the Pottiaceae: mosses of harsh environments. *Bulletin of the Buffalo Society of Natural Sciences* 32: i–vi, 1–378.
- Zander, R. H. 1995. Phylogenetic relationships of *Hyophiladelphus* gen. nov. (Pottiaceae, Musci) and a perspective on the cladistic method. *The Bryologist* 98: 363–374.
- Zander, R. H. 2007a. *Barbula*. In: Flora of North America Editorial Committee (eds.), *Flora of North America* 27: 528–534. Oxford University Press, New York.
- Zander, R. H. 2007b. *Gymnostomiella*. In: Flora of North America Editorial Committee (eds.), *Flora of North America* 27: 609–610. Oxford University Press, New York.
- Zander, R. H. 2007c. *Gyroweisia*. In: Flora of North America Editorial Committee (eds.), *Flora of North America* 27: 523–525. Oxford University Press, New York.
- Zander, R. H. 2007d. *Microbryum*. In: Flora of North America Editorial Committee (eds.), *Flora of North America* 27: 627–631. Oxford University Press, New York.
- Zander, R. H. 2007e. *Pseudocrossidium*. In: Flora of North America Editorial Committee (eds.), *Flora of North America* 27: 569–572. Oxford University Press, New York.
- Zander, R. H. 2007f. *Tortula*. In: Flora of North America Editorial Committee (eds.), *Flora of North America* 27: 586–603. Oxford University Press, New York.
- Zander, R. H. 2007g. *Trichostomum*. In: Flora of North America Editorial Committee (eds.), *Flora of North America* 27: 488–494. Oxford University Press, New York.
- Zander, R. H. 2007h. *Weissia*. In: Flora of North America Editorial Committee (eds.), *Flora of North America* 27: 512–518. Oxford University Press, New York.
- Zander, R. H. 2013. A Framework for Post-Phylogenetic Systematics. [iv] 209 pp. Zetetic Publications, St. Louis, MO.
- Zander, R. H. 2017. *Anoetangium sikkimense* (Pottiaceae, Bryophyta) new to the New World from Alaska, and its macroevolutionary relationships. *The Bryologist* 120: 435–440.
- Zander, R. H. 2021. A new species of *Hyophila* (Pottiaceae, Bryophyta), *Teniolophora* synonymized with *Hyophila*, and *Plaubelia* clarified. *The Bryologist* 124: 469–474.
- Zander, R. H. 2023. *Fractal Evolution, Complexity and Systematics*. Zetetic Publications, St. Louis, Missouri.
- Zander, R. H. 2024. Minimally monophyletic genera are the cast-iron building blocks of evolution. *Ukrainian Botanical Journal* 81: 87–99.
- Zander, R. H. & R. T. Caners. 2017. *Didymodon constrictus* (Pottiaceae, Bryophyta) new to the New World from Alberta, Canada. *The Bryologist* 120: 287–292.
- Zander, R. H. & P. M. Eckel. 2007. *Anoetangium*. In: Flora of North America Editorial Committee (eds.), *Flora of North America* 27: 521–523. Oxford University Press, New York.
- Zander, R. H. & P. M. Eckel. 2017. Notes on Pleuroweisiae (Pottiaceae, Bryophyta) in North America. *Evansia* 34: 61–64.
- Zander, R. H. & P. M. Eckel. 2019. *Pseudosymblypharis* and *Oxystegus* species (Pottiaceae) new to northwestern North America, and refutation of molecular *Chionoloma*. *The Bryologist* 122: 568–577.
- Zander, R. H. & T. A. J. Hedderson. 2016. Re-evaluation of *Hymenostylium xanthocarpum* (Hook.) Brid., and *Ardeuma* R.H. Zander & Hedd., a new name for all other species of *Hymenostylium* (Pottiaceae, Bryophyta). *Journal of Bryology* 38: 295–301.
- Zander, R. H. & W. A. Weber. 1997. *Didymodon anserinocapitatus* (Musci, Pottiaceae) new to the New World. *The Bryologist* 100: 237–238.
- Żarnowiec, J. 2001. A Taxonomic Monograph of the *Drepanocladus aduncus* Group (Bryopsida: Amblystegiaceae). Bielsko-Biała, Poland: Łódź Technical University, Bielsko-Biała Branch. xii + 248 pp.

manuscript received October 16, 2024; accepted October 24, 2024.

Supplementary documents online:

Supplementary Table S1. Alphabetical list of genera, species, subspecies, varieties and forms of mosses from the continental United States and Canada and their familial affiliations.

Supplementary Table S2. Alphabetical list of families of mosses from the continental United States and Canada and their generic inclusions.

Supplementary Table S3. Alphabetical list of families of mosses from the continental United States and Canada and their ordinal affiliations.

Supplementary Table S4. Alphabetical list of orders of mosses from the continental United States and Canada and their familial inclusions.