



Ontogenetic dietary shifts in the summer feeding intensity of brown trout in relation to fish condition

Authors: Sánchez-Hernández, Javier, and Cobo, Fernando

Source: Folia Zoologica, 62(2) : 110-114

Published By: Institute of Vertebrate Biology, Czech Academy of Sciences

URL: <https://doi.org/10.25225/fozo.v62.i2.a4.2013>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Ontogenetic dietary shifts in the summer feeding intensity of brown trout in relation to fish condition

Javier SÁNCHEZ-HERNÁNDEZ^{1,2*} and Fernando COBO^{1,2}

¹ Department of Zoology and Physical Anthropology, Faculty of Biology, University of Santiago de Compostela, Campus Sur s/n, 15782 Santiago de Compostela, Spain; e-mail: javier.sanchez@usc.es

² Station of Hydrobiology “Encoro do Con”, Castroagudín s/n, 36617 Vilagarcía de Arousa, Pontevedra, Spain

Received 14 November 2012; Accepted 3 April 2013

Abstract. We examined condition factor, hepatosomatic index and stomach fullness in brown trout to study if feeding intensity can be related to fish condition. Trout were collected at three locations during the summer in temperate rivers (Galicia, NW Spain). Our findings suggest that the feeding intensity is inversely related with the fish condition because the stomach fullness decreases with fish age and size and the condition factor is the lowest in young-of-the-year (YOY). In general, no significant differences among age classes were found in the hepatosomatic index, except in one river (the River Lengüelle) in which YOY shows the highest value. The high feeding intensity of YOY during summer could be related with the increases in fish condition and survival in the later autumn and winter.

Key words: *Salmo trutta*, feeding activity, condition factor, hepatosomatic index, ontogeny

Introduction

The intake of food by fishes has a great importance to the fish development, and when the daily intake is above the maintenance level, the total energy content of the fish increases and growth occurs (Elliott 1994). Hence, several studies emphasize the importance of the feeding intensity in the acquisition of energetic reserves (e.g., Rikardsen et al. 2006), feature that is especially important in juveniles because the corporal reserve is higher in older fishes (e.g., Nomura et al. 2001).

Studies of the feeding ecology of *S. trutta* are frequent across the world (e.g., Jensen et al. 2004, Fochetti et al. 2008, Sánchez-Hernández et al. 2011), and the fullness index have been widely used to study the feeding intensity in salmonids (Kara & Alp 2005, Shustov et al. 2012, Cobo et al. 2013). Although the shifts in the feeding behaviour of brown trout during the ontogeny are well documented (e.g., Fochetti et al. 2008, Sánchez-Hernández & Cobo 2012, Sánchez-Hernández et al. 2013), other aspects of its feeding behaviour such as the ontogenetic shifts in the feeding intensity remain unclear. In this context, Kara & Alp (2005) found that the brown trout between 5.7 cm and 32 cm fed most intensively, whilst the intensity declined above 32 cm length. So, the aims of the present study were (1) to study if the feeding intensity

of brown trout varies with age during the summer and (2) to verify whether this difference in the feeding intensity among age classes could be related to fish condition.

Material and Methods

Individuals were collected using pulsed D.C. backpack electrofishing equipment (Hans Grassl GmbH, ELT60II) following the standardised electric fishing procedures described in the European CEN directive Water analysis – Fishing with electricity (EN 14011; CEN 2003) for wadable rivers. Brown trout were collected in three wadable riffle sections in oligotrophic rivers draining granitic catchments in Galicia (NW Spain) (Martínez-Ansemil & Membiela 1992) during three consecutive days of September 2007: the River Anllóns (43°13' N and 8°53' W), the River Furelos (42°52' N and 8°01' W) and the River Lengüelle (42°58' N and 8°27' W).

A total of 139 individuals were captured (Table 1), measured to the nearest 1 mm and weighed to the nearest 1 g. Age of each individual was determined by scale reading. In the laboratory, individuals were dissected in order to extract the stomach and the liver. The condition factor (*CF*) was calculated using the formula $CF = 100 W/L^3$, where *W* is the wet weight (g) before evisceration and *L* is the fork length (cm).




Fig. 1. Plots of curve estimation procedure between variables (condition factor, hepatosomatic index and stomach fullness index) and fork length in total using pooled data for brown trout. Observed (dots) and cubic regression model (lines).

Hepatosomatic index (*HSI*) was calculated as: $HSI = (W_l/W_b) \times 100$, where W_l is the wet liver weight and W_b is the wet body weight. The stomach fullness index (*f*) was calculated as $f = (W_s/W) \times 100$, where W_s is the total stomach content mass (g) and W is the fish mass (g).

The non-parametric Kruskal-Wallis tests for non-normal data were used for detecting differences in the *CF*, *HSI* and *f* among age classes in each studied river and in total using pooled data. In order to explore the possibility of a nonlinear relationship between variables (*CF*, *HSI* and *f*) and fork length the curve estimation procedure was used, which compared 11 different models (linear, quadratic, cubic, exponential...). The model with the highest adjusted R^2 was chosen. All tests were considered statistically significant at p level < 0.05 . Statistical analyses were conducted using the programme IBM SPSS Statistics 20.

Results

Regarding morphological indices, in spite of *CF* was lower in young-of-the-year (YOY) than the other age classes in the three studied rivers, the significant differences were found only in the River Furelos (Table 2). Opposite, only significant differences in the *HSI* were found in the River Lengüelle in which the age-0+ shows the highest value (Table 2). With pooled data (total), differences among age classes were significant for *CF* and *HSI* (Table 2).

In the three studied rivers, results of the stomach fullness index showed high variations among age classes, being higher in age-0+ than the other age classes (Table 2). According to the curve estimation procedure the relationship between variables (*CF*, *HSI* and *f*) and fork length was best described by a cubic regression model (Fig. 1). As expected, the relation between the stomach fullness and fish size was negative in all cases. However, only in the River Furelos the relationship between *CF* and *HSI* and fish length was significant, being the correlation positive for *CF* and negative for *HSI* (Table 3).

Discussion

The energetic reserves in the body of the fish changes depending on the age and season (e.g., Nomura et al. 2001, Rikardsen et al. 2006). Several researchers have demonstrated that the condition factor and the hepatosomatic index serve to assess the overall fish condition, providing information on energy reserves (Herbinger & Friars 1991, Van der Oost et al. 2003, Cobo et al. 2013). In spite that statistical differences between rivers and age-class might be due to the differences in sample size; in our case, only in the River Lengüelle the *HSI* decreases with fish age. Results that may reflect the storage of glycogen in the liver in YOY as a result of the higher food intake, and similar than studies about starvation and refeeding under laboratory conditions (e.g., Machado et al.

Table 1. Size and the number of brown trout used in the present study. Mean \pm standard error (SE). Data are presented for each age class and in total.

	Age-0+	Age-1+	Age-2+	Age-3+	Age-4+	Total
the River Anllóns						
Mean size (cm)	7.9 \pm 1.45	14.7 \pm 0.48	19.2 \pm 0.52	28.5 \pm 0.39	-	17.8 \pm 0.99
Maximum size (cm)	9.4	17.9	22.3	30	-	30
Minimum size (cm)	6.5	10.9	17.7	27	-	6.5
Number of trout (<i>n</i>)	2	18	9	6	0	35
the River Furelos						
Mean size (cm)	6.6 \pm 0.16	14.5 \pm 0.32	21.7 \pm 1.07	26.2 \pm 0.20	34.6	11.5 \pm 0.81
Maximum size (cm)	8.7	17	24	26.6	-	34.6
Minimum size (cm)	4.8	12	18	26	-	4.8
Number of trout (<i>n</i>)	37	20	5	3	1	66
the River Lengüelle						
Mean size (cm)	6.6 \pm 0.19	12.7 \pm 0.19	19.7 \pm 0.49	25.5 \pm 0.42	-	16.1 \pm 0.96
Maximum size (cm)	7.2	13.9	22.9	26.8	-	26.8
Minimum size (cm)	5.9	11.7	16.5	25	-	5.9
Number of trout (<i>n</i>)	6	12	16	4	0	38
Total number of trout (<i>n</i>)	45	50	30	13	1	

Table 2. Condition factor (*CF*), hepatosomatic index (*HSI*) and stomach fullness index (*f*). Mean \pm standard error (SE). Data are presented for each age class. Kruskal-Wallis test for comparisons between age classes within rivers and rivers within age classes. Significant differences are marked in bold.

	Age-0+	Age-1+	Age-2+	Age-3+	Age-4+	Kruskal-Wallis test
Condition factor						
the River Anllóns	1.24 \pm 0.049	1.35 \pm 0.025	1.32 \pm 0.032	1.25 \pm 0.008	-	<i>H</i> = 6.48; <i>p</i> = 0.090
the River Furelos	1.10 \pm 0.025	1.33 \pm 0.172	1.34 \pm 0.031	1.26 \pm 0.069	1.34	<i>H</i> = 30.92; <i>p</i> < 0.001
the River Lengüelle	1.29 \pm 0.058	1.43 \pm 0.037	1.36 \pm 0.018	1.31 \pm 0.030	-	<i>H</i> = 5.68; <i>p</i> = 0.128
Total (pooled data)	1.13 \pm 0.023	1.36 \pm 0.015	1.34 \pm 0.014	1.27 \pm 0.018	1.34	<i>H</i> = 46.90; <i>p</i> < 0.001
Kruskal-Wallis test	<i>H</i> = 8.09; <i>p</i> = 0.017	<i>H</i> = 4.96; <i>p</i> = 0.084	<i>H</i> = 0.58; <i>p</i> = 0.746	<i>H</i> = 2.06; <i>p</i> = 0.357	-	-
Hepatosomatic index						
the River Anllóns	0.63 \pm 0.070	0.66 \pm 0.100	0.67 \pm 0.050	0.58 \pm 0.200	-	<i>H</i> = 3.49; <i>p</i> = 0.322
the River Furelos	0.66 \pm 0.05	0.64 \pm 0.028	0.59 \pm 0.092	0.48 \pm 0.091	0.24	<i>H</i> = 4.97; <i>p</i> = 0.290
the River Lengüelle	1.05 \pm 0.156	0.61 \pm 0.054	0.78 \pm 0.057	0.66 \pm 0.099	-	<i>H</i> = 9.49; <i>p</i> = 0.023
Total (pooled data)	0.71 \pm 0.049	0.64 \pm 0.039	0.72 \pm 0.038	0.58 \pm 0.039	0.24	<i>H</i> = 24.04; <i>p</i> < 0.001
Kruskal-Wallis test	<i>H</i> = 7.10; <i>p</i> = 0.029	<i>H</i> = 4.27; <i>p</i> = 0.118	<i>H</i> = 3.47; <i>p</i> = 0.176	<i>H</i> = 1.09; <i>p</i> = 0.577	-	-
Stomach fullness						
the River Anllóns	5.20 \pm 2.979	0.72 \pm 0.092	0.85 \pm 0.433	0.11 \pm 0.034	-	<i>H</i> = 19.67; <i>p</i> < 0.001
the River Furelos	9.02 \pm 0.643	1.09 \pm 0.139	1.00 \pm 0.246	1.15 \pm 0.24	1.36	<i>H</i> = 48.29; <i>p</i> < 0.001
the River Lengüelle	1.81 \pm 0.958	1.02 \pm 0.129	0.57 \pm 0.204	0.16 \pm 0.050	-	<i>H</i> = 19.51; <i>p</i> = 0.001
Total (pooled data)	7.9 \pm 0.665	0.94 \pm 0.074	0.73 \pm 0.171	0.36 \pm 0.135	1.36	<i>H</i> = 125.5; <i>p</i> < 0.001
Kruskal-Wallis test	<i>H</i> = 13.32; <i>p</i> = 0.01	<i>H</i> = 6.79; <i>p</i> = 0.033	<i>H</i> = 4.63; <i>p</i> = 0.099	<i>H</i> = 8.65; <i>p</i> = 0.013	-	-

Table 3. Spearman's rank correlation (R^2) according to the curve estimation procedure between variables (condition factor, hepatosomatic index and stomach fullness index) and fish length. Data are presented for each river and in total (pooled data). Significant differences are marked in bold.

	R^2	p -value	Type of regression model	Regression model equation
Condition factor				
the River Anllóns	0.15	0.177	Cubic	$y = 0.069x^3 - 0.003x^2 + 0.001x + 0.907$
the River Furelos	0.52	< 0.001	Cubic	$y = 0.001x^3 - 0.005x^2 + 0.166x + 0.553$
the River Lengüelle	0.11	0.274	Cubic	$y = 0.001x^3 - 0.005x^2 + 0.085x + 0.928$
Total (pooled data)	0.40	< 0.001	Cubic	$y = 0.001x^3 - 0.006x^2 + 0.124x + 0.554$
Hepatosomatic index				
the River Anllóns	0.04	0.698	Cubic	$y = -0.001x^3 + 0.005x^2 - 0.109x + 1.453$
the River Furelos	0.06	0.039	Linear	$y = -0.010x + 0.746$
the River Lengüelle	0.41	< 0.001	Cubic	$y = -0.001x^3 + 0.036x^2 - 0.575x + 3.507$
Total (pooled data)	0.07	0.026	Cubic	$y = -0.001x^3 + 0.008x^2 - 0.130x + 1.282$
Stomach fullness				
the River Anllóns	0.70	< 0.001	Cubic	$y = -0.003x^3 + 0.189x^2 - 3.692x + 23.838$
the River Furelos	0.86	< 0.001	Cubic	$y = -0.004x^3 + 0.238x^2 - 4.899x + 32.214$
the River Lengüelle	0.38	< 0.001	Exponential	$y = -0.121^x + 3.330$
Total (pooled data)	0.76	< 0.001	Cubic	$y = -0.003x^3 + 0.199x^2 - 4.116x + 27.521$

1988). As expected, the correlation between CF and fish length was positive in the River Furelos and in total (pooled data), and also CF changes depending on the age as it can be seen in Table 2. These findings are in agreement with previous studies, and in salmonids, as in many other fish species; there is normally a shift in the condition factor and energetic reserves during the ontogeny, being lower in young fishes (e.g., Nomura et al. 2001).

Our results demonstrate that the feeding intensity decreases with the fish age and length. However, as proved by the results obtained by several authors, for brown trout, as in many other salmonids species, the stomach fullness may vary considerably during the year (e.g., Lagarrigue et al. 2002, Kara & Alp 2005). So, under natural conditions, stomach fullness index was the lowest in autumn, then increased from winter to summer (Lagarrigue et al. 2002). In our case, surveys was carried out at the end of the summer season (September) coinciding with the low water level of the studied rivers, and the interpretation of the results of this study should take into account the phenology of feeding intensity. Indeed, although the data of the present study is limited to a September, the end of the summer is the most interesting moment to survey because energetic reserves at this moment will be determinant for the survival on the coming winter.

The acquisition of energetic reserves in fishes is extremely linked with the feeding intensity (e.g., Rikardsen et al. 2006). Studies developed in brown

trout have demonstrated that fishes between 5.7 cm and 32 cm fed most intensively, whilst the intensity declined above 32 cm length (Kara & Alp 2005). In contrast with the findings found by Kara & Alp (2005), in our case the feeding intensity was the highest in YOY (with sizes between 4.8 and 9.4 cm) and declined drastically in the age-1+ (with sizes between 10.9 and 17.9 cm). Additionally, the present study demonstrated that the relation between the stomach fullness and fish size was negative.

In brown trout, the peak feeding periods also corresponded with an increase in both condition factor and lipid content (Rikardsen et al. 2006). Hence, the negative relation between the stomach fullness and fish size and the positive relation between the condition factor and fish length found in this study could indicate that the feeding intensity is related with the fish condition, as other researchers have found in previous studies (e.g., Rikardsen et al. 2006). In this context, it is important to note that the high feeding intensity of YOY during the end of the summer could be related with the increases in fish condition and survival in the later autumn and winter.

Acknowledgements

This work has been partially supported by the project 10PXIB2111059PR of the Xunta de Galicia and the project MIGRANET of the Interreg IV B SUDOE (South-West Europe) Territorial Cooperation Programme (SOE2/P2/E288). Protocols used in this study conform to the ethical laws of the regional government (Xunta de Galicia). The authors are also grateful to two anonymous referees for their helpful comments.

Literature

- CEN 2003: Water quality – sampling of fish with electricity. European standard – EN 14011: 2003. *European Committee for Standardization, Brussels*.
- Cobo F., Sánchez-Hernández J., Vieira-Lanero R. & Servia M.J. 2013: Organic pollution induces domestication-like characteristics in feral populations of brown trout (*Salmo trutta*). *Hydrobiologia* 705: 119–134.
- Elliott J.M. 1994: Quantitative ecology and the brown trout. *Oxford University Press, Oxford*.
- Fochetti R., Argano R. & Tierno de Figueroa J.M. 2008: Feeding ecology of various age-classes of brown trout in River Nera, Central Italy. *Belg. J. Zool.* 138: 128–131.
- Herbinger C.M. & Friars G.W. 1991: Correlation between condition factor and total lipid content in Atlantic salmon, *Salmo salar* L., parr. *Aquacul. Fish. Manage.* 22: 527–529.
- Jensen H., Bøhn T., Amundsen P.-A. & Aspholm P. 2004: Feeding ecology of piscivorous brown trout (*Salmo trutta* L.) in a subarctic watercourse. *Ann. Zool. Fenn.* 41: 319–328.
- Kara C. & Alp A. 2005: Feeding habits and diet composition of brown trout (*Salmo trutta*) in the upper streams of river Ceyhan and river Euphrates in Turkey. *Turk. J. Vet. Anim. Sci.* 29: 417–428.
- Lagarigue T., Céréghino R., Lim P., Reyes-Marchant P., Chappaz R., Lavandier P. & Belaud A. 2002: Diel and seasonal variations in brown trout (*Salmo trutta*) feeding patterns and relationship with invertebrate drift under natural and hydropeaking conditions in a mountain stream. *Aquat. Living Resour.* 15: 129–137.
- Machado C.R., Garofalo M.A.R., Roselino J.E.S., Kettelhut I.C. & Migliorini R.H. 1988: Effects of starvation, refeeding, and insulin on energy-linked metabolic processes in catfish (*Rhamdia hilarii*) adapted to a carbohydrate-rich diet. *Gen. Comp. Endocrinol.* 71: 429–437.
- Martínez-Ansemil E. & Membiela P. 1992: The low mineralized and fast turnover watercourses of Galicia. *Limnetica* 8: 125–130.
- Nomura T., Myers K.W., Kondzela C.M., Murphy J.M., Honma H. & Carlson H.R. 2001: Variation in lipid content in the muscle of chum and pink salmon in the Gulf of Alaska in May 1999. *Bull. Nat. Salmon Resources Center* 4: 13–18.
- Rikardsen A.H., Amundsen P.-A., Knudsen R. & Sandring S. 2006: Seasonal marine feeding and body condition of sea trout *Salmo trutta* (L.) at its northern distribution area. *ICES J. Mar. Sci.* 63: 466–475.
- Sánchez-Hernández J. & Cobo F. 2012: Summer differences in behavioural feeding habits and use of feeding habitat among brown trout (Pisces) age classes in a temperate area. *Ital. J. Zool.* 79: 468–478.
- Sánchez-Hernández J., Servia M.J., Vieira-Lanero R. & Cobo F. 2013: Ontogenetic dietary shifts in a predatory freshwater fish species: the brown trout as an example of a dynamic fish species. In: Türker H. (ed.), *New advances and contributions to fish biology. InTech*: 271–298.
- Sánchez-Hernández J., Vieira-Lanero R., Servia M.J. & Cobo F. 2011: First feeding diet of young brown trout fry in a temperate area: disentangling constraints and food selection. *Hydrobiologia* 663: 109–119.
- Shustov Yu.A., Baryshev I.A. & Belyakova E.E. 2012: Specific features of the feeding of juvenile Atlantic salmon (*Salmo salar* L.) in the subarctic Varzuga River and its small tributaries (Kola Peninsula). *Inland Water Biol.* 5: 288–292.
- Van der Oost R., Beyer J. & Vermeulen N.P.E. 2003: Fish bioaccumulation and biomarkers in environmental risk assessment: a review. *Environ. Toxicol. Pharmacol.* 13: 57–149.