



New and noteworthy bryophyte records for the flora of Yemen Additions to the Bryophyte Flora of the Arabian Peninsula and Socotra 8.

Authors: Kürschner, Harald, and Sollman, Philip

Source: Willdenowia, 34(1) : 301-308

Published By: Botanic Garden and Botanical Museum Berlin (BGBM)

URL: <https://doi.org/10.3372/wi.34.34126>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

doi:10.3372/wi.34.34126 (available via <http://dx.doi.org/>)

HARALD KÜRSCHNER & PHILIP SOLLMAN

New and noteworthy bryophyte records for the flora of Yemen Additions to the Bryophyte Flora of the Arabian Peninsula and Socotra 8.

Abstract

Kürschner, H. & Sollman, P.: New and noteworthy bryophyte records for the flora of Yemen. Additions to the Bryophyte Flora of the Arabian Peninsula and Socotra 8. – Willdenowia 34: 301-308. – ISSN 0511-9618; © 2004 BGBM Berlin-Dahlem.

Based on recent bryophyte collections eight new records are added to the bryophyte flora of mainland Yemen, increasing the number of known species to 133 (one hornwort, 39 liverworts, 93 mosses). First records for the Arabian Peninsula among the new findings are *Archidium alternifolium* (Archidiaceae) and *Tuerckheimia svihlae* (Pottiaceae), which are briefly described and illustrated.

Introduction

Our knowledge of the bryophyte flora of mainland Yemen (for Socotra Island see Kürschner 2003a) is still very incomplete and despite various collecting activities, the country appears to be bryologically undercollected. As a first step towards a better understanding of this neglected group of plants and to act as a stimulus for future bryological research, the Bryophyte Flora of the Arabian Peninsula and Socotra (Kürschner 2000a) was published, listing 103 taxa (one hornwort, 31 liverworts, 71 mosses) for the Yemeni mainland.

Ongoing bryophyte collecting activities within the last three years in the frame of the Yemeni German BIOTA (BIODiversity monitoring Transect Analysis) Project (http://www.biota-africa.de/800/biota_east/structure_east.htm) has resulted in a number of new records (e.g., Kürschner & al. 2001, Kürschner 2003b-c, Kürschner & Ochya 2003), which increased the bryophyte flora of mainland Yemen to 125 species (one hornwort, 39 liverworts, 85 mosses). The continuous determination of the large collections led to the discovery of eight additional species so far unknown to mainland Yemen. New records of taxa are an important first step towards a flora of a country. In this respect, the following eight new records are a further contribution to reach this ambitious aim.

The voucher material is deposited at B, duplicates are in the herbarium H. Kürschner (Berlin) and, as far as available, in Leiden (L). Authors of botanical names are abbreviated according to Brummitt & Powell (1992).

New records for mainland Yemen

Archidiaceae

Archidium alternifolium (Hedw.) Mitt. – Fig. 1

Plants yellow to yellowish green, 3–10 mm high, in dense, glossy tufts or loose gregarious patches. *Stems* simple, later branched by fertile innovations, often capitate-foliate. *Leaves* erect-spreading, ovate-lanceolate to narrowly triangular-lanceolate, 0.7–2 mm long, acuminate to $\pm$ subulate, grading into the much longer perichaetial leaves; margins plane, entire to serrulate; costa percurrent to short-excurrent; laminal cells rhomboidal to linear rhomboidal, $45\text{--}70 \times 9\text{--}12\ \mu\text{m}$, becoming shorter above; basal cells rectangular, at margin quadrate to short-rectangular; perichaetial leaves ovate-lanceolate to narrowly lanceolate, long-acuminate to subulate, 2–3.5 mm long; paroicous. *Capsules* immersed, globose, terminal, 1–2 per stem, 0.5–0.8 mm in diameter, cleistocarpous. *Spores* very large, 125–250 μm in diameter, irregularly polyhedral, 8–30 per capsule.

AL-MAHRA: Fartak Mts, Jabal Karmoun, $15^{\circ}50'\text{N}$, $51^{\circ}59'\text{E}$, 780 m, on soil in isolated island-like patches of *Anogeissus dhofarica*, 6.10.2001, *H. Kürschner* 01-1383.

A northern, somewhat polymorphic species frequently growing on damp, often sandy soils at the margins of temporary pools, on ditch banks, pastures, among grasses and fields. Widely distributed in the northern hemisphere and known from W and Central Europe eastwards to Greece and W Russia, northwards to Faeroer and Iceland, also from the Azores and Canary Islands (Hill & al. 1992), SE United States (Florida, Louisiana north to Missouri, Ohio, N Carolina; Crum & Anderson 1981), N Africa (Algeria, Morocco, Tunisia, Ros & al. 1999) and S Asia (Turkey; Frey & Kürschner 1991). Former reports from China could not be confirmed by Gao & Crosby (1999) and all so determined specimens from Africa south of the Sahara should be referred to *A. ohioense* Schimp. (O'Shea 1999, Snider 1975).

Pottiaceae

Barbula consanguinea (Thwaites & Mitt.) A. Jaeger

AL-MAHRA: Hawf Mts, Uteq area near Al-Ayn, Jabal Chattan, $16^{\circ}38.9'\text{N}$, $52^{\circ}57.6'\text{E}$, 780 m, on soil beneath rocks in *Anogeissus dhofarica* woodland, 29.9.2001, *H. Kürschner* 01-1296; Shah'rut area near Damqawt, $16^{\circ}33.5'\text{N}$, $52^{\circ}46.2'\text{E}$, 700 m, on soil in *Anogeissus dhofarica* woodland, 2.10.2001, *H. Kürschner* 01-1333, 01-1345; N Fartak Mts, Jabal Karmoun, $15^{\circ}50'\text{N}$, $51^{\circ}59'\text{E}$, 780 m, on soil in isolated *Anogeissus dhofarica* patches, 6.10.2001, *H. Kürschner* 01-1379, 01-1382; *ibid.*, 950 m, 6.10.2001, *H. Kürschner* 01-1400; S Fartak Mts, northeast of Khadifud, $15^{\circ}39'\text{N}$, $52^{\circ}12'\text{E}$, 780 m, on soil in isolated *Anogeissus dhofarica* patches, 10.10.2001, *H. Kürschner* 01-1417.

Our concept of this very variable taxon follows Eddy (1990) and Sollman (2000), conflicting with Zander (1993), who listed *Barbula consanguinea* as a synonym of *B. javanica* Dozy & Molk. However, both taxa are clearly distinct by the laminal cells (mamillose and pellucid in *B. javanica*, pluripapillose and obscure in *B. consanguinea*; cf. Sollman 2000). *B. consanguinea* is considered as a pantropical species frequent and widespread on damp soils and rocks throughout tropical Asia, Africa and Central America. Previous Arabian records are from the neighbouring Oman (Dhofar; Kürschner 2000b).

Gymnostomiella vernicosa var. *tenerum* (Müll. Hal. ex Dusén) Arts

AL-MAHRA: Hawf Mts, Jabal Lusaka northeast of Faydami, west of Al-Fatk, 550 m, isolated *Anogeissus dhofarica* patches, on soil in rock crevices near a spring, 5.10.2001, *H. Kürschner* 01-1361.

A pantropical species reported from the Americas (Lesser Antilles), Africa [Cape Verde Islands, Central African Republic, Comoro Islands, Democratic Republic of Congo (Zaire), Kenya, Liberia, Réunion, Tanzania, Zambia; O'Shea 1999] and Asia (Burma, China, India, Indonesia, Nepal,




Fig. 1. *Archidium alternifolium* – 1: distribution (open circles: regional and country records); 2: habit; 3: leaves; 4: perichaetial leaves; 5: leaf apex; 6: mid-laminal cells; 7: basal laminal cells; 8: cross-section of costa. – All drawn from *Kürschner 01-1383* (herb. Kürschner).

Sri Lanka, Thailand; Arts 1998). Previous Arabian records are from the neighbouring Oman (Dhofar, Arts 1998).

***Hymenostylum crassinervium* Broth. & Dixon**

AL-MAHRA: Hawf Mts, Uteq area near Al-Ayn, Jabal Chattan, 16°38.9'N, 52°57.6'E, 780 m, on soil in *Anogeissus dhofarica* woodland, 29.9.2001, *H. Kürschner* 01-1282; *ibid.*, 700 m, 30.9.2001, *H. Kürschner* 01-1304, 01-1308; Shah'rut area near Damqawt, 16°33.5'N, 52°46.2'E, 700 m, on soil beneath rocks in *Anogeissus dhofarica* woodland, 2.10.2001, *H. Kürschner* 01-1335; *ibid.*, 700 m, 2.10.2001, *H. Kürschner* 01-1336; Jabal Lusaka, northeast of Faydami, west of Al-Fatk, 550 m, on soil beneath rocks in *Anogeissus dhofarica* remnants, 5.10.2001, *H. Kürschner* 01-1358; N Fartak Mts, Jabal Karmoun, 15°50'N, 51°59'E, 780 m, on soil beneath rocks in isolated *Anogeissus dhofarica* patches, 6.10.2001, *H. Kürschner* 01-1365, 01-1367; *ibid.*, 950 m, 6.10.2001, *H. Kürschner* 01-1402; S Fartak Mts, northeast of Khadifud, 15°39'N, 52°12'E, 950 m, on soil beneath rocks in isolated *Anogeissus dhofarica* patches, 10.10.2001, *H. Kürschner* 01-1409; *ibid.*, 780 m, on soil in isolated *Anogeissus dhofarica* patches, 10.10.2001, *H. Kürschner* 01-1419.

An afrotropical species, reported from the Democratic Republic of Congo (Zaire), Kenya, Somalia, Tanzania, Zambia and Zimbabwe (O'Shea 1999). Previous Arabian records are from the neighbouring Oman (Dhofar, Kürschner & al. 2001).

***Tuerckheimia svihlae* (E. B. Bartram) R. H. Zander – Fig. 2, 3**

Plants light to dark green, growing in short-tufts, to 1-1.5 cm high. *Leaves* incurved, weakly twisted when dry, widely spreading when moist, oblong to linear-lanceolate, 2-3.5 mm long; margins plane, entire; apex narrowly acute; costa short excurrent and ending in an apiculus; costa in cross-section circular, ventrally bulging, with 2 stereid bands, both strong, with 2-4 guide cells; upper laminal cells rounded-quadrate to hexagonal, mostly 10-14 µm width, with massive, compound papillae centred over the lumen; basal cells differentiated, thin-walled, often hyaline, rectangular, smooth. *Sporophytes* not seen in Yemen material.

Tuerckheimia is very similar to *Trichostomum* and also *Molendoa* in habit but differs from both by the massive papillae centred over the lumen of the rather large upper laminal cells (Fig. 3). In *Trichostomum*, they are smaller and covered by crowded, multiplex, flattened papillae (Zander 1993). For the differences from *Molendoa* see especially Saito (1972).

AL-MAHRA: S Fartak Mts, northeast of Khadifud, 15°39'N, 52°12'E, 500 m, on soil in rock fissures; 10.10.2001, *H. Kürschner* 01-1407; Hawf Mts, northeast of Al-Ayn, Uteq area, 16°38.9'N, 52°57.6'E, 780 m, on soil in *Anogeissus dhofarica* forest, 30.9.2001, *H. Kürschner* 01-1290.

A species known from the SE United States and Mexico [as *Tuerckheimia angustifolia* (K. Saito) R. H. Zander, nom. inval.], E Asia (Japan, Korea, Taiwan) and SE Asia (Burma, as *Oxystegus svihlae* (E. B. Bartram) Gangulee, Philippines]. According to O'Shea (1999) and Ros & al. (1999), *Tuerckheimia svihlae* has not been recorded from Africa. *Trichostomum abyssinicum* (Thér.) R. H. Zander obviously was erroneously recorded as a synonym of *Tuerckheimia svihlae* by Zander (1993: 344; Crosby & Magill 1997). We have not yet studied type material of *Trichostomum abyssinicum*, nor seen any material so-determined. However, according to the protologue of *Weissia abyssinica* Thér. (Thériot 1928), which is the basionym of *T. abyssinicum*, it is clearly different.

Bryaceae

***Bryum dichotomum* Hedw. (Syn.: *B. bicolor* Dicks.)**

LAHJ: Jabal Eraf, summit plateau, 13°06'N, 44°15'E, 1380-1400 m, on soil in *Juniperus procera* woodland, 10.3.2002, *H. Kürschner* 02-43.

A subcosmopolitan species, ubiquitous in northern and southern temperate zones. Previous Arabian records are from Saudi Arabia and Socotra (Kürschner 2000a).




Fig. 2. *Tuerckheimia svihlae* – 1: distribution (open circles: regional and country records); 2: habit; 3: leaves; 4: leaf apex; 5: mid-laminal cells; 6: basal laminal cells; 7: cross-section of leaf. – All drawn from Kürschner 01-1290 (herb. Kürschner).




Fig. 3. *Tuerckheimia svihlae* – 1: mid-laminal cells with massive papillae centred over the lumen; 2: leaf margin (right) with massive papillae and costa (left). – Scale bars = 10 µm; SEM photographs from Kürschner 01-1290.

***Bryum capillare* Hedw.**

AL-MAHRA: Hawf Mts, Uteq area near al-Ayn, Jabal Chattan, 16°38.9'N, 52°57.6'E, 780 m, on soil in *Anogeissus dhofarica* woodland, 1.10.2001, H. Kürschner 01-1319.

A subcosmopolitan species of northern origin. Previous Arabian records are from Saudi Arabia and Socotra (Kürschner 2000a).

Funariaceae

Entosthodon muhlenbergii (Turner) Fife

DHAMAR: Jabal Uthmar, near Mirsa'a, c. 5 km southwest of Uthmar, 14°28.7'N, 44°00.2'E, 1620-1720 m, in loamy pockets between rocks, 12.8.2002, *N. Kilian, P. Hein & Al-Naggan* YP 2615b.

A circum-Tethyan species, widely distributed in Europe, N Africa, North America and SW Asia. Previous Arabian records are from Saudi Arabia and the United Arab Emirates (Kürschner 2000a).

Acknowledgements

The field excursion of the first author was carried out within the frame of the Yemeni-German BIOTA Project funded by the German Federal Ministry of Education and Research (bmb+f). The authors are grateful to Dr M. El-Mashjary, Environmental Protection Authority (EPA), Head Office Sana'a, to Dr M. A. Hubaishan, Agricultural Research and Extension Authority (AREA), research station Mukalla, Dr N. Kilian (BGBM Berlin) and P. Hein (Berlin) for their assistance in the field and to H. Lünser (Berlin) for drawing the figures.

References

- Arts, T. 1998: A revision of the moss genus *Gymnostomiella* Fleisch. – *J. Bryol.* **20**: 411-427.
- Brummit, R. K. & Powell, C. E. 1992: Authors of plant names. – Kew.
- Crosby, M. R. & Magill, R. E. 1997: Index of mosses 1993-1995. – Monogr. Syst. Bot. Missouri Bot. Gard. **62**.
- Crum, H. A. & Anderson, L. E. 1981: Mosses of eastern North America **1**. – New York.
- Eddy, A. 1990: A handbook of Malesian mosses **2**. – London.
- Frey, W. & Kürschner, H. 1991: Conspectus bryophytorum orientalium et arabicorum. An annotated catalogue of the bryophytes of Southwest Asia. – Bryoph. Bibl. **39**.
- Gao, C. & Crosby, R. M. 1999: Moss flora of China **1**. – St Louis.
- Hill, M. O., Preston, C. D. & Smith, A. J. E. 1992: Atlas of the bryophytes of Britain and Ireland **2**. Mosses (except *Diplolepideae*). – Colchester (Essex).
- Kürschner, H. 2000a: Bryophyte flora of the Arabian Peninsula and Socotra. – Bryoph. Bibl. **55**.
- 2000b: *Claopodium prionophyllum* (C. Müll.) Broth., a remarkable tropical Asian species in the moss flora of the Arabian Peninsula and further additions for Oman, Saudi Arabia and Yemen. Studies in Arabian bryophytes 26. – Nova Hedwigia **70**: 127-134.
- 2003a: Nineteen new records to the bryophyte flora of Socotra Island. Additions to the bryophyte flora of the Arabian Peninsula and Socotra 5. – *Willdenowia* **33**: 445-458.
- 2003b: Further new bryophytes to Socotra and Yemen mainland, including *Papillaria crocea* (Hampe) A. Jaeger (*Meteoriaceae*, *Bryopsida*), a palaeotropical Asian species. Additions to the Bryophyte Flora of the Arabian Peninsula and Socotra 3. – *Nova Hedwigia* **76**: 257-267.
- 2003c: The Riccietum jovet-astii – argenteolimbatae ass. nov. of the Jabal Arays area, Yemen – life strategies of a remarkable xerotropical African bryophyte community. – *Nova Hedwigia* **76**: 399-413. [[CrossRef](#)]
- , Buck, W. R. & Sollman, P. 2001: Two tropical species new to the bryophyte flora of the Arabian Peninsula. Additions to the Bryophyte Flora of the Arabian Peninsula and Socotra 2. – *Nova Hedwigia* **73**: 253-259.
- & Ochyra, R. 2003: *Erpodium glaziovii* (*Erpodiaceae*, *Bryopsida*) and further novelties from the Arabian Peninsula. Additions to the bryophyte flora of the Arabian Peninsula and Socotra 4. – *Willdenowia* **33**: 205-210.
- O'Shea, B. J. 1999: Checklist of the mosses of sub-Saharan Africa (version 3, 11/99). – Trop. Bryol. Res. Rep. **1**.

- Ros, M. R., Cano, M. J. & Guerra, J. 1999: Bryophyte checklist of Northern Africa. – J. Bryol. **21**: 207-244.
- Saito, K. 1972: Taxonomic and phytogeographic studies of specimens previously reported as *Molendoa sendtneriana*. – J. Hattori Bot. Lab. **36**: 163-170.
- Snider, J. A. 1975: A revision of the genus *Archidium* (*Muscii*) – J. Hattori Bot. Lab. **39**: 105-201.
- Sollman, P. 2000: Studies on *Barbula consanguinea* (Thw. & Mitt.) Jaeg. sensu Eddy, a pan-tropical species. – Trop. Bryol. **19**: 17-23.
- Thériot, M. I. 1928: Musci novi africani. – Bull. Mus. Hist. Nat. Paris **34**: 115-118.
- Zander, R. H. 1993: Genera of the *Pottiaceae*: mosses of harsh environments. – Bull. Buffalo Soc. Nat. Sci. **32**.

Addresses of the authors:

PD Dr Harald Kürschner, Freie Universität Berlin, Institut für Biologie, Systematische Botanik und Pflanzengeographie, Altensteinstr. 6, D-14195 Berlin, Germany; e-mail: kuersch@zedat.fu-berlin.de

Philip Sollman, Notarisappel 2, NL-9076 LB St Annaparochie, The Netherlands.