



Spatiotemporal variation of ringed seal blubber cortisol levels in the Canadian Arctic

Authors: Ogloff, Wesley R., Anderson, Randi A., Yurkowski, David J., Debets, Cassandra D., Anderson, W. Gary, et al.

Source: Journal of Mammalogy, 103(5) : 1208-1219

Published By: American Society of Mammalogists

URL: <https://doi.org/10.1093/jmammal/gyac047>



Spatiotemporal variation of ringed seal blubber cortisol levels in the Canadian Arctic

WESLEY R. OGLOFF,^{1,*} RANDI A. ANDERSON,^{2,†} DAVID J. YURKOWSKI,^{1,3} CASSANDRA D. DEBETS,³ W. GARY ANDERSON,³ AND STEVEN H. FERGUSON^{1,3}

¹Freshwater Institute, Fisheries and Oceans Canada, 501 University Crescent, Winnipeg, MB R3T 2N6, Canada

²Office of the Minister of Natural Resources Canada, Ottawa, ON K1A 0E4, Canada

³Department of Biological Sciences, University of Manitoba, 66 Chancellors Circle, Winnipeg, MB R3T 2N2, Canada

*To whom correspondence should be addressed: wesley.ogloff@live.ca

†WRO and RAA contributed equally to this work and are co-first authors.

Climate change in the Arctic has widespread and complex effects on the health of animals and their populations. We used radioimmunoassay to measure blubber cortisol in ringed seals (*Pusa hispida*) sampled in Ulukhaktok, Inuvialuit Settlement Region, Northwest Territories, Canada (spring, 2002, 2004–2005, 2007–2012) and Arviat, Nunavut, Canada (autumn, 2003–2012) to examine chronic stress relative to biology (age, sex, length), body condition (blubber depth), and diet ($\delta^{13}\text{C}$, $\delta^{15}\text{N}$, and isotopic niche size). Ulukhaktok ringed seals had higher cortisol concentrations overall (0.46 ± 0.04 ng/g) than Arviat ringed seals (0.36 ± 0.03 ng/g), and these higher concentrations were associated with higher muscle $\delta^{15}\text{N}$ and lower blubber thickness. In contrast, blubber cortisol concentrations for Arviat ringed seals decreased with blubber depth and increased with age, though testing of age effects individually suggests that age-related patterns are weak. Annual mean cortisol concentration increased from 2003 to 2012 in Arviat ringed seals, but low sample sizes precluded analysis of annual patterns for Ulukhaktok ringed seals. The trend of increased cortisol over time in Arviat ringed seals suggests that they might be experiencing greater chronic stress over time, which could have implications for numerous population health metrics including reproduction and pup recruitment.

Key words: Arctic, Arviat, blubber, body condition, chronic stress, cortisol, *Pusa hispida*, ringed seal, stable isotopes, Ulukhaktok

Climate change in the Arctic has led to a reduction in the quantity and quality of sea ice, a key habitat component for numerous species ranging from algae- and ice-associated invertebrates to polar bears (*Ursus maritimus*; Laidre et al. 2015; Hamilton et al. 2017; Hop et al. 2020; Hwang et al. 2020). This reduction in sea ice has allowed for the expansion of human impacts, including increased shipping and industrial activities, exposing animals to an increased risk of contaminant exposure, ship strikes, and noise pollution (Ghosh and Rubly 2015; Hauser et al. 2018; McWhinnie et al. 2018). Higher air and sea-surface temperatures introduce physiological challenges that could lead to hyperthermia in extreme cases and, more commonly, facilitate the northward shift of temperate and subarctic species (Evans and Bjørge 2013; Pinsky et al. 2013; Pecl et al. 2017), including pathogens (Cohen et al. 2018). Competition for food and

habitat can arise as southern and marginal species become more common in Arctic environments, impacting the food web at multiple levels (Kortsch et al. 2015; Pecl et al. 2017; Yurkowski et al. 2018). Foraging may become more energetically expensive as habitat becomes more fragmented, and energetic return may be limited for predators that must switch to prey that is more limited in quantity, accessibility, or energetic quality. These and other climate-driven changes will likely present new and compounding challenges for ice-associated Arctic species, which can then lead to population- and ecosystem-wide consequences over time, such as reduced population abundances and impacts on biodiversity (Kovacs et al. 2011; Laidre et al. 2015).

In response to challenging or threatening conditions, animals might experience a stress response, with the release of cortisol and other glucocorticoids facilitating a shift in energy balance

© The Author(s) 2022. Published by Oxford University Press on behalf of the American Society of Mammalogists.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

to maintain survival or homeostasis, a process that can be maladaptive when stress is chronic (Sapolsky et al. 2000; Romero and Butler 2007; Busch and Hayward 2009). Among the main drivers of chronic stress in animals are changes to habitat and food availability, both of which are threats in the warming Arctic (Kovacs and Lydersen 2008; Doney et al. 2012; IPCC 2014) and have implications for the variation of individual- and population-level responses to environmental change. Glucocorticoid concentration may be used as an indicator of stress, ideally when experimentally validated or paired with other measures of condition (Romero and Butler 2007; MacDougall-Shackleton et al. 2019). Indicators of stress at the individual level can then be used to infer animal health (Atkinson et al. 2015), which, given adequate sample sizes, can indicate the general health of the population. Acute levels of cortisol can be measured in tissues with fast turnover times, such as blood (e.g., Ortiz et al. 2001; DeRango et al. 2019), making these tissues ideal candidates for examining the effects of animal handling during tagging or medical procedures or of the postweaning stress levels of seal pups (Ortiz et al. 2001; Champagne et al. 2012b; Atkinson et al. 2015; DeRango et al. 2019). In contrast, by measuring cortisol in metabolically inert or slow-turnover tissues (e.g., hair, whisker, blubber, and skin; Kershaw and Hall 2016; Bechshoft et al. 2020; Keogh et al. 2020a, 2020b), researchers can get an indication of chronic stress (Atkinson et al. 2015). Chronic stress can cause immunosuppression, altered fight-or-flight responses, lowered body condition, and reduced reproduction and/or offspring recruitment, which have implications for long-term fitness and population health (Gulland et al. 1999; Burek et al. 2008; Ferguson et al. 2017; Romero and Gormally 2019). Arctic marine mammals, like the ringed seal (*Pusa hispida*), are at particular risk of chronic stress, as their life histories are tightly linked with sea ice (McLaren 1958; Kovacs et al. 2011).

Ringed seals are abundant and circumpolar, making them an ideal candidate species to monitor ecosystem changes in the Arctic (Laidre et al. 2008; Hazen et al. 2019). In the summer open-water period, ringed seals feed intensively to build their blubber layer in preparation for the winter, when resources are more limited (Smith 1987; Freitas et al. 2008; Yurkowski et al. 2016b). In autumn, ringed seals continue to forage and can move long distances to overwintering areas (Harwood et al. 2012; Ogloff et al. 2021)—at this time, their condition is typically at its highest of the year (McLaren 1958; Young and Ferguson 2013). During the winter, foraging strategies shift from pelagic to benthic in some areas (Young and Ferguson 2013), and adult ringed seals establish territories and maintain breathing holes in the sea ice, whereas subadults inhabit pack ice areas or move into open water (McLaren 1958; Kelly et al. 2010; Crawford et al. 2012; Harwood et al. 2015). The final stages of gestation, as well as parturition and lactation, occur in the spring and have considerable energetic costs for females (Ryg et al. 1990; Hammill et al. 1991; Lydersen 1995). Following the nursing and mating period, ringed seals also undergo the energetically expensive process of molting, where they shed and regrow their fur (McLaren 1958). During this time, ringed seals are at their poorest condition of the year and have a markedly higher resting metabolic rate (McLaren 1958; Young and Ferguson 2013; Thometz et al. 2020). At southern latitudes, such as Hudson Bay, Canada, declines in ringed seal reproduction

and pup recruitment have been observed between 2003 and 2013 (Ferguson et al. 2017), though the mechanisms of these changes are unknown. The present gaps in our understanding of these changes highlight the importance of long-term health monitoring of ringed seals and other marine mammal populations. Some factors of importance for monitoring include morphometric and condition data, chemical tracers (e.g., stable isotopes, contaminants, trace metals), and biomarkers (e.g., hormones, fatty acids, highly branched isoprenoids).

Here, we examine indicators of chronic stress in ringed seals from two Arctic locations in northern Canada over a period of ~10 years (Arviat: 2003–2012; Ulukhaktok: 2002, 2004–2005, 2007–2012). We measured cortisol concentrations in blubber and tested for relationships with a number of biological and demographic factors. Blubber depth was included in analyses as it is a common proxy for body condition in marine mammals (Crawford et al. 2015; Ostertag et al. 2018). Stable isotopes ($\delta^{15}\text{N}$, $\delta^{13}\text{C}$, and isotopic niche) were included as indicators of long-term foraging, as diet and nutritional state are important indicators of animal health. For example, poor nutritional state has been shown to correlate with higher blubber cortisol concentrations in California sea lions (*Zalophus californianus*; Beaulieu-McCoy et al. 2017). Biological and demographic factors—age, length (as a metric of body size), and sex—were included in analyses due to the roles they play in glucocorticoid production and their influence on the pressures faced by individuals (Dantzer et al. 2014). For instance, northern elephant seal pups (*Mirounga angustirostris*) experience high stress, with associated high cortisol levels, during the postweaning fast (Ortiz et al. 2001); some adult female northern right whales experience higher cortisol levels during pregnancy than during intercalving intervals (Hunt et al. 2017); cortisol measurements of pregnant bottlenose dolphins (*Tursiops truncatus*) increase with time postconception (Steinman et al. 2016); adult and subadult male Weddell seals (*Leptonychotes weddelli*) experience high cortisol levels as a cost of territoriality and competition for mates (Bartsh et al. 1992); smaller, thinner belugas (*Delphinapterus leucas*) have been shown to have higher blubber cortisol concentrations than larger individuals (Loseto et al. 2017); blubber cortisol in Indo-Pacific humpback dolphin (*Sousa chinensis*) calves is positively correlated with body length (Guo et al. 2022); and higher hair cortisol concentrations in polar bears have been associated with smaller, lighter, poorer condition individuals (Macbeth et al. 2012). Ringed seals in the present study were sampled in Arviat in autumn, following the summer foraging period, and in Ulukhaktok in spring, when ringed seals are in their molting phase (McLaren 1958). Specifically, we set out to test (i) which factors relate to ringed seal blubber cortisol levels and (ii) whether there is spatial, seasonal, and long-term variation in blubber cortisol levels of ringed seals.

MATERIALS AND METHODS

Biological sampling.— Ringed seals were sampled during Inuit subsistence harvests in Arviat, Nunavut, and Ulukhaktok, Northwest Territories, from 2002 to 2012 (Fig. 1). In Ulukhaktok, located within the Inuvialuit Settlement




Fig. 1.—Map highlighting our study regions, Arviat, Nunavut (circle), and Ulukhaktok, Northwest Territories (diamond), within the Canadian Arctic. Ringed seals in the present study were sampled in Arviat in autumn (October and November; 2003–2012) and in Ulukhaktok in spring (May and June; 2002, 2004–2005, 2007–2012). The black square in the inset shows the position of the study area within North America.

Region, ringed seals were mainly harvested in the spring and summer (May and June; 2002, 2004–2005, 2007–2012), whereas the harvest in Arviat mainly occurred in autumn (October and November; 2003–2012). Ringed seal samples and data from Arviat were also used in [Ferguson et al. \(2017\)](#). Morphometric measurements were taken, and tissues were sampled in field by hunters. Standard length (straight line) was measured as the distance from the tip of the nose to the end of the tail ([Committee on Marine Mammals 1967](#)). Samples of blubber—including the hair and skin, through to the outer muscle layer, representing a cross section of all three blubber layers ([Strandberg et al. 2008](#))—and muscle were taken. Lower jaws were collected for age estimation using canine teeth, which were extracted and sectioned to allow counting of the dentin layers. This procedure was conducted in Ulukhaktok or at Matson's Lab in Manhattan, MT. Ringed seals aged ≥ 6 years were considered adults, 1–5 years were subadults, and < 1 year were pups ([Holst et al. 1999](#); [Krafft et al. 2006](#)). Tissue samples were stored at -20°C until they were shipped to the Freshwater Institute in Winnipeg, MB, where they were subsequently stored at -30°C until analysis.

Cortisol extraction and analysis.— We visually assessed blubber sample quality for all samples used in this study, including only those which showed no signs of degradation (e.g., yellowing of the tissue, odor; [Trana et al. 2015](#)). We subsampled even, cross-sectional pieces (innermost to outermost, all layers) of tissue from the blubber sample to account for any differences in cortisol concentrations between blubber layers, as they have been shown to differ metabolically and functionally ([Strandberg et al. 2008](#)). Samples were then freeze-dried at -50°C for ~ 48 h. Cortisol extraction and radioimmunoassay were conducted using the procedures outlined in [Trana et al. \(2015\)](#). We determined cortisol extraction efficiency by adding a known volume of tritium-labeled cortisol to a sample, processing along with the others, and measuring the level of radioactivity remaining. Blubber mean cortisol extraction efficiency for this study was $60 \pm 4\%$ ($n = 11$). Inter-assay variation was $20 \pm 5\%$ ($n = 23$), and intra-assay variation was $6 \pm 3\%$ ($n = 5$). Cortisol results are reported as ng/g of blubber tissue.

Stable isotope analysis.— We subsampled ~ 3 – 5 g of ringed seal muscle tissue from each muscle sample for stable isotope analysis. Subsamples were freeze-dried at -50°C for 48 h and

homogenized using a mortar and pestle. Lipids were extracted using a 2:1 chloroform:methanol solvent mixture, based on the established Bligh and Dyer (1959) method. Dried, homogenized muscle samples were placed into 10-ml centrifuge tubes, and 5 ml of solvent was added. We vortexed samples for ~1 min and allowed them to sit overnight in a 30°C water bath. Samples were then centrifuged for 5 min at 3,000 rpm, and the supernatant was poured off and discarded. We rinsed the remaining pellets with an additional 5 ml of chloroform:methanol, vortexed, and centrifuged again without the overnight soak. This process was repeated, and sample pellets were then dried overnight under a fume hood.

We rehomogenized lipid-extracted muscle tissue and weighed 400–600 µg of tissue into tin capsules. Ratios of $^{13}\text{C}:^{12}\text{C}$ and $^{15}\text{N}:^{14}\text{N}$ were measured using a Thermo Finnigan Delta^{plus} mass spectrometer equipped with an elemental analyzer at the Great Lakes Institute for Environmental Research, University of Windsor. Ratios of $^{13}\text{C}:^{12}\text{C}$ and $^{15}\text{N}:^{14}\text{N}$ were measured in samples and in Pee Dee Belemnite and atmospheric nitrogen standards, respectively. Relative differences of isotopic ratios (δ), measured in ‰, were determined using the following formula: $\delta^xE = \left(\frac{R_{\text{sample}} - R_{\text{standard}}}{R_{\text{standard}}} \right) \times 1,000$, where xE refers to ^{13}C or ^{15}N , and R refers to the ratio of heavy to light isotopes of element E in either the sample or the standard.

Body condition—As a proxy for body condition, we measured the depth of the blubber (to the nearest 10th of a cm) on the ventral side of the seal at the sternum using a ruler.

Statistical analysis.—We grouped seals by sampling location for analyses. All groups were checked for normality and homoscedasticity prior to statistical analysis using Shapiro–Wilk tests of normality and Levene’s tests of homogeneity of variance. Cortisol data were highly non-normal, so a log-transformation was applied for subsequent analyses. Outliers in the cortisol data were removed by calculating Z-scores and removing values with Z-scores $\geq |3|$, resulting in omission of three values. We used the SIBER package v2.1.4 (Jackson et al. 2011) in R version 4.1.0 (R Core Team 2019) to calculate isotopic niche sizes (standard ellipse area corrected for small sample sizes; SEA_C) by location, year, and age-class. The standard ellipse area represents a standard deviation around the bivariate mean and encompasses ~40% of data, considered representative of the core population (Jackson et al. 2011). Yearly SEA_C was also non-normal and was log-transformed prior to statistical

tests. We used tests of difference to examine age-class (categorical) effects; specifically, we used ANOVA, Kruskal–Wallis rank-sum tests, t -tests, or Wilcoxon rank-sum tests—depending on sample sizes, normality, and homoscedasticity—to test for differences in cortisol, condition, and stable isotopes between ringed seal age groups.

Generalized linear mixed models (GLMMs) were run using the lmer function in the lme4 package v1.1-23 (Bates et al. 2015) in R version 4.1.0 (R Core Team 2019) to test for predictive factors of ringed seal blubber cortisol, with model runs on both an annual timescale (to allow for inclusion of annual isotopic niche size and to examine patterns over time) and irrespective of year. When not grouped by year, year was included in the models as a random effect, and age was included as a continuous variable. Models were assessed using Akaike’s information criteria. The inclusion of interaction terms was explored, but, ultimately, stepwise removal of variables during data exploration resulted in all interaction terms being removed early on, with the same end points as when interactions were excluded. For this reason, and to keep analyses simple and concise, interaction terms are not presented here. We assessed multicollinearity in our full models using variance inflation factors (VIFs) calculated via the check_collinearity function in the R package performance v0.8.0 (Lüdecke et al. 2021). For models using data throughout the sampling period, all VIFs were low (≤ 1.7 for Arviat and < 3.2 for Ulukhaktok). For models assessing annual mean values across variables, VIFs for some variables were moderate (8.4 and 6.6 for age-class and length, respectively); however, upon removal of age-class in subsequent model runs, all VIFs were ≤ 2.7 . Model performance for each model was assessed using conditional R -squared values (R^2_C —indicating the variance explained by the entire model, including both fixed and random effects), calculated using the r.squaredGLMM function in the R package MuMIn v1.43.17 (Bartoń 2020). Results are reported as mean \pm standard error (SE) throughout.

RESULTS

A total of 449 ringed seals were sampled from 2002 to 2012 (Table 1). For Arviat, there were no samples collected in 2002, and samples from Ulukhaktok were absent in 2003 and 2006. Overall, mean cortisol measurements were low (0.40 ± 0.03 ng/g; 0.01–2.66 ng/g), with the highest mean

Table 1.—Sample sizes and means (SE) of blubber cortisol, blubber depth, muscle $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$, and yearly isotopic niche breadths for ringed seals sampled in Arviat, Nunavut (autumn; 2003–2012), and Ulukhaktok, Northwest Territories (spring; 2002, 2004–2005, 2007–2012).

Age-class	Blubber cortisol		Blubber depth at sternum		Stable isotopes			Annual isotopic niches		
	<i>n</i>	(ng/g)	<i>n</i>	(cm)	<i>n</i>	$\delta^{13}\text{C}$ (‰)	$\delta^{15}\text{N}$ (‰)	C:N	<i>n</i>	SEA_C (‰ ²)
Arviat										
Adult	66	0.38 (0.05)	101	5.7 (0.1)	102	−20.42 (0.07)	+16.00 (0.08)	3.47 (0.02)	9	1.42 (0.22)
Subadult	40	0.32 (0.08)	76	4.7 (0.2)	77	−20.60 (0.09)	+15.70 (0.10)	3.48 (0.03)	9	1.65 (0.36)
Pup	48	0.35 (0.06)	86	3.9 (0.1)	87	−20.59 (0.07)	+14.87 (0.10)	3.46 (0.03)	9	1.20 (0.32)
Ulukhaktok										
Adult	101	0.45 (0.04)	141	2.6 (0.1)	126	−20.58 (0.04)	+16.72 (0.05)	3.37 (0.01)	7	0.54 (0.05)
Subadult	11	0.52 (0.12)	14	1.9 (0.2)	11	−20.81 (0.18)	+15.72 (0.25)	3.38 (0.01)	—	—

concentration (0.52 ± 0.12 ng/g; 0.11–1.28 ng/g) found in subadults from Ulukhaktok (Fig. 2; Table 1). In contrast, subadults from Arviat had the lowest mean concentration of cortisol (0.32 ± 0.08 ng/g; 0.02–2.66 ng/g; Fig. 2; Table 1). Log-cortisol did not differ significantly by age-class in Arviat ringed seals (ANOVA: $F_{2,151} = 0.98$, $P = 0.38$; Table 1), or between adults and subadults from Ulukhaktok (Welch two-sample t -test: $t_{13,14} = -0.95$, $P = 0.36$; Table 1). Blubber depth differed significantly among age-classes for Arviat ringed seals (ANOVA: $F_{2,260} = 47.05$, $P < 0.0001$), with post hoc tests revealing a significant increase in blubber depth with age (Tukey's multiple comparison test: $P < 0.001$; Fig. 2; Table 1). Similarly, Ulukhaktok adults had significantly thicker blubber than subadults (Welch two-sample t -test: $t_{16,13} = 4.33$, $P < 0.001$; Fig. 2; Table 1). Both $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ differed significantly between age groups for Arviat ringed seals (Kruskal–Wallis rank-sum tests: $\chi^2_2 = 9.88$, $P < 0.01$ and $\chi^2_2 = 64.65$, $P < 0.0001$, respectively). The $\delta^{13}\text{C}$ values in Arviat adults were significantly higher than both subadults (Dunn's multiple comparison test: $P = 0.03$) and pups ($P = 0.01$; Table 1). For $\delta^{15}\text{N}$, all pairwise age-class comparisons differed significantly (Dunn's multiple comparison test: $P < 0.05$; Table 1), with increases in $\delta^{15}\text{N}$ with age. For Ulukhaktok

ringed seals, $\delta^{13}\text{C}$ did not differ significantly between adults and subadults (Welch two-sample t -test: $t_{10,84} = 1.22$, $P = 0.25$), but $\delta^{15}\text{N}$ was significantly greater in adults than subadults (Wilcoxon rank-sum test: $W = 1186.5$, $P < 0.0001$; Table 1).

Isotopic niches were largest for Arviat subadults and smallest for Ulukhaktok adults (Fig. 3; Table 1). There was no significant difference in mean annual SEA_C among Arviat ringed seal age-classes (Kruskal–Wallis rank-sum test: $\chi^2_2 = 1.53$, $P = 0.47$), though Arviat adults had a significantly higher mean SEA_C than Ulukhaktok adults (Wilcoxon rank-sum test: $W = 58$, $P < 0.01$; Table 1). We were unable to calculate and compare mean annual niches for Ulukhaktok subadults due to limited samples.

GLMMs revealed different predictive factors explaining log-cortisol concentration for both locations. The most-supported model predicting cortisol concentrations in Arviat ringed seals included age (continuous) and blubber depth, whereas blubber depth and $\delta^{15}\text{N}$ best predicted cortisol concentration in Ulukhaktok ringed seals (Tables 2 and 3). For both locations, the relationship between cortisol concentration and blubber depth was negative, indicating greater cortisol levels for ringed seals with lower body condition (Table 3). For Arviat ringed seals, cortisol concentration increased with seal




Fig. 2.—Boxplots indicating (A) blubber cortisol concentrations (ng/g) and (B) blubber depth (cm), measured at the sternum, of ringed seals sampled in Arviat, Nunavut (autumn; 2003–2012), and Ulukhaktok, Northwest Territories (spring; 2002, 2004–2005, 2007–2012). Solid horizontal lines on each boxplot indicate the median, and the dotted line represents the mean. Whiskers denote the highest and lowest points that fall within $1.5 \times$ the interquartile range; points outside of this are outliers. The lower and upper limits of the boxes represent the first and third quartiles.




Fig. 3.—Biplots of $\delta^{13}\text{C}$, $\delta^{15}\text{N}$, and isotopic niches of ringed seal muscle sampled in (A) Arviat, Nunavut (autumn; 2003–2012), and (B) Ulukhaktok, Northwest Territories (spring; 2002, 2004–2005, 2007–2012). Standard ellipse areas, corrected for small sample sizes (SEA_C) are denoted for each age group.

age, whereas, for Ulukhaktok ringed seals, cortisol concentration increased with $\delta^{15}\text{N}$ (Table 3). Mean annual cortisol levels for Arviat ringed seals increased significantly with year, with a 3.4-fold increase in cortisol over the study period, from 0.14 ng/g in 2003 to 0.47 ng/g in 2012 (Table 4; best-supported model: year slope = 0.1363 ± 0.0325 , $t = 4.192$, $P < 0.01$, null deviance = 7.5774 on 22 d.f., residual deviance = 4.1251 on 21 d.f.). Limited data prevented analysis of annual trends for Ulukhaktok ringed seals.

DISCUSSION

Given the rapid climate-driven changes in the Arctic (IPCC 2014; Box et al. 2019), it is important to monitor the health of Arctic marine species to better understand changes to

population trends. Animal health can be monitored using a variety of morphometric, chemical, and behavioral (e.g., observations of behavioral changes—lethargy, aggression, abandonment of young, altered movements/dives, etc.; Biuw et al. 2003; Huntington et al. 2017; Ostertag et al. 2018) methods. In the present study, we measured the levels of cortisol, as an indicator of chronic stress, found within the blubber of ringed seals sampled during two different seasons and in two different locations, representing ringed seals during their presumptive peak condition of the year (i.e., Arviat ringed seals) as well as at their lowest condition of the year (i.e., Ulukhaktok ringed seals; McLaren 1958; Young and Ferguson 2013). We found significant differences in blubber cortisol concentrations and body condition among locations, seasons, and age-classes. In addition, we found that body condition (blubber depth), diet ($\delta^{13}\text{C}$,

Table 2.—Summary of general linear mixed models testing the effects of measured variables on blubber cortisol in ringed seals sampled in Arviat, Nunavut (autumn; 2003–2012), and Ulukhaktok, Northwest Territories (spring; 2002, 2004–2005, 2007–2012). Included are the ranked models, regression coefficients (intercept and factors), number of estimated parameters (d.f.), log-likelihood (logLik), Akaike's information criterion corrected for small sample sizes (AIC_c), ΔAIC_c , AIC_c weights (Weight), and conditional R -squared (R^2_c) for each model. The “+” in the Sex column indicates its inclusion in the model.

Model	Intercept	Age (years)	$\delta^{13}C$ (‰)	$\delta^{15}N$ (‰)	Standard length (cm)	Sex	Blubber depth (cm)	d.f.	logLik	AIC_c	ΔAIC_c	Weight	R^2_c
Arviat													
A_5	-1.08	0.02309					-0.1404	5	-223.461	457.3	0	0.988	0.07
A_4	-0.4482	0.02984			-0.007013		-0.124	6	-226.97	466.5	9.19	0.01	0.07
A_3	-1.394	0.03	-0.04502		-0.006804		-0.125	7	-228.046	470.9	13.54	0.001	0.06
A_2	-1.478	0.02818	-0.04984		-0.006356	+	-0.1117	9	-226.864	473	15.67	0	0.08
A_1	-1.363	0.02819	-0.04834	-0.006948	-0.006198	+	-0.1114	10	-228.043	477.7	20.32	0	0.08
Ulukhaktok													
U_5	-5.474			0.3672			-0.6808	5	-80.968	172.8	0	0.98	0.41
U_4	-6.587			0.5352	-0.01402		-0.6794	6	-83.696	180.6	7.83	0.02	0.47
U_3	-6.056	0.01137		0.5242	-0.01859		-0.6699	7	-86.731	189.2	16.34	0	0.47
U_2	-6.264	0.01107		0.5326	-0.01771	+	-0.6688	8	-87.394	193	20.19	0	0.47
U_1	-6.139	0.01106	0.005306	0.5324	-0.01781	+	-0.6689	9	-87.818	196.4	23.63	0	0.47

Table 3.—Best-supported general linear mixed model output examining effects on the log-cortisol of ringed seals sampled in Arviat, Nunavut (autumn; 2003–2012), and Ulukhaktok, Northwest Territories (spring; 2002, 2004–2005, 2007–2012). Note that values under the *SE* column header for random effects represent standard deviation rather than standard error (*SE*), denoted by *.

Parameter	Estimate	<i>SE</i>	d.f.	<i>t</i> -value	<i>P</i>
Arviat					
(Intercept)	-1.08049	0.31161	62.23	-3.47	0.000959
Blubber depth (cm)	-0.14038	0.06628	141.7	-2.12	0.035917
Age (years)	0.02309	0.01079	143.1	2.14	0.034045
Random effects					
Year variance	0.02806	0.1675*			
Ulukhaktok					
(Intercept)	-5.4742	1.9658	70.67	-2.79	0.00687
Blubber depth (cm)	-0.6808	0.1292	70.99	-5.27	0.0000014
$\delta^{15}N$ (‰)	0.3672	0.1208	70.57	3.04	0.00332
Random effects					
Year variance	0.0921	0.3035*			

Table 4.—Summary of general linear mixed models testing the effects of measured variables on yearly mean blubber cortisol in ringed seals sampled in Arviat, Nunavut (autumn; 2003–2012). Included are the ranked models, regression coefficients (intercept and factors), number of estimated parameters (d.f.), log-likelihood (logLik), Akaike's information criterion corrected for small sample sizes (AIC_c), ΔAIC_c , AIC_c weights (Weight), and conditional R -squared (R^2_c) for each model. The “+” in the Age-class column indicates its inclusion in the model.

Model	Intercept	Age-class	Standard length (cm)	SEA_c (‰ 2)	Blubber depth (cm)	Year	d.f.	logLik	AIC_c	ΔAIC_c	Weight	R^2_c
A.Y_5	-275					0.1363	3	-12.874	33	0	0.555	0.44
A.Y_4	-273.6				0.1241	0.1353	4	-12.155	34.5	1.52	0.259	0.46
A.Y_3	-254.8		-0.01476		0.3052	0.1263	5	-11.081	35.7	2.68	0.145	0.50
A.Y_2	-244.9		-0.01328	-0.09791	0.307	0.1213	6	-10.529	38.3	5.3	0.039	0.51
A.Y_1	-233.8	+	-0.01897	-0.09287	0.28	0.1162	8	-10.425	47.1	14.13	0	0.48

$\delta^{15}N$, and isotopic niches), and morphometric and demographic factors (age, size, and sex) explained our observed patterns in cortisol concentrations.

Ringed seals sampled in Ulukhaktok during spring had higher absolute blubber cortisol concentrations than those sampled in Arviat in autumn. This can likely be explained by the timing of sampling, as Ulukhaktok ringed seals were hunted following the spring fasting, molting, and pupping period, when their body condition is at or near its lowest (McLaren 1958; Young and Ferguson 2013). A similar pattern has also been observed in harbor seals (*Phoca vitulina*) sampled during the molt (Kershaw and Hall 2016) and in western Hudson Bay polar bears sampled following the summer fast (Mislan et al. 2016). In contrast, ringed seals

in Arviat were sampled following intense foraging over the summer open-water period, when their body condition is at or near its peak (McLaren 1958; Young and Ferguson 2013). Blubber thickness was generally higher for Arviat ringed seals collected in autumn than Ulukhaktok ringed seals collected in spring, which also agrees with seasonal patterns in condition. Cortisol concentrations did not differ significantly between ringed seal age-classes in Arviat or Ulukhaktok.

Blubber thickness increased significantly with age for both sampling locations/seasons. This pattern suggests that ringed seals accumulate a thicker layer of blubber as they grow, as has been observed in phocids generally (Chabot et al. 1996; Chabot and Stenson 2002; Strandberg et al. 2008). Significant

increases in $\delta^{15}\text{N}$ with age for both sample locations could also be related to body growth, with older/larger individuals feeding at higher trophic levels than younger individuals (Ben-David and Flaherty 2012; Yurkowski et al. 2016b). The greater diving capacity of larger seals could also allow them to access larger, deeper, and higher trophic level prey sources with higher $\delta^{15}\text{N}$ (Lydersen et al. 1992; Wathne et al. 2000; Hobson et al. 2002). In contrast, higher $\delta^{13}\text{C}$ values in Arviat adults than subadults and pups might suggest that adult ringed seals feed more benthically, nearshore, or on more ice-associated prey than their younger counterparts. This latter idea is supported by the tendency for adults to outcompete subadults for primary feeding and mating habitats nearshore in the fall (McLaren 1958; Krafft et al. 2007; Crawford et al. 2012), though this pattern in $\delta^{13}\text{C}$ is not evident for Ulukhaktok ringed seals, possibly due to the relative lack of recent feeding, seasonal differences, or the unbalanced sample sizes between age-classes. The relatively larger isotopic niche of Arviat adults than Ulukhaktok adults could also be reflective of differences in recent feeding, as well as differences in prey availability and degree of specialization (Yurkowski et al. 2016a); however, spatial or seasonal differences in isotopic baselines (Arviat phytoplankton $\delta^{13}\text{C}$: -29.8% , $\delta^{15}\text{N}$: $+8.3\%$, Graham et al. 2010; Ulukhaktok phytoplankton $\delta^{13}\text{C}$: -27.6% , $\delta^{15}\text{N}$: $+5.4\%$, Roy et al. 2015) and food chain length between the two locations could also affect isotopic niche size and should be considered when interpreting our observed patterns (Vander Zanden and Fetzer 2007; Ward and McCann 2017).

For Ulukhaktok ringed seals sampled in spring, log-cortisol was best predicted by blubber depth and $\delta^{15}\text{N}$. The positive relationship between $\delta^{15}\text{N}$ and cortisol suggests a possible fasting signal, which is further supported by the negative relationship to blubber depth, indicating poorer body condition (Hobson et al. 1993; Kurle and Worthy 2001; Hertz et al. 2015; Lübcker et al. 2020). Thinner ringed seals had higher $\delta^{15}\text{N}$ and higher cortisol concentrations, suggesting that, as body condition decreased, ringed seals metabolized more of their own tissue, becoming enriched in ^{15}N and possibly experiencing greater stress (Hobson et al. 1993; Hertz et al. 2015; Doi et al. 2017). Although a similar relationship has been found in numerous species of marine mammal (e.g., polar bears, Polischuk et al. 2001; northern fur seals *Callorhinus ursinus*, Kurle and Worthy 2001; southern elephant seals *Mirounga leonina*, Lübcker et al. 2020), other studies have shown more uncertain and variable patterns (Das et al. 2004; Gómez-Campos et al. 2011; Aguilar et al. 2014; Spurlin et al. 2019), and it has been suggested that $\delta^{15}\text{N}$ is not a reliable predictor of fasting in some marine mammals, as many factors—such as gestation, lactation, and differing isoscapes between feeding locations—can confound the relationships (Gómez-Campos et al. 2011; Aguilar et al. 2014). However, given the supporting blubber measurements and the time of year Ulukhaktok ringed seals were collected in this study, a fasting signal is most likely, as fasting during the breeding season is common in some marine mammal species (Mesnick and Ralls 2009). Many female pinnipeds reduce or eliminate time

spent foraging during this period, instead mobilizing stored energy to support lactation (Costa 1993; Champagne et al. 2006), whereas males may fast while competing for mates and defending territory in order to maximize reproductive success (Galimberti et al. 2007; Champagne et al. 2012a).

For Arviat ringed seals sampled in autumn, cortisol was best predicted by blubber depth and age, with cortisol increasing with age and decreasing with blubber depth; however, when testing for differences in cortisol levels among Arviat ringed seal age-classes, no significant differences were found. Given the low R^2 values for these models, it is possible that the effects of age on blubber cortisol are only weak, with older individuals tending to have slightly higher cortisol concentrations in their blubber compared to their younger, more naïve counterparts. As these seals are generally in their best body condition of the year (McLaren 1958; Young and Ferguson 2013), it is uncertain whether this effect is biologically significant, and additional factors not measured here (e.g., contaminants, movement patterns, food web metrics, etc.) are likely involved.

Annual mean cortisol levels of Arviat ringed seals significantly increased over the ~10-year study period, indicating that these ringed seals are possibly experiencing greater chronic stress over time. Climate change in Hudson Bay has resulted in warmer waters (Galbraith and Larouche 2011; Brand et al. 2014), a longer open-water season (Kowal et al. 2017; Andrews et al. 2018), and altered prey assemblages, evidenced by changes in the diet of predators over time (Gaston et al. 2003; Chambellant et al. 2013; Gaston and Elliott 2014). Reduced snow and ice quality has also been documented in some Arctic locations (Webster et al. 2014), impacting ringed seal pup survival (Iacozza and Ferguson 2014). These environmental changes have the potential to lead to reduced fitness of ringed seals, and, consequently, population declines. Warmer waters might challenge the physiological capabilities of ringed seals and/or their prey, leading to altered energy expenditure and intake. A longer open-water season can impact several phases of the ringed seal life cycle; for example, earlier breakup can impact ringed seal ecology by reducing the duration and spatial extent of sea-ice platforms for molting, pupping, and nursing, potentially leading to increased pup mortality. Changes to prey assemblages could lead to increased competition (Ogloff et al. 2019) for limited prey and alter dietary composition, which can have negative implications if new prey are of lower energetic quality and digestibility. Though the mechanisms are not fully understood, previous studies have already suggested linkages between climate change and reductions in ringed seal ovulation rates, pregnancy rates, and pup recruitment, as well as increased levels of stress indicators and observations of sick seals, all of which appear to have contributed to declines in abundance (Ferguson et al. 2005, 2017; Stirling 2005).

Blubber cortisol measurements in this study were low (range: 0.01–2.66 ng/g) compared to other published blubber cortisol measurements from pinnipeds. For example, cortisol concentrations in the blubber of harbor seals in Scotland ranged from 37.8 ng/g throughout the non-molting period to

1,553.6 ng/g during the molt (Kershaw and Hall 2016), and mean blubber cortisol concentrations measured in California sea lions ranged from 8.1 ± 2.1 ng/g in bycaught animals to 256.3 ± 115.7 and 235.8 ± 68.5 ng/g for dead- and live-stranded animals, respectively (Beaulieu-McCoy et al. 2017). The low cortisol values measured in the present study could be related to biological factors or to measurement error. Namely, the 60% extraction efficiency could explain low cortisol values, but this extraction efficiency is within the range of values regularly published (e.g., 30% from beluga blubber, Loseto et al. 2017; 63% from harbor seal blubber, Kershaw and Hall 2016; 77% from beluga blubber, Trana et al. 2016; 73% from humpback whale *Megaptera novaeangliae* blubber, Beaulieu-McCoy et al. 2017). Alternately, low cortisol measurements could suggest that there is little cortisol to be measured in blubber relative to other tissues. Indeed, it has been shown that blubber actively metabolizes steroid hormones in some other marine mammals (Galligan et al. 2018); however, Champagne et al. (2017) found that blubber cortisol levels can be a reliable indicator of circulating cortisol levels in bottlenose dolphins. An additional possible explanation is that, because ringed seals are well-adapted to highly variable and extreme conditions, they are resilient to stressors and are well-equipped to handle low body condition in the spring (Ferguson and Higdon 2006). For instance, they regularly perform extensive long-distance movements (Teilmann et al. 1999; Harwood et al. 2012; Hamilton et al. 2015; Ogloff et al. 2021) and energetically expensive molting and fasting phases, where they can lose up to ~60% of their body fat (McLaren 1958; Ryg et al. 1990; Young and Ferguson 2013). As such, low levels of stress during challenging conditions could be an evolved trait of ringed seals and other animals in highly variable environments, allowing them to remain sensitive and responsive to additional stressors, such as predation (Champagne et al. 2015). Similarly, low cortisol measurements have been observed in other Arctic marine mammals (Macbeth et al. 2012; Trana et al. 2016; Watt et al. 2016), lending support to this idea.

Increases in measured cortisol over time are a noteworthy trend given that the climate will likely continue to warm (IPCC 2014). Although the data here are not extensive enough to link increased cortisol levels in Arviat ringed seals with warmer conditions, further research and attention is warranted. Given that cortisol measurements were overall low, it is possible that this population of ringed seals is not presently experiencing harmful levels of stress; however, the increasing extent and number of pressures that will co-occur with warmer conditions might exacerbate acute and chronic stress levels. On an individual level, this might lead to inhibited or reduced reproduction, reduced growth, and reduced immune function (Sapolsky et al. 2000; Tilbrook et al. 2000; Sheriff et al. 2010; Dantzer et al. 2014), though these effects are not always consistent in animals (Crespi et al. 2013).

In addition, we acknowledge that cortisol measurements are not synonymous with stress and that the relationship is much more complex (Busch and Hayward 2009;

MacDougall-Shackleton et al. 2019). There are many markers other than cortisol that are implicated in the stress response, including aldosterone, thyroid hormones, progesterone, and corticosteroid-binding globulin (Atkinson et al. 2015; Beaulieu-McCoy et al. 2017). Furthermore, cortisol will influence gene expression associated with numerous cellular pathways (MacDougall-Shackleton et al. 2019), and, in many animals, there are daily and seasonal fluctuations of circulating levels of cortisol unrelated to a specific response to a stressor. For instance, during winter hibernation, cortisol concentrations in black bears increase to assist with utilization of fat stores (Harlow et al. 1990). It is possible that, for ringed seals, seasonal differences in cortisol represent a biochemical adaptation to food scarcity, aiding in the mobilization of stored energy, rather than a stress response. The higher cortisol concentrations in Ulukhaktok seals support this idea, though spatial differences and other factors likely also play a role, and further investigation will be needed to draw conclusions on this matter.

We believe that pairing indicators of body condition with cortisol addresses some of the concerns with linking cortisol to stress; however, future studies incorporating other hormones and markers involved in the stress response would be beneficial and would further support our conclusions. Additional study might also address limitations with relation to sample sizes for Ulukhaktok subadults and pups as well as seasonality of sampling. The different timing of sampling precluded direct comparison among sites and seasons, but continued sampling might allow for comparisons across latitudes, seasons, and age-classes.

ACKNOWLEDGMENTS

We gratefully acknowledge the contributions of the Hunters and Trappers Associations of Ulukhaktok, Northwest Territories, and Arviat, Nunavut, as well as their hunters for their work collecting seal samples, especially Frank Nutarasugnik and John Alikamak. Thanks also to Blair Dunn and Brent Young for their coordination of community sampling, and to members of the Great Lakes Institute for Environmental Research for stable isotope analysis.

CONFLICT OF INTEREST

The authors declare no conflicts of interest.

FUNDING

Funding for this research came from the Fisheries and Oceans Canada (SHF), ArcticNet (SHF), the Natural Sciences and Engineering Research Council of Canada Discovery Grant (WGA-2020-05328), and the Faculty of Graduate Studies and Biological Sciences Department at the University of Manitoba (RAA).

LITERATURE CITED

- Aguilar A., Giménez J., Gómez-Campos E., Cardona L., Borrell A. 2014. $\delta^{15}\text{N}$ value does not reflect fasting in mysticetes. *PLoS ONE* 9:e92288.
- Andrews J., Babb D., Barber D.G. 2018. Climate change and sea ice: shipping in Hudson Bay, Hudson Strait, and Foxe Basin (1980–2016). *Elementa: Science of the Anthropocene* 6:1–23.
- Atkinson S., Crocker D., Houser D., Mashburn K. 2015. Stress physiology in marine mammals: how well do they fit the terrestrial model? *Journal of Comparative Physiology, B. Biochemical, Systemic, and Environmental Physiology* 185:463–486.
- Bartoń K. 2020. MuMIn: multi-model inference. R package version 1.43.17. <https://cran.r-project.org/package=MuMIn>. 14-04-2020.
- Bartsh S.S., Johnston S.D., Siniff D.B. 1992. Territorial behavior and breeding frequency of male Weddell seals (*Leptonychotes weddelli*) in relation to age, size, and concentrations of serum testosterone and cortisol. *Canadian Journal of Zoology* 70:680–692.
- Bates D., Mächler M., Bolker B.M., Walker S.C. 2015. Fitting linear mixed-effects models using lme4. *Journal of Statistical Software* 67:1–48.
- Beaulieu-McCoy N.E., Sherman K.K., Trego M.L., Crocker D.E., Kellar N.M. 2017. Initial validation of blubber cortisol and progesterone as indicators of stress response and maturity in an otariid; the California sea lion (*Zalophus californianus*). *General and Comparative Endocrinology* 252:1–11.
- Bechshøft T.O., Wright A.J., Styrihave B., Houser D. 2020. Measuring and validating concentrations of steroid hormones in the skin of bottlenose dolphins (*Tursiops truncatus*). *Conservation Physiology* 8:coaa032.
- Ben-David M., Flaherty E.A. 2012. Stable isotopes in mammalian research: a beginner's guide. *Journal of Mammalogy* 93:312–328.
- Biuw M., McConnell B., Bradshaw C.J.A., Burton H., Fedak M. 2003. Blubber and buoyancy: monitoring the body condition of free-ranging seals using simple dive characteristics. *Journal of Experimental Biology* 206:3405–3423.
- Bligh E.G., Dyer W.J. 1959. A rapid method of total lipid extraction and purification. *Canadian Journal of Biochemistry and Physiology* 37:911–917.
- Box J.E., ET AL. 2019. Key indicators of Arctic climate change: 1971–2017. *Environmental Research Letters* 14:045010.
- Brand U., Came R.E., Affek H., Azmy K., Mooi R., Layton K. 2014. Climate-forced change in Hudson Bay seawater composition and temperature, Arctic Canada. *Chemical Geology* 388:78–86.
- Burek K.A., Gulland F.M.D., O'Hara T.M. 2008. Effects of climate change on Arctic marine mammal health. *Ecological Applications* 18:126–134.
- Busch D.S., Hayward L.S. 2009. Stress in a conservation context: a discussion of glucocorticoid actions and how levels change with conservation-relevant variables. *Biological Conservation* 142:2844–2853.
- Chabot D., Stenson G.B. 2002. Growth and seasonal fluctuations in size and condition of male Northwest Atlantic harp seals *Phoca groenlandica*: an analysis using sequential growth curves. *Marine Ecology Progress Series* 227:25–42.
- Chabot D., Stenson G.B., Cadigan N.G. 1996. Short- and long-term fluctuations in the size and condition of harp seal (*Phoca groenlandica*) in the Northwest Atlantic. *NAFO Scientific Council Studies* 26:15–32.
- Chambellant M., Stirling I., Ferguson S.H. 2013. Temporal variation in western Hudson Bay ringed seal *Phoca hispida* diet in relation to environment. *Marine Ecology Progress Series* 481:269–287.
- Champagne C.D., Kellar N.M., Crocker D.E., Wasser S.K., Booth R.K., Trego M.L., Houser D.S. 2017. Blubber cortisol qualitatively reflects circulating cortisol concentrations in bottlenose dolphins. *Marine Mammal Science* 33:134–153.
- Champagne C.D., Crocker D.E., Fowler M.A., Houser D.S. 2012a. Fasting physiology of the pinnipeds: the challenges of fasting while maintaining high energy expenditure and nutrient delivery for lactation. In: McCue M.D., editor. *Comparative physiology of fasting, starvation, and food limitation*. Springer, Berlin, Heidelberg; p. 309–336.
- Champagne C.D., Houser D.S., Costa D.P., Crocker D.E. 2012b. The effects of handling and anesthetic agents on the stress response and carbohydrate metabolism in northern elephant seals. *PLoS ONE* 7:e38442.
- Champagne C.D., Houser D.S., Crocker D.E. 2006. Glucose metabolism during lactation in a fasting animal, the northern elephant seal. *American Journal of Physiology—Regulatory, Integrative and Comparative Physiology* 291:1129–1137.
- Champagne C.D., Tift M., Houser D., Crocker D. 2015. Adrenal sensitivity to stress is maintained despite variation in baseline glucocorticoids in moulting seals. *Conservation Physiology* 3:1–11.
- Cohen R.E., ET AL. 2018. Marine host-pathogen dynamics: influences of global climate change. *Oceanography* 31:182–193.
- Committee on Marine Mammals. 1967. Standard measurements of seals. *Journal of Mammalogy* 48:459–462.
- Costa D.P. 1993. The relationship between reproductive and foraging energetics and the evolution of the Pinnipedia. *Symposia of the Zoological Society of London* 66:293–314.
- Crawford J.A., Frost K.J., Quakenbush L.T., Whiting A. 2012. Different habitat use strategies by subadult and adult ringed seals (*Phoca hispida*) in the Bering and Chukchi seas. *Polar Biology* 35:241–255.
- Crawford J.A., Quakenbush L.T., Citta J.J. 2015. A comparison of ringed and bearded seal diet, condition and productivity between historical (1975–1984) and recent (2003–2012) periods in the Alaskan Bering and Chukchi seas. *Progress in Oceanography* 136:133–150.
- Crespi E.J., Williams T.D., Jessop T.S., Delehanty B. 2013. Life history and the ecology of stress: how do glucocorticoid hormones influence life-history variation in animals? *Functional Ecology* 27:93–106.
- Dantzer B., Fletcher Q.E., Boonstra R., Sheriff M.J. 2014. Measures of physiological stress: a transparent or opaque window into the status, management and conservation of species? *Conservation Physiology* 2:cou023.
- Das K., Siebert U., Fontaine M., Jauniaux T., Holsbeek L., Bouquegneau J.M. 2004. Ecological and pathological factors related to trace metal concentrations in harbour porpoises *Phocoena phocoena* from the North Sea and adjacent areas. *Marine Ecology Progress Series* 281:283–295.
- DeRango E.J., Greig D.J., Gálvez C., Norris T.A., Barbosa L., Elorriaga-Verplancken F.R., Crocker D.E. 2019. Response to capture stress involves multiple corticosteroids and is associated with serum thyroid hormone concentrations in Guadalupe fur seals (*Arctocephalus philippii townsendi*). *Marine Mammal Science* 35:72–92.
- Doi H., Akamatsu F., González A.L. 2017. Starvation effects on nitrogen and carbon stable isotopes of animals: an insight from meta-analysis of fasting experiments. *Royal Society Open Science* 4:170633.
- Doney S.C., ET AL. 2012. Climate change impacts on marine ecosystems. In: Carlson C.A., Giovanni S.J., editors. *Annual review*

- of marine science. Annual Reviews, Palo Alto, California, USA; p. 11–37.
- Evans P.G.H., Bjørge A. 2013. Impacts of climate change on marine mammals. *MCCIP Science Review* 2013:134–148. doi:10.14465/2013.arc15.134-148.
- Ferguson S.H., Higdon J.W. 2006. How seals divide up the world: environment, life history, and conservation. *Oecologia* 150:318–329.
- Ferguson S.H., Stirling I., McLoughlin P. 2005. Climate change and ringed seal (*Phoca hispida*) recruitment in Western Hudson Bay. *Marine Mammal Science* 21:121–135.
- Ferguson S.H., Young B.G., Yurkowski D.J., Anderson R., Willing C., Nielsen O. 2017. Demographic, ecological, and physiological responses of ringed seals to an abrupt decline in sea ice availability. *PeerJ* 5:e2957.
- Freitas C., Kovacs K.M., Ims R.A., Fedak M.A., Lydersen C. 2008. Ringed seal post-moulting movement tactics and habitat selection. *Oecologia* 155:193–204.
- Galbraith P.S., Larouche P. 2011. Reprint of “Sea-surface temperature in Hudson Bay and Hudson Strait in relation to air temperature and ice cover breakup, 1985–2009”. *Journal of Marine Systems* 88:463–475.
- Galimberti F., Sanvito S., Braschi C., Boitani L. 2007. The cost of success: reproductive effort in male southern elephant seals (*Mirounga leonina*). *Behavioral Ecology and Sociobiology* 62:159–171.
- Galligan T.M., Schwacke L.H., McFee W.E., Boggs A.S.P. 2018. Evidence for cortisol–cortisone metabolism by marine mammal blubber. *Marine Biology* 165:114.
- Gaston A.J., Elliott K.H. 2014. Seabird diet changes in northern Hudson Bay, 1981–2013, reflect the availability of schooling prey. *Marine Ecology Progress Series* 513:211–223.
- Gaston A.J., Woo K., Hipfner J.M. 2003. Trends in forage fish populations in Northern Hudson Bay since 1981, as determined from the diet of nestling thick-billed murres *Uria lomvia*. *Arctic* 56:227–233.
- Ghosh S., Rubly C. 2015. The emergence of Arctic shipping: issues, threats, costs, and risk-mitigating strategies of the Polar Code. *Australian Journal of Maritime & Ocean Affairs* 7:171–182.
- Gómez-Campos E., Borrell A., Aguilar A. 2011. Nitrogen and carbon stable isotopes do not reflect nutritional condition in the striped dolphin. *Rapid Communications in Mass Spectrometry* 25:1343–1347.
- Graham B.S., Koch P.L., Newsome S.D., McMahon K.W., Aurioles D. 2010. Using isoscapes to trace the movements and foraging behavior of top predators in oceanic ecosystems. In: West J.B., Bowen G.J., Dawson T.E., Tu K.P., editors. *Isoscapes: understanding movement, pattern, and process on Earth through isotope mapping*. 1st ed. Springer Netherlands, Dordrecht, The Netherlands; p. 299–318.
- Gulland F.M.D., Haulena M., Lowenstine L.J., Munro C., Graham P.A., Bauman J., Harvey J. 1999. Adrenal function in wild and rehabilitated Pacific harbor seals (*Phoca vitulina richardii*) and in seals with phocine herpesvirus-associated adrenal necrosis. *Marine Mammal Science* 15:810–827.
- Guo Y., Gui D., Zhang X., Liu W., Xie Q., Yu X., Wu Y. 2022. Blubber cortisol-based approach to explore the endocrine responses of Indo-Pacific humpback dolphins (*Sousa chinensis*) to diet shifts and contaminant exposure. *Environmental Science and Technology* 56:1069–1080.
- Hamilton C.D., Kovacs K.M., Ims R.A., Aars J., Lydersen C. 2017. An Arctic predator–prey system in flux: climate change impacts on coastal space use by polar bears and ringed seals. *Journal of Animal Ecology* 86:1054–1064.
- Hammill M.O., Lydersen C., Ryg M.S., Smith T.G. 1991. Lactation in the ringed seal (*Phoca hispida*). *Canadian Journal of Fisheries and Aquatic Sciences* 48:2471–2476.
- Hamilton C.D., Lydersen C., Ims R.A., Kovacs K.M. 2015. Predictions replaced by facts: a keystone species’ behavioural responses to declining arctic sea-ice. *Biology Letters* 11:1–6.
- Harlow H.J., Beck T.D.I., Walters L.M., Greenhouse S.S. 1990. Seasonal serum glucose, progesterone, and cortisol levels of black bears (*Ursus americanus*). *Canadian Journal of Zoology* 68:183–187.
- Harwood L.A., Smith T.G., Auld J.C. 2012. Fall migration of ringed seals (*Phoca hispida*) through the Beaufort and Chukchi seas, 2001–02. *Arctic* 65:35–44.
- Harwood L.A., Smith T.G., Auld J.C., Melling H., Yurkowski D.J. 2015. Seasonal movements and diving of ringed seals, *Pusa hispida*, in the Western Canadian Arctic, 1999–2001 and 2010–11. *Arctic* 68:193–209.
- Hauser D.D.W., Laidre K.L., Stern H.L. 2018. Vulnerability of arctic marine mammals to vessel traffic in the increasingly ice-free northwest passage and Northern Sea Route. *Proceedings of the National Academy of Sciences of the United States of America* 115:7617–7622.
- Hazen E.L., Abrahms B., Brodie S., Carroll G., Jacox M.G., Savoca M.S., Scales K.L., Sydeman W.J., Bograd S.J. 2019. Marine top predators as climate and ecosystem sentinels. *Frontiers in Ecology and the Environment* 17:565–574.
- Hertz E., Trudel M., Cox M.K., Mazumder A. 2015. Effects of fasting and nutritional restriction on the isotopic ratios of nitrogen and carbon: a meta-analysis. *Ecology and Evolution* 5:4829–4839.
- Hobson K.A., Alisauskas R.T., Clark R.G. 1993. Stable-nitrogen isotope enrichment in avian tissues due to fasting and nutritional stress: implications for isotopic analyses of diet. *The Condor* 95:388.
- Hobson K.A., Fisk A., Karnovsky N., Holst M., Gagnon J.M., Fortier M. 2002. A stable isotope ($\delta^{13}\text{C}$, $\delta^{15}\text{N}$) model for the North Water food web: implications for evaluating trophodynamics and the flow of energy and contaminants. *Deep Sea Research Part II: Topical Studies in Oceanography* 49:5131–5150.
- Holst M., Stirling I., Calvert W. 1999. Age structure and reproductive rates of ringed seals (*Phoca hispida*) on the northwestern coast of Hudson Bay in 1991 and 1992. *Marine Mammal Science* 15:1357–1364.
- Hop H., ET AL. 2020. Changes in sea-ice protist diversity with declining sea ice in the Arctic Ocean from the 1980s to 2010s. *Frontiers in Marine Science* 7:243.
- Hunt K.E., Lysiak N.S., Moore M., Rolland R.M. 2017. Multi-year longitudinal profiles of cortisol and corticosterone recovered from baleen of North Atlantic right whales (*Eubalaena glacialis*). *General and Comparative Endocrinology* 254:50–59.
- Huntington H.P., Quakenbush L.T., Nelson M. 2017. Evaluating the effects of climate change on Indigenous marine mammal hunting in Northern and Western Alaska using traditional knowledge. *Frontiers in Marine Science* 4:319.
- Hwang B., Aksenov Y., Blockley E., Tsamados M., Brown T., Landy J., Stevens D., Wilkinson J. 2020. Impacts of climate change on global sea ice. *MCCIP Science Review* 2020:208–227.
- Iacozza J., Ferguson S.H. 2014. Spatio-temporal variability of snow over sea ice in western Hudson Bay, with reference to ringed seal pup survival. *Polar Biology* 37:817–832.
- IPCC. 2014. In: Core Writing Team, Pachauri R.K., Meyer L.A., editors. *Climate change 2014: synthesis report. Contribution of Working Groups I, II and III to the fifth assessment report of the Intergovernmental Panel on Climate Change*. Geneva, Switzerland.

- Jackson A.L., Inger R., Parnell A.C., Bearhop S. 2011. Comparing isotopic niche widths among and within communities: SIBER—Stable Isotope Bayesian Ellipses in R. *Journal of Animal Ecology* 80:595–602.
- Kelly B.P., Badajos O.H., Kunasranta M., Moran J.R., Martinez-Bakker M., Wartzok D., Boveng P. 2010. Seasonal home ranges and fidelity to breeding sites among ringed seals. *Polar Biology* 33:1095–1109.
- Keogh M.J., Charapata P., Karpovich S., Jones A., Sprowls C., Marshall C.D. 2020a. Reproductive and stress-related hormones in whiskers from two North Pacific phocids: harbor and ringed seals. *Marine Mammal Science* 36:1322–1333.
- Keogh M.J., Gastaldi A., Charapata P., Melin S., Fadely B.S. 2020b. Stress-related and reproductive hormones in hair from three north Pacific otariid species: Steller sea lions, California sea lions and northern fur seals. *Conservation Physiology* 8:coaa069.
- Kershaw J.L., Hall A.J. 2016. Seasonal variation in harbour seal (*Phoca vitulina*) blubber cortisol—a novel indicator of physiological state? *Scientific Reports* 6:21889.
- Kortsch S., Primicerio R., Fossheim M., Dolgov A.V., Aschan M.M. 2015. Climate change alters the structure of arctic marine food webs due to poleward shifts of boreal generalists. *Proceedings of the Royal Society of London, B: Biological Sciences* 282:20151546.
- Kovacs K.M., Lydersen C. 2008. Climate change impacts on seals and whales in the North Atlantic Arctic and adjacent shelf seas. *Science Progress* 91:117–150.
- Kovacs K.M., Lydersen C., Overland J.E., Moore S.E. 2011. Impacts of changing sea-ice conditions on Arctic marine mammals. *Marine Biodiversity* 41:181–194.
- Kowal S., Gough W.A., Butler K. 2017. Temporal evolution of Hudson Bay sea ice (1971–2011). *Theoretical and Applied Climatology* 127:753–760.
- Krafft B.A., Kovacs K.M., Frie A.K., Haug T., Lydersen C. 2006. Growth and population parameters of ringed seals (*Pusa hispida*) from Svalbard, Norway, 2002–2004. *ICES Journal of Marine Science* 63:1136–1144.
- Krafft B.A., Kovacs K.M., Lydersen C. 2007. Distribution of sex and age groups of ringed seals *Pusa hispida* in the fast-ice breeding habitat of Kongsfjorden, Svalbard. *Marine Ecology Progress Series* 335:199–206.
- Kurle C.M., Worthy G.A.J. 2001. Stable isotope assessment of temporal and geographic differences in feeding ecology of northern fur seals (*Callorhinus ursinus*) and their prey. *Oecologia* 126:254–265.
- Laidre K.L., ET AL. 2015. Arctic marine mammal population status, sea ice habitat loss, and conservation recommendations for the 21st century. *Conservation Biology* 29:724–737.
- Laidre K.L., Stirling I., Lowry L.F., Wiig O., Heide-Jørgensen M.P., Ferguson S.H. 2008. Quantifying the sensitivity of Arctic marine mammals to climate-induced habitat change. *Ecological Applications* 18:S97–S125.
- Loseto L.L., Pleskach K., Hoover C., Tomy G.T., Desforjes J.-P., Halldorson T., Ross P.S. 2017. Cortisol levels in beluga whales (*Delphinapterus leucas*): setting a benchmark for Marine Protected Area monitoring. *Arctic Science* 372:358–372.
- Lübcker N., Whiteman J.P., Millar R.P., de Bruyn P.J.N., Newsome S.D. 2020. Fasting affects amino acid nitrogen isotope values: a new tool for identifying nitrogen balance of free-ranging mammals. *Oecologia* 193:53–65.
- Lüdecke D., Ben-Shachar M., Patil I., Waggoner P., Makowski D. 2021. performance: an R package for assessment, comparison and testing of statistical models. *Journal of Open Source Software* 6:3139.
- Lydersen C. 1995. Energetics of pregnancy, lactation and neonatal development in ringed seals (*Phoca hispida*). *Developments in Marine Biology* 4:319–327.
- Lydersen C., Ryg M.S., Hammill M.O., O'Brien P.J. 1992. Oxygen stores and aerobic dive limit of ringed seals (*Phoca hispida*). *Canadian Journal of Zoology* 70:458–461.
- Macbeth B.J., Cattet M.R.L., Obbard M.E., Middel K., Janz D.M. 2012. Evaluation of hair cortisol concentration as a biomarker of long-term stress in free-ranging polar bears. *Wildlife Society Bulletin* 36:747–758.
- MacDougall-Shackleton S.A., Bonier F., Romero L.M., Moore I.T. 2019. Glucocorticoids and “stress” are not synonymous. *Integrative Organismal Biology* 1:obz017.
- McLaren I.A. 1958. The biology of the ringed seal (*Phoca hispida* Schreher) in the Eastern Canadian Arctic. *Bulletin of the Fisheries Research Board of Canada* 118:97.
- McWhinnie L.H., Halliday W.D., Insley S.J., Hilliard C., Canessa R.R. 2018. Vessel traffic in the Canadian Arctic: management solutions for minimizing impacts on whales in a changing northern region. *Ocean and Coastal Management* 160:1–17.
- Mesnick S.L., Ralls K. 2009. In: Perrin W.F., Würsig B., Thewissen J.G.M., editors. *Mating systems. Encyclopedia of marine mammals*. 3rd ed. San Diego, California, USA; p. 712–719.
- Mislan P., Derocher A.E., Louis V.L., Richardson E., Lunn N.J., Janz D.M. 2016. Assessing stress in Western Hudson Bay polar bears using hair cortisol concentration as a biomarker. *Ecological Indicators* 71:47–54.
- Ogloff W.R., Ferguson S.H., Fisk A.T., Marcoux M., Hussey N.E., Jaworenko A., Yurkowski D.J. 2021. Long-distance movements and associated diving behaviour of ringed seals (*Pusa hispida*) in the eastern Canadian Arctic. *Arctic Science* 7:494–511.
- Ogloff W.R., Yurkowski D.J., Davoren G.K., Ferguson S.H. 2019. Diet and isotopic niche overlap elucidate competition potential between seasonally sympatric phocids in the Canadian Arctic. *Marine Biology* 166:103.
- Ortiz R.M. 2001. Effects of prolonged fasting on plasma cortisol and TH in postweaned northern elephant seal pups. *American Journal of Physiology—Regulatory, Integrative and Comparative Physiology* 280:R790–R795.
- Ostertag S.K., Loseto L.L., Snow K., Lam J., Hynes K., Gillman D.V. 2018. “That’s how we know they’re healthy”: the inclusion of traditional ecological knowledge in beluga health monitoring in the Inuvialuit Settlement Region. *Arctic Science* 320:1–29.
- Pecl G.T., ET AL. 2017. Biodiversity redistribution under climate change: impacts on ecosystems and human well-being. *Science* 355:eaai9214.
- Pinsky M.L., Worm B., Fogarty M.J., Sarmiento J.L., Levin S.A. 2013. Marine taxa track local climate velocities. *Science* 341:1239–1242.
- Polischuk S.C., Hobson K.A., Ramsay M.A. 2001. Use of stable-carbon and -nitrogen isotopes to assess weaning and fasting in female polar bears and their cubs. *Canadian Journal of Zoology* 79:499–511.
- R Core Team. 2019. R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <https://www.r-project.org>.
- Romero L.M., Butler L.K. 2007. Endocrinology of stress. *International Journal of Comparative Psychology* 20:89–95.
- Romero L.M., Gormally B.M.G. 2019. How truly conserved is the “well-conserved” vertebrate stress response? *Integrative and Comparative Biology* 59:273–281.

- Roy V., Iken K., Gosselin M., Tremblay J.E., Bélanger S., Archambault P. 2015. Benthic faunal assimilation pathways and depth-related changes in food-web structure across the Canadian Arctic. *Deep Sea Research Part I: Oceanographic Research Papers* 102:55–71.
- Ryg M., Smith T.G., Øritsland N.A. 1990. Seasonal changes in body mass and body composition of ringed seals (*Phoca hispida*) on Svalbard. *Canadian Journal of Zoology* 68:470–475.
- Sapolsky R.M., Romero L.M., Munck A.U. 2000. How do glucocorticoids influence stress responses? Integrating permissive, suppressive, stimulatory, and preparative actions. *Endocrine Reviews* 21:55–89.
- Sheriff M.J., Krebs C.J., Boonstra R. 2010. The ghosts of predators past: population cycles and the role of maternal programming under fluctuating predation risk. *Ecology* 91:2983–2994.
- Smith T.G. 1987. The Ringed Seal, *Phoca hispida*, of the Canadian Western Arctic. *Canadian Bulletin of Fisheries and Aquatic Science* 216:81.
- Spurlin S.M., Peterson S.H., Crocker D.E., Costa D.P. 2019. Nitrogen and carbon stable-isotope ratios change in adult northern elephant seals (*Mirounga angustirostris*) during the breeding and molting fasts. *Marine Mammal Science* 35:707–717.
- Steinman K.J., Robeck T.R., O'Brien J.K. 2016. Characterization of estrogens, testosterone, and cortisol in normal bottlenose dolphin (*Tursiops truncatus*) pregnancy. *General and Comparative Endocrinology* 226:102–112.
- Stirling I. 2005. Reproductive rates of ringed seals and survival of pups in Northwestern Hudson Bay, Canada, 1991–2000. *Polar Biology* 28:381–387.
- Strandberg U., Käkälä A., Lydersen C., Kovacs K.M., Grahl-Nielsen O., Hyvärinen H., Käkälä R. 2008. Stratification, composition, and function of marine mammal blubber: the ecology of fatty acids in marine mammals. *Physiological and Biochemical Zoology* 81:473–485.
- Teilmann J., Born E., Acquarone M. 1999. Behaviour of ringed seals tagged with satellite transmitters in the North Water polynya during fast-ice formation. *Canadian Journal of Zoology* 77:1934–1946.
- Thometz N.M., Hermann-Sorensen H., Russell B., Rosen D.A.S., Reichmuth C. 2020. Molting strategies of Arctic seals drive annual patterns in metabolism. *Conservation Physiology* 9:coaa112.
- Tilbrook A.J., Turner A.I., Clarke I.J. 2000. Effects of stress on reproduction in non-rodent mammals: the role of glucocorticoids and sex differences. *Reviews of Reproduction* 5:105–113.
- Trana M.R., Roth J.D., Tomy G.T., Anderson W.G., Ferguson S.H. 2015. Influence of sample degradation and tissue depth on blubber cortisol in beluga whales. *Journal of Experimental Marine Biology and Ecology* 462:8–13.
- Trana M.R., Roth J.D., Tomy G.T., Anderson W.G., Ferguson S.H. 2016. Increased blubber cortisol in ice-entrapped beluga whales (*Delphinapterus leucas*). *Polar Biology* 39:1563–1569.
- Ward C.L., McCann K.S. 2017. A mechanistic theory for aquatic food chain length. *Nature Communications* 8:2028.
- Wathne J.A., Haug T., Lydersen C. 2000. Prey preference and niche overlap of ringed seals *Phoca hispida* and harp seals *P. groenlandica* in the Barents Sea. *Marine Ecology Progress Series* 194:233–239.
- Watt C.A., Orr J., Ferguson S.H. 2016. A shift in foraging behaviour of beluga whales *Delphinapterus leucas* from the threatened Cumberland Sound population may reflect a changing Arctic food web. *Endangered Species Research* 31:259–270.
- Webster M.A., Rigor I.G., Nghiem S. V., Kurtz N.T., Farrell S.L., Perovich D.K., Sturm M. 2014. Interdecadal changes in snow depth on Arctic sea ice. *Journal of Geophysical Research, Oceans* 119:5395–5406.
- Young B.G., Ferguson S.H. 2013. Seasons of the ringed seal: pelagic open-water hyperphagy, benthic feeding over winter and spring fasting during molt. *Wildlife Research* 40:52–60.
- Yurkowski D.J., Ferguson S.H., Choy E.S., Loseto L.L., Brown T.M., Muir D.C.G., Semeniuk C.A.D., Fisk A.T. 2016a. Latitudinal variation in ecological opportunity and intraspecific competition indicates differences in niche variability and diet specialization of Arctic marine predators. *Ecology and Evolution* 6:1666–1678.
- Yurkowski D.J., Ferguson S.H., Semeniuk C.A.D., Brown T.M., Muir D.C.G., Fisk A.T. 2016b. Spatial and temporal variation of an ice-adapted predator's feeding ecology in a changing Arctic marine ecosystem. *Oecologia* 180:631–644.
- Yurkowski D.J., Hussey N.E., Ferguson S.H., Fisk A.T. 2018. A temporal shift in trophic diversity among a predator assemblage in a warming Arctic. *Royal Society Open Science* 5:180259.
- Vander Zanden M.J., Fetzner W.W. 2007. Global patterns of aquatic food chain length. *Oikos* 116:1378–1388.

Submitted 24 August 2021. Accepted 23 April 2022.

Associate Editor was Elizabeth Flaherty.