



---

## **The Cretaceous Scelionid Genus *Proteroscelio* Brues (Hymenoptera: Platygastroidea)**

Authors: Johnson, Norman F., Musetti, Luciana, and Masner, Lubomír

Source: American Museum Novitates, 2008(3603) : 1-7

Published By: American Museum of Natural History

URL: [https://doi.org/10.1206/0003-0082\(2008\)3603\[1:TCSGPB\]2.0.CO;2](https://doi.org/10.1206/0003-0082(2008)3603[1:TCSGPB]2.0.CO;2)

---

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

# AMERICAN MUSEUM *Novitates*

PUBLISHED BY THE AMERICAN MUSEUM OF NATURAL HISTORY  
CENTRAL PARK WEST AT 79TH STREET, NEW YORK, NY 10024

Number 3603, 7 pp., 1 figure

April 9, 2008

## The Cretaceous Scelionid Genus *Proteroscelio* Brues (Hymenoptera: Platygastridae)

NORMAN F. JOHNSON<sup>1</sup>, LUCIANA MUSETTI<sup>2</sup>, AND LUBOMÍR MASNER<sup>3</sup>

### ABSTRACT

The genus *Proteroscelio* Brues is redescribed and *P. gravatus*, n. sp., is described from Lebanese amber (Aptian age, 112–122 mya). The relationships between *Proteroscelio* and other scelionids is discussed. The described species of fossil platygastroids are tabulated. The taxa represented by the unavailable names “*Eopteromalites fushunensis*” Hong, “*Leptogasterites brunneus*” Hong, “*L. furvus*” Hong, and “*Sinilongicapito guchengziensis*” Hong, recently described from Fushun, Liaoning, China (50 mya), should all be classified as scelionids. The replacement name *Sinoprotelenomus* Zhang n. name is proposed for *Protelenomus* Zhang, 1989 (preoccupied by *Protelenomus* Kieffer, 1906).

### INTRODUCTION

Despite a paucity of described species, the parasitoid wasp superfamily Platygastridae (Scelionidae and Platygastridae) is cited as one of the dominant groups of Hymenoptera found in Cretaceous fossils (Grimaldi et al., 2002; Zherikhin and Sukacheva, 1971). Only five species from that time have been named to date (Carpenter et al., 1937; Nel and Azar,

2005; Schlüter, 1978), and most described fossil platygastroids are from Baltic amber inclusions (see appendix). A clarification of the relationships of the superfamily with the other major monophyletic groups of Hymenoptera would contribute substantially to elucidation of the phylogeny of all of the suborder Apocrita (Austin et al., 2005). Therefore, understanding the diversity and characteristics of these earliest elements of

<sup>1</sup>Research Associate, Division of Invertebrate Zoology, American Museum of Natural History; Professor, Department of Entomology, The Ohio State University, 1315 Kinnear Road, Columbus, OH 43212 (johnson.2@osu.edu).

<sup>2</sup>Research Associate, Department of Entomology, The Ohio State University, 1315 Kinnear Road, Columbus, OH 43212 (musetti.2@osu.edu).

<sup>3</sup>Research Associate, Division of Invertebrate Zoology, American Museum of Natural History; Research Associate, Agriculture and Agri-Food Canada, Research Branch, K. W. Neatby Building, Ottawa, Ontario K1A 0C6, Canada.

Platygastroidea promises to be an important element in the goal of resolving hymenopteran relationships.

Charles T. Brues described the genus *Proteroscelio* from a single fossil from Cedar Lake, Manitoba (Carpenter et al., 1937). In his description he drew attention to the significance of the 14-merous antenna, a characteristic at that time unknown among extant species. Brues referred to an undescribed species in Baltic amber also with 14-merous antenna, a taxon he later described as the genus *Archaeoscelio* (Brues, 1940). Today, a third genus is known with this feature, *Nixonia* Masner, an extant group of 14 species from tropical Africa and Asia (Johnson and Masner, 2006).

We describe and illustrate below the type species of *Proteroscelio*, *P. antennalis* Brues, and also describe a new, similar, but significantly older species, *P. gravatus*, n. sp. This new species is from Lebanese amber of lower Aptian age (112–122 mya).

## MATERIALS

The new species described below was originally recognized as a species of *Proteroscelio* by L. Masner. This work is based on specimens in the following collections:

MNHN	A. Nel: Laboratoire d'Entomologie, Muséum National d'Histoire Naturelle, Paris, France
ROMC	J. Waddington: Department of Palaeobiology, Royal Ontario Museum, Toronto, Canada

Abbreviations and terms used in text: **A1**, **A2**, ... **A14**: antennomere 1, 2, ... 14; **claval formula**: distribution of the large, multiporous basiconic sensilla on the underside of apical antennomeres of the female, with the segment interval specified followed by the number of sensilla per segment (Bin, 1981); **epomial carina**: the vertical portion of epomium on the pronotum; **pronotal humeral carina**: the horizontal portion of epomium on the pronotum; **S1**, **S2**, ... **S6**: metasomal sterna 1, 2, ... 6; **T1**, **T2**, ... **T7**: metasomal tergum 1, 2, ... 7. Morphological terminology follows Masner (1980) and Mikó et al. (2007). Dates for geological periods are from Grimaldi and Engel (2005).

## *PROTEROSCELIO* BRUES


*Proteroscelio* Brues, in Carpenter et al., 1937: 39. Original description. Type: *Proteroscelio antennalis* Brues, by monotypy and original designation. Walker, 1934: pl. 1, fig. 2; Muesebeck and Walkley, 1956: 392, citation of type species; Carpenter, 1992: 471, diagnosis; Johnson, 1992: 467, catalog of species.

**DESCRIPTION:** Body length 1.1–1.7 mm, gracile, mesosoma dorsoventrally depressed, head anteroposteriorly compressed.

Head wider than long, strongly transverse when viewed dorsally; vertex rounded, hyper-occipital carina absent; occipital carina absent; lateral ocellus distinctly separated from inner orbit by distance greater than one ocellar diameter; compound eye large, apparently bare. frons flat, without frontal depression, median longitudinal carina absent; no inter-antennal prominence visible; submedian carina absent; orbital carina absent; lower frons without fanlike striae; width of interocular space very broad, distinctly greater than eye height; clypeal region, mouthparts not clearly visible; antenna 14-merous; radicle inserted apically into base of A1, more or less parallel to longitudinal axis of A1; apical antennomeres forming a clava, either laterally compressed or generally enlarged; claval formula at least A7–A14/1–2–2–2–2–2–1.

Mesosoma strongly depressed; pronotum in dorsal view campanulate, ecarinate; netrion present as a narrow fusiform sclerite, closed ventrally; anterior margin of mesoscutum meeting pronotum dorsally; mesoscutum semioval in outline; notauli absent; parapsidal lines not distinguishable; skaphion absent; transscutal articulation well developed; scutellum semicircular, unarmed, flattened; mesopleuron strongly inclined anteriorly; mesopleural depression broad, deep; mesopleural carina absent; details of sculpture of meso-, metapleural area obscured; legs elongate, slender; femora weakly incrassate; trochantellus present on all legs; outer surface of tibiae without visible spines; tibial spur formula 1–2–2, spurs on mid- and hind tibiae fine, short; tarsal formula 5–5–5; tarsomeres tapering in width apically, cylindrical; pretarsal claws simple; forewing extending nearly to apex of metasoma; R slightly, but distinctly bent at origin of basal vein, extending through basal 0.6 of length of forewing, without large





Figs. 1–2. *Proteroscelio antennalis* Brues, holotype female. 1, Ventrolateral habitus. 2, Head and mesosoma, ventrolateral view. Figs. 3–4. *Proteroscelio gravatus*, n. sp. 3, Dorsolateral habitus. 4, Ventrolateral habitus. Scale bars in millimeters.

A7–A13 cylindrical, evenly expanded; T3 longest metasomatic tergite, slightly longer than T2.

**DIAGNOSIS:** Distinguished from *P. antennalis* by the very short funicular segments.

**MATERIAL EXAMINED:** Holotype female: “Ambre de HAMMANA/MDEIRIJ, LIBAN, Aptien inférieur, Collection Dany AZAR, Echantillon no 23.” Single block of amber mounted beneath a cover slip on a microscope slide. Deposited in MNHN.

**ETYMOLOGY:** The epithet *gravatus* is from the Latin meaning “weighed down”, referring to the massive antennal clava.

**COMMENTS:** The Aptian age, indicated on the specimen label, is defined in Grimaldi and Engel (2005) as extending from 112–122 mya. In the same text, the age of Lebanese amber is said to be between 120–135 mya. Nel and

Azar (2005) cite the age of *Cretaxenomerus*, collected at the same Hammana locality, to be from 125–135 mya.

## DISCUSSION

In the original description, Brues (in Carpenter et al., 1937) remarked that *Proteroscelio* “resembles the remarkable Austromalayan genus *Platyscelio* somewhat in the form of the antennae in the less strongly flattened head and thorax, but it [*Proteroscelio*] is much less highly modified.” The direct comparison of these two groups is superficial: the only real similarity lies in the flattened head and mesosoma.

For many years the consensus of opinion was that the most basal lineages of Scelionidae were the genera placed in the tribes Nixonini

(*Nixonia*) and Sparasionini (*Sparasion* Latreille, *Scelioromorpha* Ashmead, *Archaeoteleia* Masner, *Electroteleia* Brues) (e.g., Masner, 1976; Kozlov and Kononova 1983). Recent works (e.g., Austin and Field, 1997; Johnson and Masner, 2006; Masner et al., 2007) challenge this notion: Scelionidae may be paraphyletic because *Sparasion* and *Scelioromorpha* may be most closely related to Platygasteridae; *Archaeoteleia* possesses a tubular *Scelio*-type ovipositor and apparently is not closely related to other sparasionines; and an additional genus—*Neuroscelio* Dodd—must also be considered when addressing this question.

*Proteroscelio* generally has been considered to be a “primitive” scelionid, most likely because of the age of the fossil and the 14-merous antennae. It also has the 1–2–2 tibial spur formula, but the paired spurs on the mid and hind legs are fine and short, much more delicate than the robust structures in living genera. The metasoma is more or less evenly segmented, unlike most scelionids in which one segment—usually the second or third—is significantly longer than any other. *Proteroscelio* does have a fine but clearly developed malar sulcus, lacks the epomial and transverse carinae on the pronotum, lacks a clear bulla in the submarginal vein, and has the apex of the radicle inserted into the base of the scape more or less parallel to the long axis of the scape. It seems that the most appropriate interpretation is that *Proteroscelio* represents an ancient, but already highly specialized lineage.

A summary of the known fossil taxa of Platygastroidea is presented in the appendix. Few of these have been either critically or even recently examined, and their placement in this superfamily may well be subject to change. The names “*Eopteromalites*” Hong, “*Leptogasterites*” Hong, and “*Sinilongicapito*” Hong were originally described in the Chalcidoidea (Hong, 2002). They are actually members of the Scelionidae; this judgment is based upon the line drawings and photographs of the original amber specimens. However, these names are not available under the International Code of Zoological Nomenclature because the author did not specify a depository for the type specimens. He only generally

stated in the preface that “most” specimens are deposited in the Geological Museum of China.

#### ACKNOWLEDGMENTS

We thank J.-Y. Rasplus (Montpelier, France) and J. Heraty (Riverside, CA) for making available for study the specimen in Lebanese amber; to J.S. Noyes (London, UK) for alerting us to Hong (2002); and to Zhang Junfeng (Nanjing) for the contribution of the name *Sinoprotelenomus*. This material is based upon work supported in part by the National Science Foundation under grants No. DEB-0344034 and DEB-0614764.

#### REFERENCES

- Austin, A.D., N.F. Johnson, and M. Dowton. 2005. Systematics, evolution, and biology of scelionid and platygasterid wasps. *Annual Review of Entomology* 50: 553–582.
- Austin, A.D., and S.A. Field. 1997. The ovipositor system of scelionid and platygasterid wasps (Hymenoptera: Platygastroidea): comparative morphology and phylogenetic implications. *Invertebrate Taxonomy* 11: 1–87.
- Bin, F. 1981. Definition of female antennal clava based on its plate sensilla in Hymenoptera Scelionidae Telenominae. *Redia* 64: 245–261.
- Brues, C.T. 1940. Fossil parasitic Hymenoptera of the family Scelionidae in Baltic amber. *Proceedings of the American Academy of Arts and Sciences* 74: 69–90.
- Carpenter, F.M. 1992. Arthropoda 4. Superclass Hexopoda. *Treatise on Invertebrate Paleontology*. Part R. Boulder, CO: Geological Society of America, 655 pp.
- Carpenter, F.M., J.W. Folsom, E.O. Essig, A.C. Kinsey, C.T. Brues, M.W. Boesel, and H.E. Ewing. 1937. Insects and arachnids from Canadian amber. *University of Toronto Studies, Geological Series* 40: 7–62.
- Cockerell, T.D.A. 1909. Descriptions of Hymenoptera from Baltic amber. *Schriften der Physikalisch-Ökonomischen Gesellschaft zu Königsberg* 50: 1–20.
- Cockerell, T.D.A. 1915. Miocene fossil insects. *Proceedings of the Academy of Natural Sciences of Philadelphia* 66: 634–648.
- Cockerell, T.D.A. 1921. Fossil arthropods in the British Museum—V. Oligocene Hymenoptera from the Isle of Wight. *Annals and Magazine of Natural History* (9)7: 1–25.

- Grimaldi, D.A., and M.S. Engel. 2005. Evolution of the insects. Cambridge: Cambridge University Press, 755 pp.
- Grimaldi, D.A., M.S. Engel, and P.C. Nascimbene. 2002. Fossiliferous Cretaceous amber from Myanmar (Burma): its rediscovery, biotic diversity, and paleontological significance. *American Museum Novitates* 3361: 1–71.
- Hong, Y.-C. 2002. [Annals of amber insects of China.]. Beijing: Beijing Ke Xue Ji Shu Chu Ban She, 653 pp.
- Hope, F.W. 1837. Observations on succinic insects. *Transactions of the Entomological Society of London* 2: 46–57.
- Johnson, N.F. 1992. Catalog of world Proctotrupeoidea excluding Platygastriidae. *Memoirs of the American Entomological Institute* 51: 1–825.
- Johnson, N.F., and L. Masner. 2006. Revision of world species of the genus *Nixonia* (Hymenoptera: Platygastroidea, Scelionidae). *American Museum Novitates* 3518: 1–32.
- Kieffer, J.J. 1906. Description de quelques nouveaux serphides. *Bulletin de la Société d'Histoire Naturelle de Metz* 25: 1–7.
- Kozlov, M.A., and S.V. Kononova. 1983. [Telenominae of the fauna of the USSR]. Leningrad: Nauka, 336 pp.
- Maneval, H. 1938. Trois serphoïdes de l'ambre de la Baltique. *Revue Française d'Entomologie* 5: 107–116.
- Masner, L. 1969. A scelionid wasp surviving unchanged since Tertiary (Hymenoptera: Proctotrupeoidea). *Proceedings of the Entomological Society of Washington* 71: 397–400.
- Masner, L. 1976. Revisionary notes and keys to world genera of Scelionidae (Hymenoptera: Proctotrupeoidea). *Memoirs of the Entomological Society of Canada* 97: 1–87.
- Masner, L. 1980. Key to the genera of Scelionidae of the Holarctic region, with descriptions of new genera and species (Hymenoptera: Proctotrupeoidea). *Memoirs of the Entomological Society of Canada* 113: 1–54.
- Masner, L., N.F. Johnson, and A. Polaszek. 2007. Redescription of *Archaeoscelio* Brues and description of three new genera of Scelionidae (Hymenoptera): a challenge to the definition of the family. *American Museum Novitates* 3550: 1–24.
- Meunier, F. 1905. Nouvelles recherches sur quelques diptères et hyménoptères du copal fossile "dit de Zanzibar". *Revue Scientifique du Bourbonnais et du Centre de la France* 1905: 204–215.
- Meunier, F. 1917. Über einige Proctotrypidae (Bethylinae, Ceraphroninae und Scelioninae) aus dem subfossilen und dem rezenten Kopal von Zanzibar und von Madagaskar. *Zeitschrift der Deutschen Geologischen Gesellschaft* 68: 391–395.
- Mikó, I., L. Vilhelmsen, N.F. Johnson, L. Masner, and Z. Péntes. 2007. Skeletomusculature of Scelionidae (Hymenoptera: Platygastroidea): head and mesosoma. *Zootaxa* 1571: 1–78.
- Muesebeck, C.F.W., and L.M. Walkley. 1956. Type species of the genera and subgenera of parasitic wasps comprising the superfamily Proctotrupeoidea (order Hymenoptera). *Proceedings of the U.S. National Museum* 105: 319–419.
- Nel, A., and D. Azar. 2005. The oldest parasitic Scelionidae: Teleasinae (Hymenoptera: Platygastroidea). *Polskie Pismo Entomologiczne* 74: 333–338.
- Nel, A., and J. Prokop. 2005. New fossil Scelionidae (Insecta: Hymenoptera) in early Paleogene amber from eastern Moravia (Czech Republic) and northern France. *Polskie Pismo Entomologiczne* 74: 339–347.
- Schlüter, T. 1978. Zur Systematik und Palökologie harzkonservierter Arthropoda einer Taphozönose aus dem Cenomanium von NW-Frankreich. *Berliner geowissenschaftliche Abhandlungen (A)* 9: 1–150.
- Statz, G. 1938. Neue Funde parasitischer Hymenopteren aus dem Tertiär von Rott am Siebengebirge. *Decheniana* 98((A)): 71–154.
- Szabó, J.B., and J. Oehlke. 1986. Neue Proctotrupeoidea aus dem Baltischen Bernstein. *Beiträge zur Entomologie* 36: 99–106.
- Walker, T.L. 1934. Chemawinite or Canadian amber. *University of Toronto Studies Geological Series* 36: 5–10.
- Zhang, J.-F. 1989. [Fossil insects from Shanwang, Shandong, China]. Jinan: Shandong Science and Technology Publishing House.
- Zherikhin, V.V., and I.D. Sukacheva. 1971. [On Cretaceous insect-bearing ambers (retinites) of northern Siberia.] *In* *Doklady 24 Ezhegodnom Chtenii Pamyati N.A. Kholodkovskogo*: 3–48. Leningrad Nauka.

## APPENDIX 1

**Described species of fossil and subfossil  
Platygastridae**

Dates for geological periods are from Grimaldi and Engel (2005).

## Cretaceous Period

Aptian Age (Lebanese amber: 112–122 mya)  
*Cretoxenomerus jankotejai* Nel and Azar, 2005  
*Proteroscelio gravatus*, n. sp.

Cenomanian Age (unnamed formation: 100 mya)  
*Cenomanoscelio pulcher* Schlüter, 1978

Campanian Age (Cedar Lake, Canada: 80 mya)  
*Baeomorpha dubitata* Brues, 1937  
*Baryconus fulleri* Brues, 1937  
*Proteroscelio antennalis* Brues, 1937

## Tertiary Period

Paleocene to Middle Eocene (Bilé Karpaty, Belověž Formation, Czech Republic (<65 mya)  
*Moravoscelio bednariki* Nel and Prokop, 2005

Lower Eocene Epoch (Les Quesnoy, France: 55.6 mya)  
*Galloscelio pumilio* Nel and Prokop, 2005

Eocene Epoch (Fushun, Liaoning, China: 50 mya)

“*Eopteromalites fushunensis*” Hong, 2002  
“*Leptogasterites brunneus*” Hong, 2002  
“*Leptogasterites furvus*” Hong, 2002  
“*Sinilongicapito guchengziensis*” Hong, 2002

Middle Eocene Epoch (Baltic amber: 44 mya)

*Aneurobaeus collaris* Brues, 1940  
*Archaeoscelio filicornis* Brues, 1940  
*Archaeoscelio rugosus* Brues, 1940  
*Brachyscelio cephalotes* Brues, 1940  
*Brachyscelio dubius* Brues, 1940  
*Ceratobaeoides acuminatus* Brues, 1940  
*Ceratoteleia caudata* Brues, 1940  
*Ceratoteleia proleptica* Brues, 1940  
*Ceratoteleia succinophila* Brues, 1940  
*Chromoteleia theobaldi* Maneval, 1938

*Cobaloscelio cuspidatus* Johnson and Masner, 2007

*Cobaloscelio speculifer* Johnson and Masner, 2007

*Dissolcus electra* Brues, 1940  
*Electroteleia stigmatica* Brues, 1940  
*Hadronotoides dubitatus* Brues, 1940  
*Hadronotus electrinus* Cockerell, 1909  
*Hoploteleia doddii* Brues, 1940  
*Idris ilonkae* Szabó and Oehlke, 1986  
*Macroteleia renatae* Szabó and Oehlke, 1986  
*Mirotelenomus angulatus* Brues, 1940  
*Parabaeus pusillus* Brues, 1940  
*Proplatyscelio depressus* Brues, 1940  
*Pseudobaeus fecundulus* Brues, 1940  
*Sembilanocera clavata* Brues, 1940  
*Sparaison amabilis* Brues, 1940  
*Sparaison simplicifrons* Brues, 1940  
*Trachelopteron angulipenne* Brues, 1940  
*Uroteleia synthetica* Brues, 1940  
Late Eocene Epoch (Florissant: 38 mya)  
*Palaeoteleia oxyura* Cockerell, 1915

Eocene-Oligocene boundary (Isle of Wight: 38 mya)

*Macroteleia veterna* Cockerell, 1921

Late Oligocene Epoch (Chiapas amber: 23–28 mya)

*Palaeogryon muesebecki* Masner, 1969

Oligocene or Miocene Epoch (Rott: 20–28 mya)

*Platygasterites femoralis* Statz, 1938  
*Platygasterites spinosa* Statz, 1938  
*Scelionites capitatus* Statz, 1938

Middle Miocene Epoch (~18 mya)

*Sinoprotelenomus miocenicus* (Zhang,) **new combination**. *Sinoprotelenomus* **new name** is proposed by Dr. Zhang Junfeng for *Protelenomus* Zhang, 1989, preoccupied by *Protelenomus* Kieffer, 1906

Subfossil copal

*Acutibaeus bellicosus* Meunier, 1917  
*Calotelea aurantia* Hope, 1837  
*Ceratobaeus incertus* Meunier, 1905



Complete lists of all issues of the *Novitates* and the *Bulletin* are available at World Wide Web site <http://library.amnh.org/pubs>. Inquire about ordering printed copies via e-mail from [scipubs@amnh.org](mailto:scipubs@amnh.org) or via standard mail from: American Museum of Natural History, Library—Scientific Publications, Central Park West at 79th St., New York, NY 10024. TEL: (212) 769-5545. FAX: (212) 769-5009.

Ⓢ This paper meets the requirements of ANSI/NISO Z39.48-1992 (Permanence of Paper).