



Formicidae of the Andaman and Nicobar Islands (Indian Ocean: Bay of Bengal)

Authors: Mohanraj, Prashanth, Ali, Musthak, and Veenakumari, K.

Source: Journal of Insect Science, 10(172) : 1-11

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.010.14132>



Formicidae of the Andaman and Nicobar Islands (Indian Ocean: Bay Of Bengal)

Prashanth Mohanraj^{1,3a*}, Musthak Ali^{2b} and K. Veenakumari^{1c}

¹Central Agricultural Research Institute, P.B. No. 181, Port Blair 744 101, Andamans, India

²Department of Entomology, University of Agricultural Sciences, GKVK, Bangalore-560065, India

³Present address: Project Directorate of Biological Control, P.B. No.2491, Hebbal, Bangalore-560024, India

Abstract

Ants on the Andaman and Nicobar Islands, India were surveyed. These collections doubled the number of ant species recorded from these islands (from 59 to 125). Records include five endemic species, but no endemic genera. The surveys were fairly superficial, and it is likely many species remain to be discovered on these islands.

Keywords: Ants, Andamans, Nicobars

Correspondence: ^{a*} veenaprashi@rediffmail.com, ^b tmmusthakali@yahoo.com, ^c veenapmraj@rediffmail.com,

*Corresponding author

Received: 16 December 2008, **Accepted:** 15 January 2010

Copyright : This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.

ISSN: 1536-2442 | Vol. 10, Number 172

Cite this paper as:

Mohanraj P, Ali M, Veenakumari K. 2010. Formicidae of the Andaman and Nicobar Islands (Indian Ocean: Bay Of Bengal). *Journal of Insect Science* 10:172 available online: insectscience.org/10.172

Introduction

It has been 3 decades since McVean (1976) evaluated the status of zoological studies on the Andaman and Nicobar islands (India) and observed that many insect species needed to be studied as they have been "hitherto neglected [in studies on] the natural history of these islands". Along with a number of other insect groups, the Formicidae have continued to remain largely neglected, even after this observation was made. The only papers on the Formicidae of these islands during this period were by Chhotani and Maiti (1977) and Tiwari and Jonathan (1986a,b). Together these reports added a mere 11 species to the ant fauna already known from these islands, of which nine were new records and two were new species.

Mayr (1865) was perhaps the first to describe Formicidae from these islands, mainly from the Nicobars, based on collections made by the Austrian frigate *Novara*. Years later Forel (1903) listed a total of 39 species of ants, which included species from Mayr (1865), as well as the descriptions of three new species. The new descriptions included two species from the Andamans and one from the Nicobars. In the same year, Bingham (1903), while dealing with the ants of the Indian subcontinent in the Fauna of British India series, mentioned a mere two species from the Nicobars and none from the Andamans. Emery (1911, 1912, 1921, 1922) made some references to ants from these islands. The collection of Dr. N. Annandale, made during November and December 1923 from Mount Harriet (S. Andaman), was described by Mukherji and Ribeiro (1925). This included 15 species, of which two were recorded at the generic level only. Chhotani and Maiti (1977), working on another collection, the Zoological

Survey of India that was collected between February and April 1964, reported 15 species of ants, of which 10 were new records. Tiwari and Jonathan (1986a, 1986b) described two new species, one each from South Andaman and Great Nicobar. These species belong to genera that were not known previously from these islands. In his revisionary studies, Bolton (1987) added one more species to the ant fauna of these islands. In short, a total of 59 Formicidae species (of which, four were identified to the generic level only) were recorded from these islands between 1865 and 1987. No papers on the ants of these islands have appeared since then.


The Island Setting

Area

The Andaman - Nicobar chain of islands situated between 6° 45' - 13° 30' N and 92° - 94° E in the Bay of Bengal, Indian Ocean (Figure 1A, B) consists of 572 islands, islets and rocks (Anonymous 1986). The Andaman islands (6408 km²), which are more than three times larger than the Nicobars (1841 km²), are separated from the latter by the 150-km-wide Ten Degree Channel. In turn, each island group is fragmented by straits and channels of varying widths and depths.

Tectonics and sea level changes

These are true oceanic islands lying along the 6000 km long Sunda Arc, which extends west from the island of Sumatra to Burma in the north and marks the zone where the Indian-Australian plate is being subducted beneath the Eurasian plate (Moore et al. 1980; Curry 1989). Subduction, which is presumed to have commenced about 130 MYBP following the breakup of Gondwanaland in the early Cretaceous, resulting in the formation of oceanic ridges that were uplifted to their



current elevation as two arcs in the late Eocene or early Oligocene times (Curry et al. 1979). The outer arc ridge emerges intermittently above sea level as the Andaman-Nicobar islands, while the Barren and Narcondam islands are the emergent peaks of the submarine ridge that forms the inner volcanic arc (Hamilton 1979).

It is thought that in the Pleistocene, when sea levels were lower, there was no dry land connecting the islands to any of the adjacent land masses (Ripley and Beehler 1989). There is also no evidence for the existence of an earlier land bridge, even during Tertiary times (Nassig et al. 1996). During times of sea level lowering, however, the sea separating the Andaman islands from Burma was much narrower and shallower than that separating the southern Nicobars from Sumatra or the Malay Peninsula (Ripley and Beehler 1989).

Topography

The surface of the majority of the islands is irregular and hilly, with many narrow valleys. The hills, which follow the general direction of the islands, are oriented in the north-south direction, and from which arise numerous spurs and ridges that branch out in all directions. The hills on the east are higher than those on the west, with Saddle Peak (732 m) in North Andaman being the highest point in the Andamans and Mount Thuillier (642 m) in Great Nicobar is the highest in the Nicobars (Anonymous 1994).

Climate

The islands experience a tropical maritime climate that is strongly influenced by the Indian Ocean. They receive over 3000 mm of rainfall between May and December during both the South-West and the North-East monsoons. The mean minimum and mean maximum temperatures vary between 23° C

and 30° C with maximum temperatures of about 34° C attained in April. High humidity prevails throughout the year, ranging from more than 60 to about 90 percent.

Vegetation

The islands are densely wooded from the water's edge to the tops of the hills, except in those areas initially cleared by the European colonial powers and later by settlers from the Indian mainland. Native vegetation has been replaced to varying degrees by settlements and introduced plants (intentionally for cultivation and accidentally as weeds) on all of the 38 inhabited islands. Over the years, this has led to near extinction of the Andaman Giant Evergreen forests, one of the seven major forest types characterized by Champion and Seth (1968) as occurring on these islands.

Materials and Methods

Ants were collected from cultivated and uncultivated sites, both from the ground and the vegetation. No special methods were used. Ants were located visually while walking randomly, and they were collected in alcohol tubes using a camel hair brush/forceps. Twigs were broken to collect nesting arboreal ants.

Collection localities

Ants were collected from 14 islands in the Andamans and three in the Nicobars (Table 1) at various times between 1989 and 1998. It was, however, on the island of South Andaman that over 80% of the time was spent collecting ants. Only 13% of the time was spent in the Nicobars, with most of this time (over 98%) spent on the island of Great Nicobar. The islands from where ants had been collected earlier but which the authors were unable to visit were South Brother and North Reef in the Andamans and Pulo Milu in

the Nicobars (Mayr 1865; Forel 1903; Mukherji and Ribeiro 1925; Chhotani and Maiti 1977).

The ants collected by the first and third authors were identified by Musthak Ali, the second author.

Results and Discussion

This study raises the total species of ants known from these islands to 125, from a previous total of 59. Only 37 of the 59 previously reported species of ants were collected (Table 2). The 125 species of ants listed here belong to 41 genera. While 10 genera (*Anochetus*, *Platythyrea*, *Cerapachys*, *Bothriomyrmex*, *Dolichoderus*, *Gnamptogenys*, *Hyoponera*, *Podomyrma*, *Pristomyrmex*, and *Technomyrmex*) were recorded for the first time from these islands, seven genera (*Myopopone*, *Paratopula*, *Vollenhovia*,

Liomyrmex, *Metapone*, *Acropyga*, and *Echinopla*) found in the earlier literature were not encountered. Of these seven genera, all but *Paratopula* and *Liomyrmex* were collected from the Nicobar islands, where this survey was limited.

The fact that some Formicidae previously recorded from these islands were not collected does not imply that they have gone extinct. It only denotes that more intensive collections have to be made when most, if not all, of these species are likely to be found.

Echinopla, *Myopopone*, *Odontoponera*, and *Philidris* are genera with Malesian (or Indoaustralian) + Australasian distribution that find their western limit on these islands. Not only was *Echinopla* not found on these islands, but there is also some doubt about the occurrence of the genus on these islands. The lone species of the genus was reported from

Table 1. Islands of the Andamans and Nicobars from where ants were collected.

Island Group	Islands surveyed	Total geographical area ¹ (km ²)	Area under forest I (km ²)	Collection effort (person hours)
ANDAMAN ISLANDS				
Great Andaman	Landfall	29.48	29.48	0.25
	*East	3.55	3.55	0.25
	*North Andaman	1375.99	1247.51	45
	*Middle Andaman	1535.5	1348.37	45
	*Baratang	297.6	283.2	1
	*South Andaman	1348.2	883.4	2550
	Smith	24.7	15.79	0.25
	Kyd	8	8	0.25
	North Reef	348	348	-
	*Ross	0.78	Nil	0.25
*Chatham	0.16	Nil	0.25	
Ritchie's Archipelago	*Havelock	113.93	95.6	20
	*Neil	189.6	6.74	20
Labyrinth Is.	Alexandra	4.97	4.97	0.5
Little Andaman	*Little Andaman	731.57	706.49	60
	South Brother	1.24	1.24	-
	Narcondam	7.38	7.38	6
NICOBAR ISLANDS				
North Nicobar	*Car Nicobar	126.91	NA	25
Middle Nicobar	*Nancowry	66.82	40	1
South Nicobar	*Pulo Milo	1.29	NA	-
	*Kondul	4.66	NA	0.25
	*Great Nicobar	1044.54	960.4	405

* indicates human presence on island; NA= not available; '-' indicates islands not visited by the authors of this paper; ¹ Source: Anonymous, 1998

Table 2. Preliminary list of ants of the Andaman and Nicobar islands

Sl. No.	Species	Subfamily	Old (o), Present(*)	Islands of occurrence
			Present and old (**) records	
1	<i>Acropyga acutiventris</i> Roger, 1862	Formicinae	o (1)	PM
2	<i>Aenictus binghami</i> Forel, 1901	Aenictinae	*	NA
3	<i>Aenictus fergusonii</i> Forel, 1901	Aenictinae	** (1)	SA;GN
4	<i>Aenictus ?gracilis</i> Emery, 1893	Aenictinae	*	SA
5	<i>Aenictus</i> sp.	Aenictinae	*	SA
6	<i>Aenictus</i> sp. nr. <i>laeviceps</i> (Smith, F. 1857)	Aenictinae	*	SA
7	<i>Anochetus graeffei</i> Mayr, 1870	Ponerinae	*	GN
8	<i>Anochetus</i> sp. nr. <i>yerburyi</i> Forel, 1900	Ponerinae	*	SA
9	<i>Anoplolepis gracilipes</i> (Smith, F. 1857)	Formicinae	** (1,2,3)	SA; C; S; LA; Ny;
10	<i>Aphaenogaster beccarii</i> Emery, 1887	Myrmicinae	o (2)	SA
11	<i>Aphaenogaster feae</i> Emery, 1889	Myrmicinae	*	NA; SA
12	<i>Aphaenogaster feae nicobarensis</i> Forel,	Myrmicinae	** (1)	GN
13	<i>Aphaenogaster longiceps</i> (Smith, F. 1858)	Myrmicinae	*	GN
14	<i>Aphaenogaster</i> sp.	Myrmicinae	*	GN
15	<i>Bothriomyrmex wroughtonii dalyi</i> Forel, 1895	Dolichoderinae	*	SA
16	<i>Camponotus badius</i> (Smith, F. 1857)	Formicinae	*	SA
17	<i>Camponotus compressus</i> (Fabricius, 1787)	Formicinae	** (3)	SA
18	<i>Camponotus invidus</i> Forel, 1892	Formicinae	** (3)	SA; NA
19	<i>Camponotus variegatus infuscus</i> (Forel, 1892)	Formicinae	*	N
20	<i>Camponotus irritans</i> (Smith, F. 1857)	Formicinae	** (1)	SA; CN ; GN
21	<i>Camponotus leonardi</i> Emery, 1889	Formicinae	** (1)	C; SA
22	<i>Camponotus mitis</i> Smith, F. 1858	Formicinae	o (1)	SA
23	<i>Camponotus nicobarensis</i> Mayr, 1865	Formicinae	** (1,3)	SA; NA; MA; CN
24	<i>Camponotus oblongus</i> (Smith, F. 1858)	Formicinae	o (3)	SA
25	<i>Camponotus parius</i> Emery, 1889	Formicinae	*	E; SA; NA
26	<i>Camponotus vitreus</i> (Smith, F. 1860)	Formicinae	** (1)	C; Ny; GN
27	<i>Camponotus</i> sp. nr. <i>oblongus</i> (Smith, F. 1858)	Formicinae	*	SA
28	<i>Camponotus</i> sp. 1	Formicinae	*	SA; N; GN
29	<i>Camponotus</i> sp. 2	Formicinae	*	NA; S; GN
30	<i>Cardiocondyla nuda</i> (Mayr, 1866)	Myrmicinae	** (1)	KI
31	<i>Cataulacus granulatus</i> (Latreille, 1802)	Myrmicinae	** (1)	SA
32	<i>Cataulacus muticus</i> Emery, 1889	Myrmicinae	*	GN
33	<i>Cataulacus simoni</i> Emery, 1893	Myrmicinae	o (1,3)	SA;C
34	<i>Cataulacus taprobanae</i> Smith, F. 1853	Myrmicinae	*	L; LA
35	<i>Cerapachys</i> sp.	Cerapachyinae	*	GN
36	<i>Crematogaster dohrni artifex</i> Mayr, 1879	Myrmicinae	*	SA; NA
37	<i>Crematogaster rogenhoferi</i> Mayr, 1879	Myrmicinae	** (1)	C; SA; SB
38	<i>Crematogaster wroughtonii</i> Forel, 1902	Myrmicinae	o (2)	SA
39	<i>Crematogaster</i> sp. nr. <i>anthracina</i> Smith, F.	Myrmicinae	*	GN
40	<i>Crematogaster</i> sp.	Myrmicinae	** (1)	SA; LA; GN
41	<i>Diacamma rugosum</i> (Le Guillou, 1842)	Ponerinae	** (1,2,3)	SA; NA; MA; H; Kd;
42	<i>Dolichoderus taprobanae</i> Smith, F. 1858	Dolichoderinae	*	LA; N
43	<i>Dolichoderus</i> sp.	Dolichoderinae	*	SA; N
44	<i>Echinopla lineata senilis</i> Mayr, 1862	Formicinae	o (1)	GN
45	<i>Gnamptogenys coxalis</i> (Roger, 1860)	Ponerinae	*	GN
46	? <i>Hypoponera</i> sp.	Ponerinae	*	GN
47	<i>Leptogenys diminuta</i> (Smith, F. 1857)	Ponerinae	** (3)	SA; NA
48	<i>Leptogenys minchinii</i> Forel, 1900	Ponerinae	*	SA
49	<i>Leptogenys</i> sp. (<i>diminuta</i> group)	Ponerinae	*	SA
50	<i>Liomyrmex taylori</i> Tiwari & Jonathan, 1986	Myrmicinae	o (4)	SA
51	<i>Metapone nicobarensis</i> Tiwari & Jonathan,	Myrmicinae	o (5)	GN
52	<i>Monomorium destructor</i> (Jerdon, 1851)	Myrmicinae	** (1,7)	C; Kd; E; SA
53	<i>Monomorium floricola</i> (Jerdon, 1851)	Myrmicinae	** (7)	SA; GN, N
54	<i>Monomorium orientale</i> Mayr, 1879	Myrmicinae	*	SA
55	<i>Monomorium pharaonis</i> (Linnaeus, 1758)	Myrmicinae	*	SA
56	<i>Monomorium sagei</i> Forel 1902	Myrmicinae	o (2)	SA
57	<i>Monomorium</i> sp.	Myrmicinae	*	LA
58	<i>Myopopone castanea</i> (Smith, F. 860)	Amblyoponinae	o (1)	GN
59	<i>Odontomachus simillimus</i> Smith, F. 1858	Ponerinae	*	GN
60	<i>Odontomachus</i> sp.	Ponerinae	** (1)	GN

SI. No.	Species	Subfamily	Old (o), Present(*)	Islands of occurrence
			Present and old (** records)	
61	<i>Odontoponera transversa</i> (Smith, F. 1857)	Ponerinae	** (2)	L; SA; H; A; NA; S; MA
62	<i>Oecophylla smaragdina</i> (Fabricius, 1775)	Formicinae	** (1,2,3)	SA; B; R; LA
63	<i>Oligomyrmex similis</i> (Mayr, 1862)	Myrmicinae	o (1)	CN
64	<i>Oligomyrmex</i> sp.	Myrmicinae	*	GN
65	<i>Pachycondyla luteipes</i> (Mayr, 1862)	Ponerinae	o (1)	PM
66	<i>Pachycondyla rufipes</i> (Jerdon, 1851)	Ponerinae	** (2)	SA; NA, LA
67	<i>Pachycondyla</i> sp. (<i>luteipes</i> group)	Ponerinae	*	SA; GN
68	<i>Paratopula andamanensis</i> (Forel, 1903)	Myrmicinae	o (1)	LA
69	<i>Paratrechina bourbonica</i> (Forel, 1886)	Formicinae	o (1)	GN
70	<i>Paratrechina indica</i> (Forel, 1894)	Formicinae	** (1,3)	L; SA; GN
71	<i>Paratrechina longicornis</i> (Latreille, 1802)	Formicinae	** (1,2)	C; SA; NA; Kd; GN; N
72	<i>Paratrechina yerburyi</i> (Forel, 1894)	Formicinae	*	GN
73	<i>Paratrechina</i> sp. nr. <i>indica</i> (Forel, 1894)	Formicinae	*	L; GN
74	<i>Paratrechina</i> sp.	Formicinae	*	GN
75	<i>Pheidole capellinii</i> Emery, 1887	Myrmicinae	*	SA
76	<i>Pheidole longipes</i> (Latreille, 1802)	Myrmicinae	*	GN
77	<i>Pheidole megacephala</i> (Fabricius, 1793)	Myrmicinae	** (1)	KI; GN
78	<i>Pheidole noda</i> Smith, F. 1874	Myrmicinae	o (1)	LA
79	<i>Pheidole striativentris</i> Mayr, 1879	Myrmicinae	o (1)	C
80	<i>Pheidole watsoni</i> Forel, 1902	Myrmicinae	*	GN
81	<i>Pheidole</i> sp. nr. <i>grayi</i> Forel, 1902	Myrmicinae	*	SA
82	<i>Pheidole</i> sp. nr. <i>megacephala</i> (Fabricius, 1793)	Myrmicinae	*	GN
83	<i>Pheidole</i> sp.1	Myrmicinae	*	SA; N; GN
84	<i>Pheidole</i> sp.2	Myrmicinae	*	N
85	<i>Pheidologeton affinis</i> (Jerdon, 1851)	Myrmicinae	** (1)	GN
86	<i>Pheidologeton</i> sp.	Myrmicinae	o (2)	SA
87	<i>Philidris myrmecodiae andamanensis</i> (Forel, 1903)	Dolichoderinae	** (1)	SA; LA; Kd; C
88	<i>Philidris ? laevigata</i> (Emery, 1895)	Dolichoderinae	*	SA
89	<i>Platythyrea parallela</i> (Smith, F.1859)	Ponerinae	*	SA
90	<i>Podomyrma</i> sp.	Myrmicinae	*	SA
91	<i>Polyrhachis armata</i> (Le Guillou, 1842)	Formicinae	** (3)	SA; NA; S; MA; LA; B
92	<i>Polyrhachis bicolor</i> Smith, F.1858	Formicinae	** (2)	SA; NA; S; LA
93	<i>Polyrhachis bihamata</i> (Drury, 1773)	Formicinae	** (1)	SA; LA
94	<i>Polyrhachis dives</i> Smith, F.1857	Formicinae	*	N; LA
95	<i>Polyrhachis hector</i> Smith, F. 1857	Formicinae	*	SA
96	<i>Polyrhachis illaudata</i> Walker,1859	Formicinae	** (3)	SA; NA; S; MA; N; LA
97	<i>Polyrhachis laevissima</i> Smith, F. 1858	Formicinae	** (1,2,3)	SA; Kd; LA; N
98	<i>Polyrhachis rastellata</i> (Latreille, 1802)	Formicinae	*	SA; NA; LA
99	<i>Polyrhachis tibialis</i> Smith, F. 1858	Formicinae	*	SA; NA; S
100	<i>Polyrhachis thrinax</i> Roger, 1863	Formicinae	*	SA; MA; LA
101	<i>Polyrhachis</i> sp. nr. <i>fortis</i> Emery, 1893	Formicinae	*	SA
102	<i>Polyrhachis</i> sp. nr. <i>hector</i> Smith, F. 1857	Formicinae	*	SA; S
103	<i>Polyrhachis</i> sp.nr. <i>hippomanes ceylonensis</i> Emery,1893	Formicinae	*	SA
104	<i>Polyrhachis</i> sp.1	Formicinae	*	SA; NA; S; GN
105	<i>Polyrhachis</i> sp.2	Formicinae	*	GN
106	<i>Pristomyrmex</i> sp.	Myrmicinae	*	GN
107	<i>Solenopsis geminata</i> (Fabricius, 1804)	Myrmicinae	** (1,3)	SA; CN;GN;LA
108	<i>Tapinoma andamanense</i> Forel, 1903	Dolichoderinae	** (1)	N; C
109	<i>Tapinoma melanocephalum</i> (Fabricius, 1793)	Dolichoderinae	** (2)	SA; Ny; GN; N
110	<i>Tapinoma</i> sp.	Dolichoderinae	*	SA; GN
111	<i>Technomyrmex albipes</i> (Smith, F. 1861)	Dolichoderinae	*	NI; N
112	<i>Tetramorium bicarinatum</i> (Nylander, 1846)	Myrmicinae	** (2,3,6)	GN; SA; SS
113	<i>Tetramorium lanuginosum</i> Mayr, 1870	Myrmicinae	*	GN
114	<i>Tetramorium pacificum</i> (Mayr, 1870)	Myrmicinae	*	N
115	<i>Tetramorium</i> sp. nr. <i>curvispinosum</i> Mayr,	Myrmicinae	*	N

Sl. No.	Species	Subfamily	Old (o), Present(*)	Islands of occurrence
			Present and old (**) records	
116	<i>Tetramorium indicum</i> (Forel, 1913)	Myrmicinae	o (6)	NR
117	<i>Tetramorium</i> sp. 1	Myrmicinae	*	GN
118	<i>Tetramorium</i> sp. 2	Myrmicinae	** (2)	R ; SA; GN
119	<i>Tetraponera aitkenii</i> (Forel, 1902)	Pseudomyrmecinae	*	SA
120	<i>Tetraponera allaborans</i> (Walker, 1859)	Pseudomyrmecinae	*	SA
121	<i>Tetraponera nicobarensis</i> (Forel, 1903)	Pseudomyrmecinae	o (1)	GN
122	<i>Tetraponera rufonigra testaceonigra</i> (Forel, 1903)	Pseudomyrmecinae	** (1)	SA; NA; MA; CN ;
123	<i>Tetraponera siggi nebulosa</i> (Forel, 1903)	Pseudomyrmecinae	o (1)	GN
124	<i>Tetraponera</i> sp.	Pseudomyrmecinae	*	SA; GN
125	<i>Vollenhovia oblonga laevithorax</i> Emery, 1889	Myrmicinae	o (1)	GN

Islands from where ants were collected in the past are in bold letters. Validity of names based on Bolton, 1995b and Agosti, D., and N. F. Johnson. Editors. 2005. Antbase. www.antbase.org, version (05/2005).

Andaman islands: A = Alexandra; B = Baratang; C = Chatham; E = East, H = Havelock; Kd = Kyd; L = Landfall; LA = Little Andaman; MA = Middle Andaman; NA = North Andaman; N = Narcondam; NI = Neil; NR = North Reef; R = Ross; S = Smith; SA = South Andaman; SB = South Brother; SS = Stewart Sound

Nicobar islands: CN = Car Nicobar; GN = Great Nicobar; KI = Kondul; Ny = Nancowry; PM = Pulo Milo

Numbers in parentheses indicate reports by the following: 1-Forel, 1903; 2-Mukherji and Ribeiro, 1925; 3-Chhotani and Maiti, 1977; 4-Tiwari and Jonathan, 1986a; 5-Tiwari and Jonathan, 1986b; 6-Bolton, 1977; 7-Bolton, 1987

the southern Nicobar islands as *Echinopla senilis* by Mayr in 1862. This species was later considered a variety of *E. lineata* Mayr by Emery (1896), who, however, did not furnish any reasons justifying the change. In all probability, no one has ever seen this species after 1862 (Baroni Urbani 1997 *In litt.*).

Cerapachys, *Tapinoma*, *Acropyga*, *Camponotus*, *Paratrechina*, *Crematogaster*, *Monomorium*, *Pheidole*, *Solenopsis*, and *Tetramorium* are 10 of the 15 most widely distributed genera (i.e., those that are found in all the 8 zoogeographic regions of the world) (Bolton 1995) that are found on these islands.

Hydnophytum formicarum Jack (Rubiaceae) and *Dischidia major* (Vahl.) Merr. (Asclepiadaceae) are the two myrmecophytes that are found associated with species of *Philidris* on these islands. Also, *Cataulacus* sp. was found nesting in *Dischidia* on the Andaman Islands.

Anoplolepis gracilipes and *Pheidole megacephala* are among the world's worst invasive ant species (Global Invasive Species

Database, <http://www.issg.org/database>). Both were found on these islands in the early twentieth century (Forel 1903). It is important to note that Forel's study appeared about fifty years after the British had occupied these islands. *A. gracilipes* is widely distributed in the Afro-Tropical region, and it is a well-known tramp species. It is known to have caused extensive environmental damage in island ecosystems especially in Hawaii, Seychelles, Zanzibar, and Christmas Island. Along with *P. megacephala*, these species pose a serious threat to the native invertebrate fauna of these islands.

It is interesting to note in this context that 10 species of ants were intercepted in 18 shiploads of timber transported between 2006 and 2009 from Malaysia and Myanmar to ports along the west coast of India. These were *Anoplolepis gracilipes*, *Pheidole* sp., *Diacamma rugosum*, *Camponotus compressus*, *Polyrhachis rastellata*, *Rhoptomyrmex wroughtoni*, *Aphaenogaster* sp., *Paratrechina longicornis*, and *Pheidologiton diversus* (Manickam, personal collection).

Considering that the Andaman islands have three times more area than the Nicobars, they should also be expected to harbour a richer biota than the latter island group. In fact, the data from this survey seem to reflect this trend. However, since the collection efforts were grossly uneven between the two island groups, with greater effort expended on the Andamans, this cannot be taken as a true reflection of greater species richness in the Andamans than in the Nicobars. To arrive at a truly representative picture of the relative diversities of Formicidae between the two island groups, more intensive collections will have to be made.

With transoceanic dispersal capabilities only marginally poorer than bats and better than many insect orders (like Trichoptera, Isoptera, etc.), ants are among the most successful early colonists of islands (Zimmerman 1948). This, and the fact that no studies have so far been focused exclusively on the ants of these islands, makes it almost certain that many more genera/species await discovery, particularly in the leaf litter and arboreal habitats. It is, therefore, imperative that studies are initiated on the ants of these islands before habitat destruction leads to the elimination of many species even before they are discovered.

Voucher specimens have been deposited at the University of Agricultural Sciences, GKVK, Bangalore and the Department of Entomology, Indian Agricultural Research Institute, New Delhi, India.

Acknowledgement

We are grateful to Dr. A.K. Bandyopadhyay, who, in his capacity as Director, C.A.R.I., and Port Blair, always supported us in our studies on the insect diversity of these islands. Dr.

Cesare Baroni Urbani and Dr. B. Kiauta very kindly went through an earlier version of this paper and helped us improve it considerably.

We thank Mr. P. Manickam for bringing the ant specimens intercepted by the Department of Quarantine along the West coast of India for identification to Musthak Ali. We acknowledge the assistance of Bikas Mondal in field and laboratory work and the Department of Forests, Andaman and Nicobar islands for having granted us permission (*vide* order No. CWLW/WL/47/1294 dated 4.2.1993) to collect insects for study from the National Parks and Wildlife Sanctuaries of these islands.

References

-
- Anonymous. 1986. *An Integrated Environmentally Sound Development Strategy for the Andaman and Nicobar Islands*. Planning commission, Govt. of India.
- Anonymous. 1994. *Basic Statistics: Andaman and Nicobar islands*. Statistical Bureau, Andaman and Nicobar Administration.
- Anonymous. 1998. *Forest statistics 1996-1997*. Department of Environment and Forests, Andaman and Nicobar Administration.
- Bingham CT. 1903. *The Fauna of British India, including Ceylon and Burma*. Hymenoptera, Vol. II. Ants and Cuckoo-wasps. Taylor and Francis.
- Bolton B. 1977. The ant tribe Tetramoriini. The genus *Tetramorium* Mayr in the Oriental ant Indo-Australian regions, and in Australia. *Bulletin of the British Museum (Natural History)(Entomology)* 36: 67-151.

- Bolton B. 1987. A review of the *Solenopsis* genus-group and revision of Afrotropical *Monomorium* Mayr. *Bulletin of the British Museum (Natural History)(Entomology)* 54: 263-452.
- Bolton B. 1995a. A taxonomic and zoogeographical census of the extant ant taxa (Hymenoptera: Formicidae). *Journal of Natural History* 29: 1037-1056.
- Bolton B. 1995b. *A New General Catalogue of the Ants of the World*. Harvard University Press.
- Champion SG, Seth SK. 1968. *A Revised Survey of the Forest Types of India*. Govt. of India.
- Chhotani OB, Maiti PK. 1977. Contribution to the knowledge of Formicidae of the Andaman islands. *Newsletter of Zoological Survey of India* 3(1): 17 - 20.
- Curray JR. 1989. The Sunda arc: A model of oblique plate convergence. *Netherlands Journal of Sea Research* 24(2/3): 131 - 140.
- Curray JR, Moore DG, Lawver LA, Emmel FJ, Raitt RW, Henry M, Kiechefer R. 1979. Tectonics of the Andaman Sea and Burma. In: Watkins J, Montadert L, Dickerson PW, editors. Geological and geophysical investigations of continental slopes and rises. *American Association of Petrol and Geology Memoirs* 29: 189-198.
- Emery C. 1896. *Memorie della Accademia delle Scienze dell'Istituto di Bologna* 5: 780.
- Emery C. 1911. Hymenoptera, Fam. Formicidae, Subfam. Ponerinae. In: Wystman P, editor. *Genera Insectorum*, fasc. 118. Brussels.
- Emery C. 1912. Hymenoptera, Fam. Formicidae, Subfam. Dolichoderinae. In: Wystman P, editor. *Genera Insectorum*, fasc. 137. Brussels.
- Emery C. 1921. Hymenoptera, Fam. Formicidae, Subfam. Myrmicinae. In: Wystman P, editor. *Genera Insectorum*, fasc. 174A. Brussels.
- Emery C. 1922. Hymenoptera, Fam. Formicidae, Subfam. Myrmicinae. In: Wystman P, editor. *Genera Insectorum*, fasc. 174 B, C. Brussels.
- Forel A. 1903. Les fourmis des iles Andamans et Nicobars: Rapports de cette faune avec ses voisines. *Revue Suisse de Zoologie* 11: 399-411.
- Global Invasive Species Database. Available from: <http://www.issg.org/database>
- Hamilton W. 1979. Tectonics of the Indonesian region. *United States Geological Survey Professional Paper* 1078: 64-67.
- Mayr GL. 1865. *Formicides der Reise der Oestereischen Fregatte Novara um die Erde*. p. 23, Taff. 4 Wien.
- McVean DN. 1976. *Report on Land Use in the Andaman and Nicobar Islands*. Morges.
- Moore GF, Curray JR, Moore DG. 1980. Variations in geologic structure along the Sunda fore arc, northeastern Indian Ocean, Repr. The tectonic and geological evolution of southeast Asian seas and islands. *Geophysical Monographs* 23: 145-160.
- Mukherji D, Ribeiro S. 1925. On a collection of ants (Formicidae) from the Andaman islands. *Records of Indian Museum* 27: 205-209.

Nassig WA, Lampe REJ, Kager F. 1996. Saturniidae of Sumatra (Lepidoptera). *Heterocera Sumatrana* 10: 1-174.

Ripley SD, Beehler BM. 1989. Ornithogeographic affinities of the Andaman and Nicobar islands. *Journal of Biogeography* 16: 323-332.

Tiwari RN, Jonathan JK. 1986a. A new species of *Liomyrmex* Mayr from Andaman islands (Hymenoptera: Formicidae). *Records of Zoological Survey of India* 83(1&2): 87-90.

Tiwari RN, Jonathan JK. 1986b. A new species of *Metapone* Forel from Nicobar islands (Hymenoptera: Formicidae: Myrmicinae). *Records of Zoological Survey of India* 83 (1&2): 149-153.

Zimmerman EC. 1948. *Insects of Hawaii*, Vol. 1: Introduction. University of Hawaii Press.