

# Effects of Intra- and Interpatch Host Density on Egg Parasitism by Three Species of Trichogramma

Authors: Grieshop, Matthew J., Flinn, Paul W., and Nechols, James R.

Source: Journal of Insect Science, 10(99): 1-14

Published By: Entomological Society of America

URL: https://doi.org/10.1673/031.010.9901



# Effects of intra- and interpatch host density on egg parasitism by three species of *Trichogramma*

Matthew J. Grieshop<sup>1,3a</sup>, Paul W. Flinn<sup>2b</sup>, and James R. Nechols<sup>1c</sup>

<sup>1</sup>Kansas State University, Department of Entomology, 123 West Waters Hall, Manhattan, KS66506-4004, USA <sup>2</sup>USDA-ARS, Grain Marketing and Production Research Center, 1515 College Avenue, Manhattan Kansas 66502, USA

<sup>3</sup>Current address: Michigan State University, Department of Entomology, 205 Center for Integrated Plant Systems, East Lansing, MI 48824

#### **Abstract**

Host-foraging responses to different intra- and interpatch densities were used to assess three Trichogramma spp. (Hymenoptera: Trichogrammatidae) Trichogramma deion Pinto and Oatman, T. ostriniae Pang and Chen, and T. pretiosum Riley — as potential biological control agents for the Indian meal moth, *Plodia interpunctella* Hübner (Lepidoptera: Pyralidae). Single naïve females were allowed 6 h to forage in Plexiglas arenas with four different spatial arrangements of host eggs, nine single-egg patches), nine four-egg patches, 36 single-egg patches, and 36 four-egg patches. No significant differences were found among species in the number of patches parasitized. As expected, all three species parasitized the most eggs in the 36 four-egg patch treatment and the least in the nine single-egg patch treatment. T. deion parasitized significantly more eggs than T. pretiosum on the nine four-egg patches. T. ostriniae parasitized significantly more patches when intrapatch density was greater, regardless of interpatch density. In contrast, T. deion only parasitized more patches at the greater intrapatch density when the interpatch density was low. Patch density had no effect on T. pretiosum. The spatial pattern of parasitism was more aggregated for T. deion and T. ostriniae in the 36 four-egg patches treatment compared to the 36 single-egg patches treatment. Therefore, intrapatch density was more important than interpatch density for T. ostriniae, and potentially for T. deion, but not for T. pretiosum. T. deion may be the best candidate for augmentative biological control because it parasitized either slightly or significantly more eggs than the other two species in all four treatments. Furthermore, the pattern of parasitism by T. deion in the 36 four-egg patches treatment was the most aggregated among the three species, suggesting a more thorough searching pattern. In contrast, T. pretiosum had the least aggregated pattern of parasitism and therefore may have used a more random foraging pattern.

**Keywords:** biological control, foraging behavior, *Plodia interpunctella*, spatial scale, stored products, *Trichogramma deion*, *Trichogramma ostriniae*, *Trichogramma pretiosum* 

Abbreviations: nAI, normalized aggregation index

Correspondence: a grieshop@msu.edu, b paul.plinn@ars.usda.gov, c jnechols@ksu.edu

Associate Editor: Robert Jeanne was editor of this paper.

Received: 21 May 2008, Accepted: 20 July 2009

Copyright: This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits

unrestricted use, provided that the paper is properly attributed.

ISSN: 1536-2442 | Vol. 10, Number 99

#### Cite this paper as:

Grieshop MJ, Flinn PW, Nechols JR. 2010. Effects of intra- and interpatch host density on egg parasitism by three species of *Trichogramma*. *Journal of Insect Science* 10:99 available online: insectscience.org/10.99

# Introduction

The Indian meal moth, *Plodia interpunctella* (Lepidoptera: Pyralidae) (Hübner), is a cosmopolitan pest of stored products found in bulk, processing, warehouse, and retail systems (Cox and Bell 1991). The range of products attacked by this pest includes: raw and finished cereals, nuts, dried fruit and vegetables, pulses, and garlic (Perez-Mendoza and Aguilera-Pena 2004). Moths cause damage by infesting both unprocessed, bulk well products as as finished, retail commodities. In bulk products, they are generally considered to be a nuisance pest. However, infestations in retail stores and warehouses result in significant damage, because the presence of a single larva or adult may result in the total loss of the infested product, as well as potential customer complaints or even lawsuits. The traditional management strategy for *P. interpunctella* has been the use of chemical insecticides, including: fog, fumigant, and grain protectants (Cox and Bell 1991). However, recent legislation has reduced the availability of chemical insecticides for use on stored products (Arthur and Rogers 2003).

Trichogramma (Hymenoptera: Trichogrammatidae) egg parasitoids have recently been studied as augmentative biological control agents for P. interpunctella in stored product systems, including: bulk peanut storage (Brower 1988), bulk wheat storage (Schöller et al. 1996), bakeries (Prozell and Schöller 1998; Steidle et al. 2001), as well as in warehouses and retail stores (Schöller et al. 2006). Trichogramma spp. present some unique advantages that make them well-suited for use in the high visibility/low economic threshold systems characteristic of finished stored products. These include: causing mortality to the egg stage prior to the damaging larval stage, commercial availability, and perhaps most importantly, tiny body size (< 0.5 mm in length), making them unlikely to be noticed by consumers.

Selection of an appropriate species and/or strain of biological control agent is a critically important part of any augmentative biological control program (Smith 1996; Van Driesche and Bellows 1996), and has been discussed specifically in regards choosing trichogrammatids to manage stored product moths (Schöller et al. 1994). The fact that P. interpunctella, unlike other pyralid moths such as Ostrinia nubilalis, lays its eggs both singly and in small clusters, and does so either directly on exposed products (Arbogast and Mullen 1978) or adjacent to compromised packages in response to food odors (Mullen 1994), results in a heterogeneous distribution of small (< 10) localized patches of host eggs. In general, variation in intra- and interpatch host density is known to affect parasitoid foraging behavior and success (Hassell 1982). In studies of *Trichogramma* spp., increased numbers of host eggs within patches have been shown to have a positive effect on host patch location (Morrison et al. 1980; Kfir 1983; Reznik and Umarova 1991; McCravy and Berisford 1998; Barnay et al. 1999) and foraging activity (Nordlund et al. 1981).

Therefore, the effects of four intra- and interpatch densities of *P. interpunctella* eggs were examined on the host-foraging success of individual naïve females of three *Trichogramma* species. The experimental objective was to select the most suitable species for use in an augmentative biological control program for this pest in retail stores and warehouses. Experimental assessments included measurement of overall parasitism,

as well as a simple assessment of the relative aggregation of parasitism between two intrapatch host densities. The latter assessment was perceived as important relative to the management of P. interpunctella due to this pest's tendency to lay single or small clusters of eggs adjacent to compromised food packages or product spills (Arbograst and Mullen 1978; Mullen 1994), which may result in a concentration of eggs within the vicinity of such host resources. Thus, a more aggregated pattern of parasitism is likely to be desirable, as it would lead to a more thorough exploitation of areas likely to contain hosts. This experiment was performed complement additional studies evaluating the foraging efficiency of the same three *Trichogramma* spp. in response to smaller scale habitat complexity in the form of small quantities of millet or flour (Grieshop et al. 2008), as well as their foraging response on retail shelving in the presence or absence of packaged goods (Grieshop et al. 2007b).

The three commercially available species selected for comparison were: T. deion (Pinto and Oatman), T. ostriniae Pang and Chen, and T. pretiosum (Riley). The commercial strains of *T. deion* and *T. pretiosum* used in this study were selected based on host preference tests performed by Schöller and Fields (2002). Their study demonstrated that these two species readily parasitized the eggs of P. interpunctella under both choice and nochoice conditions, while two strains of T. minutum Riley, and individual strains of T. sibericum Sorokina and Т. brassicae Bezdenko did not. Furthermore, T. pretiosum has been collected from wild stored product moth populations in peanut warehouses (Brower 1984). T. ostriniae has been extensively explored as an augmentative biological control agent for O. nubilalis in field and sweet corn (Wang et al., 1997, 1999; Hoffman et al. 2002; Wright et al. 2002), was shown capable of parasitizing the eggs of *P. interpunctella* in bulk wheat (Jeffery Gardener, personal communication), and has a history of being reared on a similar stored product pyralid *Ephestia kuhniella* (Smith 1996). Likewise all three *Trichogramma* strains used in this study had been reared on *E. kuhniella* for more than 100 generations (Synthia Penn, personal communication; Jeffery Gardener, personal communication).

# **Materials and Methods**

#### **Insects**

All insect species were maintained at the USDA-ARS Grain Marketing and Production Research Center in Manhattan, Kansas, USA, in a walk-in growth chamber set at  $26 \pm 1^{\circ}$  C,  $60 \pm 5\%$  RH, and a 16:8 light:dark photoperiod. Colonies of all three Trichogramma species were maintained on fresh (< 1 day old) eggs of E. kuehniella that had been sterilized by exposure to UV radiation. Unsterilized eggs of interpunctella were used in experiments. Both P. interpunctella and E. kuehniella were reared on a diet composed of cracked wheat, wheat shorts. honev. and glycerin (McGaughey and Beeman 1988). T. deion, T. pretiosum, and E. kuehniella starter cultures were provided by Beneficial Insectaries (www.insectary.com) in April 2003, March 2004, and February 2002, respectively. T. ostriniae were provided by Michael Hoffman's laboratory at Cornell University in November 2003. The P. interpunctella used in experiments had been in continuous culture at the Grain Marketing and Production Research Center for 10 or more years. Specimens of all species used in the experiment were deposited into the Kansas State University Museum of Entomological and Prairie Arthropod Research under voucher number 171.

Arrhenotokous strains of T. deion and T. ostriniae were used with observed sex ratios of 2:1 and 3:1 (female: male, respectively). Neither species was observed host feeding. The *T. pretiosum* strain was virtually thelytokous with an observed sex ratio of 24:1 (female: male), and was observed host Individual naïve, Trichogramma feeding. females were used in the trials. Females were less than 16 h old and had been allowed access to a 1:1 honey: water solution and mates for four hours prior to collection. Trichogramma females Individual collected using a natural hair paintbrush to drive individual females up a small strip of paper (0.25 cm wide x 4 cm long) that was then placed in an empty 8-dram shell vial.

# **Experimental design**

A three by four nested factorial design consisting of three species and four combinations of intra- and interpatch densities was used. The four treatments consisted of: nine single-egg patches (three by three grid of single-egg patches spaced 15 cm apart); nine four-egg patches (a three by three grid of four-egg patches spaced 15 cm apart); 36 single-egg patches (a six by six grid of single-egg patches spaced 7.5 cm apart); and 36 four-egg patches (a six by six grid of four-egg patches spaced 7.5 cm apart).

Experimental trials consisted of testing a single species at all four of the intra- and interpatch treatment combinations. There were 15 trials (replications) for each species with each density and a single species represented in each trial. Each trial lasted six hours and was performed in a walk-in environmental growth chamber at  $23 \pm 1^{\circ}$  C,  $45 \pm 5\%$  RH, and under constant light conditions provided by four banks of three 40 W soft white 122 cm fluorescent tubes. The environmental settings used in the experiments were based

upon temperature, humidity, and light conditions observed between April and November 2002, at four retail stores in Manhattan, KS, USA.

Treatment position on the lab bench was randomized for each experimental trial. Egg patches consisted of a 1 cm diameter cardstock disk with fresh (< 18 hr old) P. interpunctella eggs attached with a tragacanth paste, a nontoxic plant-derived glue (Merck, www.merck.com). In addition, 25-40 four-egg patches (100-160 eggs total) were kept in sealed Petri dishes for each trial to assess natural egg mortality unrelated to parasitoids. avoid cross-contamination To Trichogramma cultures, trials with T. deion were run between June and November 2003; trials with T. ostriniae were run between January and June 2004; and, for *T. pretiosum*, they were run between June and August 2004. Thus, only a single species of Trichogramma was maintained in the laboratory colony at any one time. Treatments were run in separate Plexiglas covered arenas measuring 50 x 50 x 10 cm (L x W x H) that were open on one of the 50 x 50 cm sides to form the floor of the arenas. Sentinel egg patch grids were laid out on a fresh piece of butcher paper, and a single naïve female *Trichogramma* was placed in the center of a randomly selected central quadrat. Immediately following release of the female, the Plexiglas cover was centered and placed over the grid of sentinel egg patches. An airtight seal between the edges of the arena and the paper was maintained using a strip of adhesive weather-stripping. Plexiglas covers were cleaned with distilled water prior to and immediately after experimental runs to minimize static charge.

# Data collection

Trichogramma females were removed from arenas after six hours. Egg patches were

numbered and placed in sealed Petri dishes, which were kept in a growth chamber set at 26 ± 1° C and 60 ± 7% RH. Host-egg condition was evaluated after seven days using a stereoscope (110-140x magnification). Eggs from individual egg patches were graded as hatched, parasitized, or dead; parasitized eggs were identified by their dark appearance and/or the presence of red pupal eyespots. The control egg patches were also judged at this time. All females used in the trials survived to the end of the experiment. Experimental runs with over 20% natural egg mortality were removed from the study, with additional runs made to bring the total number of replications per species to 15. A total of three, two, and two trials were repeated for T. deion, T. pretiosum, and T. ostriniae, respectively.

# Data analysis

Egg patches and egg parasitization data were analyzed using the mixed procedure (SAS 2001). Replicates with zero parasitism were removed from the analysis to reduce variability, resulting in 15, 13, and 14 replicates for T. deion, T. pretiosum, and T. ostriniae, respectively. The numbers of egg patches and eggs parasitized were analyzed in a three by four mixed model ANOVA with fixed factors consisting of species and the four treatments, and with species nested within experimental runs as a random factor. In addition, four three-way mixed model ANOVAs were run to test for differences among species within treatments, with species used as the fixed factor and species nested within experimental runs as the random factor. Three four-way ANOVAs, one for each species, were run for both the number of egg patches and eggs parasitized. Degrees of freedom were adjusted using the Kenward-Rodgers correction for mixed models with nesting, and multiple comparisons were run for all models using a Tukey-Kramer adjusted LSD (p = 0.05). Additional paired *t*-tests were run for each species comparing the number of egg patches parasitized for the nine single-egg patches treatment with the nine four-egg patches treatment, as well as the 36 single-egg patches treatment with the 36 four-egg patches treatment. Similar, paired *t*-tests were run for the number of eggs parasitized for the 36 single-egg patches treatment with the nine four-egg patches treatment. To assess whether there were differences in the number of non-responsive females among the three species and four treatments a two-way  $\chi^2$ test was performed.

In addition to patch use and parasitism, the landscape class metric "normalized aggregation index" (nAI) was calculated for egg patches parasitized by each species in the 36 single-egg patches treatment and the 36 four-egg patches treatment. The nAI equals the total length of edge (or perimeter) involving the corresponding class, given in number of cell surfaces, minus the minimum length of class edge (or perimeter) possible for a maximally aggregated class, also given in number of cell surfaces, which is achieved when the class is maximally clumped into a single, compact patch, divided by the maximum minus the minimum length of class edge (McGarigal et al. 2002). The nAI has a range from zero to one, with a value of zero indicating a maximally aggregated class and a value of one indicating a maximally dispersed class.

For the purpose of nAI calculation, patches were treated as 7.5 x 7.5 cm cells in a 6 x 6 landscape array with two class values denoting either a parasitized patch or a patch lacking parasitism. nAI was calculated using the Fragstats software package (McGarigal et al. 2002) using the equation:

$$nAI = \frac{e_i - \min e_i}{\max e_i - \min e_i}$$

Where  $e_i$  equals the actual total length of edge (or perimeter) of class i, min  $e_i$  equals the minimum possible total length of edge of class i based on the number of cells in class i, and max  $e_i$  equals the maximum possible total length of edge of class i based on the number of cells in class i. nAI values for the two treatments were analyzed for each of the three species using a paired t-test. Values were transformed using an angular transformation ( $\arcsin(y^{0.5})$ ) to stabilize variance.

# Results

# Egg patches parasitized

Significant differences were detected in the number of egg patches parasitized for treatment (F = 11.99; df = 3, 83.7; p =0.0001), but not for species or for the species by treatment interaction (F = 0.22; df = 2, 1; p = 0.8358 and F = 0.98; df = 6, 83.6; p = 0.4408, respectively) (Figure 1). The number of patches parasitized was numerically highest in the 36 single-egg patches for *T. deion* (10.11 patches) and in the 36 four-egg patches for both T. ostriniae and T. pretiosum (9.27 and 7.36 patches, respectively). The number of patches parasitized was lowest in the nine single-egg patches for all three species (Table 1). T. deion parasitized the most egg patches in the 36 four-egg patch treatment and the fewest in the two nine-patch treatments. Significant differences for *T. ostriniae* were detected between the 36 four-egg patches and all three of the other treatments. No significant differences were detected in the number of patches parasitized among treatments for *T. pretiosum* (Table 1).

Both *T. deion* and *T. ostriniae* parasitized significantly more patches in the nine fouregg patches treatment compared with the nine

single-egg patches treatment (Table 2). However, only *T. ostriniae* parasitized significantly more patches in the 36 four-egg patches treatment compared to the 36 single-egg patches treatment. No differences were observed for *T. pretiosum* regardless of patch size.

# Eggs parasitized

Significant differences were detected in the number of host eggs parasitized for treatment (F = 33.8; df = 3, 85; P = 0.0001) but not for species or species × treatment interaction (F = 0.9; df = 6, 85; p = 0.4990). However, species was nearly a significant factor (F = 2.79; df = 2, 83.7; p = 0.067) (Fig. 1). All three species parasitized the most eggs in the 36 four-egg patches treatment and the least number of eggs in the nine single-egg patches treatment. One-way ANOVA showed that T. deion parasitized significantly more eggs than T. pretiosum within the nine four-egg patches treatment (F = 3.67; df = 2, 21; p = 0.0429) (Figure 1).

With respect to patch effects on individual species, *T. deion* parasitized the most host eggs in the 36 four-egg patch treatments and the least in the nine single-egg patch treatment (Table 1). The trend in parasitism was similar for the other two *Trichogramma* species except that parasitism in the nine single-egg patch treatment was lowest, although not significantly so (Table 1).

To test whether the number of eggs parasitized was influenced by the total number of host eggs available versus the spatial pattern of host eggs (inter- and intra-patch configuration), parasitism for each species was compared for the two treatments containing 36 eggs (nine four-egg patches versus 36 single egg patches). No significant differences were detected for any of the

species, although the trends for T. deion and T. pretiosum were nearly significant (p = 0.0914 and p = 0.0738, respectively) with trends for more eggs parasitized in the nine four-egg patches treatment than in the 36 single-egg patches treatment (Table 2).


# **Number of non-responding females**

A similar pattern of non-responsive females was found among the three species with the highest number of non-responsive females found in the nine single-egg patches treatment and the lowest number found in the 36 four-

egg patches treatment ( $X^2 = 1.80$ , df = 3, p = 0.62). The exception was *T. ostriniae*, where the lowest number of non-responsive females was found in the 36 single-egg patches treatment (Table 3).

# Aggregation of parasitism

Aggregation of parasitism was significantly greater in the 36 four-egg patch treatment than in the 36 single-egg patch treatment for both T. deion (t = 4.04; df = 20; p < 0.001) and T. ostriniae (t = 2.34; df = 17; p < 0.03). While a significant difference was not detected for T. pretiosum, nAI was slightly lower for the 36 four-egg patch treatment (Figure 2).



**Figure 1.** Mean (+SEM) patches and eggs parasitized by species and treatment. Bars with different lowercase letters over them were found to be significantly different among species within the same treatment, as were whole treatments with different uppercase letters (Tukey-Kramer adjusted LSD with p < 0.05). High quality figures are available online.

# **Discussion**

Although patterns of parasitism varied considerably among species, no one species performed consistently better than the other two. The fact that all three species parasitized the most eggs where egg density was highest

(the 36 four-egg patches treatment), and the fewest where host density was lowest (the nine single-egg patches treatment) suggests that host-finding occurred in a density-dependent manner (Figure 1). The pattern of density-dependent parasitism is consistent with observations reported for *Trichogramma* 

**Table 1.** Mean ± SEM number of patches parasitized and number of eggs parasitized by *Trichogramma* species and treatment (eggs/patch).\*

Trichogramma species	Treatment	No. patches parasitized	No. eggs parasitized
T. deion	Nine single-eggs patches	2.33 ± 0.42c	2.33 ± 0.42c
	Nine four-egg patches	4.89 ± 0.70bc	15.67 ± 2.40b
	36 single-egg patches	10.11 ± 1.95a	10.11 ±
			1.95bc
	36 four-egg patches	8.69 ± 0.68ab	24.00 ± 1.81a
T. ostriniae	Nine single-egg patches	1.75 ± 0.48b	1.75 ± 0.48b
	Nine four-egg patches	5.29 ± 0.94b	10.86 ± 2.16b
	36 single-egg patches	6.55 ± 1.17b	6.55 ± 1.17b
	36 four-egg patches	9.27 ± 1.29a	22.36 ± 3.80a
T. pretiosum	Ninesingle-egg patches	2.50 ± 1.50a	2.50 ±
			1.50ab
	Nine four-egg patches	3.50 ± 0.53a	7.88 ± 1.59b
	36 single-egg patches	6.88 ± 2.37a	6.88 ±
			2.37ab
	36 four-egg patches	7.36 ± 1.61a	15.45 ± 3.34a

Data are shown as means ± SEM

\*Within a column and species, means with different letters are significantly different (Tukey-Kramer adjusted LSD P < 0.05).

**Table 2.** Paired *t*-test results for number of patches parasitized and number of eggs parasitized by trichogrammid species in test arenas (differences between means tested).

Species	Estimate ± SEM	Df	t value	P > t		
Patches parasitized: Nine single-egg patches – nine four-egg patches						
T. deion	-2.56 ± 0.93	13	-2.76	0.0163 *		
T. ostriniae	-3.54 ± 1.32	9	-2.67	0.0256 *		
T. pretiosum	-1.00 ± 1.27	8	-0.79	0.4522		
Patches parasitized: 36 single-egg patches – 36 four-egg patches						
T. deion	1.42 ± 1.80	20	0.79	0.4405		
T. ostriniae	-3.65 ± 1.49	19	-2.45	0.0244 *		
T. pretiosum	-1.23 ± 2.69	16	-0.45	0.6553		
Eggs parasitized: Nine four-egg patches – 36 single-egg patches						
T. deion	5.56 ± 3.09	16	1.8	0.0914		
T. ostriniae	4.31 ± 2.25	16	1.91	0.0738		
T. pretiosum	1.00 ± 2.85	14	0.35	0.7312		

 $*_{p} < 0.05$ 


 Table 3. Number of replicates by trichogrammid species and patch size treatment with non-responding females (out of 15).

	Number non-responding females			
Treatment	T. deion	T. ostriniae	T. pretiosum	
Nine single-egg patches	9	11	13	
Nine four-egg patches	6	8	7	
36 single-egg patches	6	4	7	
36 four-egg patches	2	5	5	

under laboratory (Kfir 1983) and field conditions (Wang and Ferro 1998). For *T. deion* and *T. ostriniae*, the number of host patches parasitized was highest when interpatch distance was lowest (36 patch treatments) (Table 1). Other authors have attributed increased host patch discovery to increased intrapatch density, both in the laboratory (Barnay et al. 1999), and in the field (Morrison et al. 1980; McCravey and Berisford 1998).

The three species responded differently in intra- and inter-patch comparisons. For *T. deion*, the number of patches parasitized was about two times greater in the nine four-egg patches treatment compared to the nine single-egg patches treatment. However, patches parasitized were similar in the 36 four-egg and single-egg patches treatments (Table 1). In contrast, *T. ostriniae* parasitized significantly more four-egg patches in both nine and 36 patch arrangements, and there were no intrapatch differences for *T. pretiosum* (Table 2).

One possible explanation for the differences in response to host densities among species is that the three species perceive "patch structure" differently. The greater spatial aggregation of patches parasitized in the 36 four-egg patches treatment compared with the nine single-egg patches treatment for both T. deion and T. ostriniae could be due to a foraging response to "meta" or super patches consisting of more than a single disk when four-eggs were present (Figure 2). In contrast, the lack of difference for T. pretiosum suggests that T. pretiosum may have a tendency to respond to the egg patches as individual patches irrespective of intrapatch density. The difference between T. pretiosum and the other two *Trichogramma* species may be the result of the latter two species exhibiting search an area-restricted response to the discovery of patches containing more hosts. For T. ostriniae this may be due to an evolutionary association with a host that lays dense egg masses versus sparsely distributed eggs.



**Figure 2.** Mean (+SEM) normalized aggregation index for one-egg-36 patch and four-egg-36 patch arenas. Values closer to zero indicate a higher level of aggregation. Pairs of bars with stars over them are significantly different (Paired t-test \* p < 0.05, \*\* p < 0.01). High quality figures are available online.

An alternative hypothesis that may explain increased patch discovery as interpatch density increases is that host patch discovery depends on the detection of a minimum level of kairomones associated with the host. Although untested in this study, the host eggs may have produced volatile compounds identifiable by Trichogramma and used as host-foraging cues since other studies have shown that Trichogramma responds to kairomones produced by or associated with host eggs, including scales from adult moths, sex pheromones, and compounds associated with egg metabolic processes (Jones et al. 1973; Morrison and Nordlund 1980; Nordlund 1981; Boo and Yang 2000). In addition, it has been hypothesized that *Trichogramma* spp. require a minimum amount of stimulus prior to engaging in normal foraging behavior (Resznik and Umarova 1991). Given that a greater number of non-responsive females were consistently found at the lowest egg density is in general agreement with this hypothesis (Table 3). Further evidence in support of this hypothesis is the nearly significant trend of a greater number of eggs parasitized by T. deion and T. ostriniae for the nine four-egg patches treatment compared to the 36 single-egg patches treatment (p =0.0914 and p = 0.0738, respectively) (Table 1). There may be different thresholds for density-dependent host-foraging responses in Trichogramma. For example, T. ostriniae exhibited greater parasitism only at the highest of the four host densities (Table 1).

The host densities and arrangements tested in this experiment were chosen because they provided an approximation of how P. *interpunctella* distributes eggs in warehouses and stores. A study by Arbogast and Mullen (1978) suggests that this distribution consists of single or small groups of eggs (< 10) per local group aggregated around a compromised

package or food product spill. An experiment performed by Grieshop et al. (2006) testing T. deion's ability to manage P. interpunctella in small arenas containing either bulk corn meal or cornmeal in compromised packaging clearly indicated that the parasitoid was largely ineffectual at reducing interpunctella populations on bulk product, but reduced it by more than 75% in packaged products. Thus, trichogrammatids that respond to low densities of hosts and thoroughly search in areas where low numbers of eggs are found (i.e., on packages or adjacent shelf space) would theoretically be better suited to manage this important stored product pest. In this case, both T. deion and T. ostriniae demonstrated a more aggregated search pattern at the highest host density, suggesting that they may perform better when foraging in finished stored product environments (Figure 2).

Two other experiments using the same three species, but conducted at different spatial scales, support the idea that T. deion is the best-suited species for biological control of P. interpunctella. The first was conducted at a smaller scale and examined the impact of microhabitat complexity on single females (Grieshop et al. 2008), while the second was conducted at a much larger scale and examined the foraging efficacy of massreleased parasitoids on retail shelving units (Grieshop et al. 2007). The failure of *T. deion* to outperform the other two species in this experiment may be due to either: 1) the difference in spatial scale among the present and previously published experiments, or 2) the homogeneous nature of the arenas used in this trial. Thorpe and Dively (1985) demonstrated that three species responded Trichogramma differently depending on the size of arena used in laboratory tests. In these trials, two species of Trichogramma performed best at a smaller scale, while the third performed best at the largest spatial scale observed. Alternatively, multiple investigators have shown that habitat complexity affects *Trichogramma* foraging success in the laboratory (Andow and Prokrym 1990; Lukianchuk and Smith 1997; Gingras and Boivin 2002; Gingras et al. 2002, 2003; Andow and Olson 2003). Thus, it is possible that physical/structural homogeneity of the arenas used in this trial may have predisposed all three species to parasitize more hosts than when they were in a more heterogeneous habitat.

Although not entirely conclusive, these results suggest that T. deion is the best-suited species for biological control of P. interpunctella. T. deion parasitized slightly more host eggs than the other two species in all four treatments and significantly more than T. pretiosum in the 36 four-egg patches treatment (Figure 1). Furthermore, while intrapatch egg density appeared to increase patch discovery for T. ostriniae at the greater interpatch density, T. deion did not appear to be affected by this factor. Because the presence of a single P. interpunctella larva, pupa, or adult can result in the total economic loss of the product (Subramanyam et al. 2001), location of individual host eggs is of critical importance for Trichogramma used in retail stores or warehouses. Other studies comparing the three species' performance at either a finer spatial scale in response to microhabitat complexity or at a broader spatial scale examining the ability of mass-released Trichogramma to locate eggs on shelving units found that T. deion parasitized/killed significantly more *P. interpunctella* eggs than the other two species (Grieshop et al. 2007, Grieshop et al. 2008).

# **Acknowledgements**

We thank Beverly Wilson (Kansas State University) and Kenlee Friesen (USDA-ARS) for their help with colony maintenance and experimental set up. We thank Kimberly With (KSU Division of Biology), Matthias Schöller (Biologische Beratung Berlin), Jim Campbell (USDA-ARS), and Lukasz Stelinski (University of Florida) for help with initial manuscript revision, as well as James Higgins (Kansas State University, Department of Statistics) for assistance with data analysis. We thank Synthia Penn (Beneficial Insectaries), Michael Hoffman and Jeffery Gardner (Cornell University) for providing us with biological material. This research was funded by: USDA/CSREES (RAMP) under agreement number 00-51101-9674. This is Contribution Number 08-150-J from the Kansas Agricultural Experiment Station.

This paper reports the results of research only. Mention of a proprietary product or trade name does not constitute a recommendation or endorsement by the US Department of Agriculture, or Kansas State University.

# References

Andow DA, Olson DM. 2003. Inheritance of host finding ability on structurally complex surfaces. *Oecologia* 136: 324-328.

Andow DA, Prokrym DR. 1990. Plant structural complexity and host-finding by a parasitoid. *Oecologia* 82: 162-165.

Arbogast RT, Mullen MA. 1978. Spatial distribution of eggs by ovipositing Indian meal moths, *Plodia interpunctella* (Hübner) (Lepidoptera: Pyralidae). *Researches on Population Ecology* 19: 148-154.

Arthur FH, Rogers T. 2003. Legislative and regulatory actions affecting insect pest management for postharvest systems in the United States. In: Credland PF, Armitage DM, Bell CH, Cogan PM, Highley E, editors. Advances in Stored Product Protection, Proceedings of the 8th International Working Conference on Stored Product Protection, July 22-26, 2002, York, UK. pp. 435-438.

Barnay O, Pizzol J, Gertz C, Kienlen JC, Hommay G, Lapchin L. 1999. Host density-dependence of discovery and exploitation rates of egg patches of *Lobesia botrana* (Lepidoptera: Tortricidae) and *Ephestia kuhniella* (Lepidoptera: Pyralidae) by the parasitoid *Trichogramma cacoeciae* (Hymenoptera: Trichogrammatidae). *Journal of Economic Entomology* 92: 1311-1320.

Boo KS, Yang JP. 2000. Kairomones used by *Trichogramma chilonis* to find *Helicoverpa assulta* eggs. *Journal of Chemical Ecology* 26: 359-375.

Brower JH. 1988. Population suppression of the almond moth and the Indian meal moth (Lepidoptera: Pyralidae) by release of *Trichogramma pretiosum* (Hymenoptera: Trichogrammatidae) into simulated peanut storages. *Journal of Economic Entomology* 81: 944-948.

Brower JH. 1984. The natural occurrence of the egg parasite, *Trichogramma*, on almond moth eggs in peanut storages in Georgia. *Journal of the Georgia Entomological Society* 19: 285-290.

Cox PD, Bell CH. 1991. Biology and ecology of moth pests of stored foods. In: JR Gorham JR, editor. *Ecology and Management of Food-Industry Pests*. pp. 181-193. FDA Technical Bulletin 4.

Gingras D, Boivin G. 2002. Effect of plant structure, host density and foraging duration on host finding by *Trichgramma evanescens* (Hymenoptera: Trichogrammatidae). *Environmental Entomology* 31: 1153-1157.

Gingras D, Dutilleul P, Boivin G. 2003. Effect of plant structure on host finding capacity of lepidopterous pests of crucifers by two *Trichogramma* parasitoids. *Biological Control* 27: 25-31.

Gingras D, Dutilleul P, Boivin G. 2002. Modeling the impact of plant structure on host-finding behavior of parasitoids. *Oecologia* 130: 396-402.

Grieshop MJ, Flinn PW, Nechols JR, Campbell JF. 2008. Effects of fine-grain habitat complexity on egg parasitism by three species of *Trichogramma*. *Biological Control* 45(3): 328-336.

Grieshop MJ, Flinn PW, Nechols JR. 2006. Biological control of Indianmeal moth (Lepidoptera: Pyralidae) on finished stored products using egg and larval parasitoids. *Journal of Economic Entomology* 99: 1080-1084.

Grieshop MJ, Flinn PW, Nechols JR, Schöller M. 2007. Host-foraging success of three species of *Trichogramma* in simulated retail environments. *Journal of Economic Entomology* 100: 591-598.

Hassel MP. 1982. Patterns of parasitism by insect parasitoids in patchy environments. *Ecological Entomology* 7: 365-377.

Hoffmann MP, Wright MG, Pitcher SA, Gardner J. 2002. Inoculative releases of *Trichogramma ostriniae* for suppression of *Ostrinia nubilalis* (European corn borer) in sweet corn: Field biology and population dynamics. *Biological Control* 25: 249-258.

Jones RL, Lewis WJ, Beroza M, Bierl BA, Sparks AN. 1973. Host-seeking stimulants (kairomones) for the egg parasite, *Trichogramma evanescens. Environmental Entomology* 2: 593-596.

Kfir R. 1983. Functional response to host density by the egg parasite *Trichogramma pretiosum*. *Entomophaga* 28: 345-353.

Lukianchuk JL, Smith SM. 1997. Influence of plant structural complexity on the foraging success of *Trichogramma minutum*: A comparison of search on artificial and foliage models. *Entomologia Experimentalis et Applicata* 84: 221-228.

McGarigal K, Cushman SA, Neel MC, Ene E. 2002. FRAGSTATS: Spatial Pattern Analysis Program for Categorical Maps. Computer software program produced by the authors at the University of Massachusetts, Amherst. Available at:

www.umass.edu/landeco/research/fragstats/fragstats.html.

McCravy KW, Berisford CW. 1998. Parasitism by *Trichogramma* spp. (Hymenoptera: Trichogrammatidae) in relation to Nantucket pine tip moth (Lepidoptera: Tortricidae) egg density and location. *Environmental Entomology* 27: 355-359.

McGaughey WH, Beeman RW. 1988. Resistance to *Bacillus thuringiensis* in colonies of Indian meal moth and almond moth (Lepidoptera: Pyralidae). *Journal of Economic Entomology* 81: 28-33.

Morrison G, Lewis WJ, Nordlund DA. 1980. Spatial differences in *Heliothis zea* egg density and the intensity of parasitism by *Trichogramma* spp.: An experimental analysis. *Environmental Entomology* 9: 79-85.

Mullen MA. 1994. Rapid determination of the effectiveness of insect resistant packaging. *Journal of Stored Product Research* 30: 95-97.

Nordlund DA, Lewis WJ, Gross HR, Beevers M. 1981. Kairomones and their use for management of entomophagous insects. XII. The stimulatory effects of host eggs and the importance of host-egg density to the effective use of kairomones for *Trichogramma pretiosum* Riley. *Journal of Chemical Ecology* 7: 909-917.

Perez-Mendoza J, Agulera-Pena M. 2004. Development, reproduction, and control of the Indian meal moth, *Plodia interpunctella* (Hubner) (Lepidoptera: Pyralidae), in stored seed garlic in Mexico. *Journal of Stored Product Research* 40: 409-421.

Prozell S and Schöller M. 1998. Insect fauna of a bakery, processing organic grain and applying *Trichogramma evanescens*Westwood. *Integrated Protection of Stored Products* (IOBC Bulletin) 21: 39-44.

Reznik S, Umarova Ya, Umarova TYa. 1991. Host population density influence on host acceptance in *Trichogramma*. *Entomologica Experimentalis et Applicata* 58: 49-54.

SAS Institute. 2000. SAS release 8.01. SAS Institute.

Schöller M, Fields P. 2002. Evaluation of North American species of Trichogramma Westwood (Hymenoptera:

Trichogrammatidae) for control of the Indian meal moth, *Plodia interpunctella* (Hübner) (Lepidoptera: Pyralidae). *Advances in Stored Product Protection. Proceedings of the 8th International Working Conference on Stored Product Protection* 22: 26.

Schöller M, Flinn PW, Grieshop MJ,

Zdarkova E. 2006. Biological control of stored product pests. In: Heaps JW, editor. *Insect management for food storage and processing. Second edition.* pp. 67-87. AACC International.

Schöller M, Hassan SA, Reichmuth Ch. 1996. Efficacy assessment of *Trichogramma* evanescens and *T. embryophagum* (Hym.: Trichogrammatidae) for control of stored products moth pests in bulk wheat. *Entomophaga* 41: 125-132.

Schöller M, Reichmuth Ch, Hassan SA. 1994. Studies on biological control of *Ephestia kuehniella* Zeller (Lep.: Pyralidae) with *Trichogramma evanescens* Westwood (Hym.: Trichogrammatidae) - host finding ability in wheat under laboratory conditions. In: Highley E, Wright EJ, Banks HJ, Champ BR, editors. *Stored Product Protection:* Proceedings of the 6th International Working Conference on Stored-Product Protection, February 1994. CAB International.

Steidle JLM, Rees D, Wright EJ. 2001. Assessment of Australian *Trichogramma* species (Hymenoptera: Trichogrammatidae) as control agents of stored product moths. *Journal of Stored Product Research* 37: 263-275.

Smith SM. 1996. Biological control with *Trichogramma*: Advances, successes, and potential of their use. *Annual Review of Entomology* 41:375-406.

Subramanyam Bh, Campbell J, Kemp K. 2001. It's in the detail for retail. *Pest Control* May: 26-28.

Thorpe KW, Dively GP. 1985. Effects of arena size on laboratory evaluations of the egg parasitoids *Trichogramma minutum*, *T. pretiosum*, and *T. exiguum* (Hymenoptera:

Trichogrammatidae). *Environmental Entomology* 14: 762-767.

Van Driesche R, Bellows TS. 2001. Augmentation of parasitoids, predators, and beneficial herbivores. In: Van Driesche R, Bellows TS, editors. *Biological Control 2<sup>nd</sup> edition*. pp. 178-234. Kluwer Academic Publishers.

Wang B, Ferro DN. 1998. Functional responses of *Trichogramma ostriniae* (Hymenoptera: Trichogrammatidae) to *Ostrinia nubilalis* (Lepidoptera: Pyralidae) under laboratory and field conditions. *Environmental Entomology* 27: 752-758.

Wang B, Ferro DN, Hosmer DW. 1999. Effectiveness of *Trichogramma ostriniae* and *T. nubilale* for controlling the European corn borer *Ostrinia nubilalis* in sweet corn. *Entomologia Experimentalis et Applicata* 91: 297-303.

Wang B, Ferro DN, Hosmer DW. 1997. Importance of plant size, distribution of egg masses, and weather conditions on egg parasitism of the European corn borer, *Ostrinia nubilalis* by *Trichogramma ostriniae* in sweet corn. *Entomologia Experimentalis et Applicata* 83: 337-345.

Wright MG, Kuhar TP, Hoffmann MP, Chenus SA. 2002. Effect of inoculative releases of *Trichogramma ostriniae* on populations of *Ostrinia nubilalis* and damage to sweet corn and field corn. *Biological Control* 23: 149-155.