



Mechanisms of the Modulation of Pacemaker Activity by GnRH Peptides in the Terminal Nerve-GnRH Neurons

Authors: Abe, Hideki, and Oka, Yoshitaka

Source: Zoological Science, 19(1) : 111-128

Published By: Zoological Society of Japan

URL: <https://doi.org/10.2108/zsj.19.111>

Mechanisms of the Modulation of Pacemaker Activity by GnRH Peptides in the Terminal Nerve-GnRH Neurons

Hideki Abe^{1,2} and Yoshitaka Oka^{1*}

¹Misaki Marine Biological Station, Graduate School of Science, The University of Tokyo, Kanagawa 238-0225, Japan.

²Present address: Department of Physiology, Tokyo Medical and Dental University, Graduate School and Faculty of Medicine, Yushima 1-5-45, Bunkyo-ku, Tokyo 113-8519, Japan

ABSTRACT—According to our working hypothesis, the terminal nerve (TN)-gonadotropin releasing hormone (GnRH) system functions as a neuromodulatory system that regulates many long-lasting changes in animal behaviors. We have already shown by using *in vitro* whole brain preparations of a small fish (dwarf gourami) that the pacemaker activities of TN-GnRH neurons are modulated biphasically by salmon GnRH, which is the same molecular species of GnRH produced by TN-GnRH neurons themselves; the modulation consists of initial transient decrease and late increase of firing frequency. In the present study, we investigated the possible involvement of Ca²⁺ release from intracellular store and voltage dependent Ca²⁺ currents in the modulation of pacemaker activities. Pharmacological blockade of Ca²⁺ release from intracellular stores or apamin-sensitive Ca²⁺-activated K⁺ current inhibited the initial transient decrease of firing frequency by sGnRH. On the other hand, bath application of Ca²⁺ channel blockers Ni²⁺ or La³⁺ slowed down the pacemaker frequency and attenuated the rate of the late increase of pacemaker frequency by GnRH. Furthermore, voltage-clamp experiments suggested that low-voltage-activated (LVA) Ca²⁺ current and high-voltage-activated (HVA) Ca²⁺ current were present in the TN-GnRH neurons, and bath application of GnRH shifted the activation threshold of HVA Ca²⁺ current to more negative potentials. These results suggest that (1) sGnRH induces Ca²⁺ release from intracellular stores and activates apamin-sensitive Ca²⁺-activated K⁺ current so that it decreases the frequency of pacemaker activity in the initial phase, (2) some kinds of Ca²⁺ currents contribute to the generation and modulation of pacemaker activities, and (3) HVA Ca²⁺ current is facilitated by sGnRH so that it increases the frequency of pacemaker activity in the late phase.

Key words: GnRH, peptide, pacemaker, neuromodulation, electrophysiology

INTRODUCTION

The gonadotropin-releasing hormone (GnRH) was originally identified as a hypophysiotropic decapeptide hormone that is produced in the preoptic area, transported to the median eminence, and facilitates the release of gonadotropins from the pituitary. However, GnRH neurons and their fibers have been found not only in the preoptic area but also in several brain areas outside the hypothalamic area. Such 'extrahypothalamic' GnRH systems have been found mainly in the terminal nerve (TN) and midbrain (Schwanzel-Fukuda and Silverman, 1990; Parhar and Iwata, 1994; Yamamoto *et al.*, 1995), and they project widely in the brain instead of the

median eminence or the pituitary.

By taking advantage of anatomical feature of the *in vitro* whole brain preparation of the dwarf gourami, we have been studying the electrophysiological characteristics of the TN-GnRH system (reviewed by Oka, 1997; Oka and Abe, 2002). We have previously shown that (1) individual cells of TN-GnRH system project widely in the entire brain from the olfactory bulb to the spinal cord (Oka and Matsushima, 1993), and (2) single TN-GnRH neurons show spontaneous pacemaker activity (Oka and Matsushima, 1993) which consists of the depolarizing phase produced by I_{Na(slow)} (Oka, 1995, 1996) and the repolarizing phase mainly produced by TEA-sensitive persistent potassium current, I_{K(v)} (Abe and Oka, 1999). Moreover, the frequency of pacemaker activity seems to vary according to the physiological conditions of the animal, and the firing frequency and/or firing mode may

* Corresponding author: Tel. +81-468-81-4105;
FAX. +81-468-81-7944.
E-mail: okay@mmb.s.u-tokyo.ac.jp

affect the efficacy of exocytosis of GnRH peptides from the GnRH neurons (Peng and Horn, 1991). On the other hand, a growing body of evidence suggests that GnRH peptide modulates the function of ion channels such as Na⁺ channels (Eisthen *et al.*, 2000), K⁺ channels (Adams and Brown, 1980) and Ca²⁺ channels (Elmslie *et al.*, 1990) and thus may regulate the excitability or neurotransmitter release of target neurons. Thus, it has been suspected that the TN-GnRH system may function as a neuromodulatory system that is involved in the regulation of long-lasting changes in the animal's behavior, e.g., motivational or arousal states (Oka and Matsushima, 1993; Oka, 1997). Therefore, the study of nature and mechanisms of modulation of the pacemaker activity of TN-GnRH neurons by hormones or transmitters will give us invaluable information about the control mechanism of neuromodulatory GnRH system.

We have previously indicated that the frequency of pacemaker activity of TN-GnRH neurons was biphasically modulated by sGnRH in a dose dependent and paracrine/autocrine manner (Abe and Oka, 2000, 2002). This biphasic modulation consisted of the transient decrease (early phase) and subsequent increase (late phase) of firing frequency. Furthermore, it was suggested that the G-protein coupled GnRH receptor in the cell membrane triggers signal transduction pathway that ended up modulating the frequency of pacemaker activity of TN-GnRH neurons. This biphasic modulation of pacemaker activity may contribute to a synchronized facilitation of pacemaker activities of neighboring TN-GnRH neurons.

In other types of neurons, it has been proposed that G-protein coupled receptors can modulate the gating properties of the ion channels. Both direct effects of G-proteins and indirect effects via diffusible second messengers have been implicated in these modulations (Hille, 1994). Generally, it has been suggested that GnRH receptors are coupled to the G_{q/11} type G-proteins, and the G_{q/11} type G-proteins enhance inositol phosphate formation and subsequent increase in [Ca²⁺]_i (Naor *et al.*, 1998; Stojilkovic *et al.*, 1994a, b; Stojilkovic and Catt, 1995a, b). In the pituitary gonadotrophs, it has been accepted that GnRH-induced activation of phospholipase C is the major signal transduction pathway of GnRH receptor-coupled processes, and the subsequent mobilization of [Ca²⁺]_i and the activation of protein kinase C (PKC) are the key elements in the control of gonadotropin secretion by pituitary gonadotrophs (Stojilkovic and Catt, 1995a, b).

On the other hand, the dependence on the extracellular Ca²⁺ of episodic release of GnRH from perfused hypothalamic neurons and GT1-7 cells suggests that GnRH secretion is controlled by Ca²⁺ entry through the plasma membrane Ca²⁺ channels (Krsmanovic *et al.*, 1992). Electrophysiological studies have demonstrated the expression of several types of plasma membrane Ca²⁺ channels in the embryonic GnRH neurons (Kusano *et al.*, 1995) and GT1 cells (Bosma, 1993; Costantin and Charles, 1999; Javors *et al.*, 1995; Van Goor *et al.*, 1999b), including transient and

sustained voltage-dependent Ca²⁺ channels.

Therefore, in the present paper, we examined the possible involvement of [Ca²⁺]_i mobilization from the intracellular stores and the plasma membrane Ca²⁺ channels in the biphasic modulations of pacemaker activity in the TN-GnRH neurons.

MATERIAL AND METHODS

Preparations

Adult male and female dwarf gourami (*Colisa lalia*), ~4 cm in standard length, were purchased from a local dealer and kept at 22 ~ 27 °C until used. The whole brain *in vitro* preparation was made according to the procedures described in the previous papers (Oka, 1995, 1996; Abe and Oka, 1999, 2000).

Electrophysiology

The *in vitro* whole brain preparation was continuously superfused with an oxygenated Ringer solution containing (in mM) 124 NaCl, 5 KCl, 1.2 KH₂PO₄, 2.4 CaCl₂, 1.3 MgSO₄, 26 NaHCO₃, and 10 glucose (pH 7.4 adjusted with NaOH) in the silicone elastomer (Shin-Etsu Silicone No.KE-106, Shin-Etsu Chemical Co., Ltd., Japan) base of a small recording chamber. Whole cell voltage- and current-clamp recordings were carried out with the use of CEZ-2300 amplifier (Nihon Kohden, Japan) and pCLAMP software (Axon instruments). Pipette resistance was ~8 MΩ, and seal resistance was >10GΩ. Series resistance was compensated as much as possible. The patch pipette was visually guided to the cluster of TN-GnRH neurons exposed on the ventral surface of the brain under a dissecting microscope (Oka and Matsushima, 1993). After gigaohm seal formation and "break in" for the whole cell recording mode, characteristic spontaneous pacemaker activity was confirmed in the current-clamp mode (see Oka and Matsushima, 1993; Abe and Oka, 2000).

For current-clamp experiments, the patch pipettes contained (in mM) 110 K-gluconate, 3 MgCl₂, 40 N-2-hydroxyethylpiperazine-N'-2-ethanesulfonic acid (HEPES), 0.3 ethylene glycol-bis (β-aminoethyl ether)-N,N,N',N'-tetraacetic acid (EGTA), 2 Na₂ATP, and 0.2 Na₂GTP (pH 7.4 adjusted with NaOH), NiCl₂ (1 mM), LaCl₃ (5 μM), Apamin (100 nM; Alomone Labs.) and salmon GnRH (sGnRH, 200 ~ 300 nM; RBI) were dissolved directly in the Ringer solution. Ruthenium red (50 μM; Sigma) and heparin (300 μg / ml; Sigma) were dissolved in the pipette solution. Pacemaker activities were digitized (2 kHz), displayed on-line with Axotape software or pClamp8 (Axon Instruments), and stored on a computer.

For voltage-clamp recording of Ca²⁺ currents, the extracellular solution contained (in mM) 95 choline Cl, 40 tetraethylammonium chloride (TEACl), 2.4 CaCl₂, 1.3 MgCl₂, 5 Glucose, 5 4-aminopyridine (4AP), and 10 HEPES (pH 7.4 adjusted with NaOH). In addition, 0.75 μM tetrodotoxin (TTX) was added to the extracellular solution. Patch pipettes were filled with a solution consisting of (mM) 90 CsCl₂, 2 MgCl₂, 20 TEACl, 10 EGTA, 10 HEPES, 2 Na₂ATP, and 0.2 Na₂GTP (pH 7.4 adjusted with CsOH). For voltage-clamp recording of tentative Ca²⁺-activated K⁺ currents, the extracellular solution contained (in mM) 115 choline Cl, 20 TEACl, 5 KCl, 2.4 CaCl₂, 1.3 MgCl₂, 5 glucose, and 10 HEPES (pH 7.4 adjusted with NaOH). 0.75 μM TTX was also added to the extracellular solution. Patch pipettes were filled with a solution consisting of (mM) 110 KCl, 3 MgCl₂, 40 HEPES, 0.3 EGTA, and 2Na₂ATP (pH 7.4 adjusted with NaOH). The linear leakage currents were digitally subtracted, either automatically with the use of the P/4 protocol, or manually, after measuring ohmic resistance in response to hyperpolarizing command pulses. The data were not corrected for the liquid junction potentials.

Data analysis

Data analysis, fitting, averaging and presentation were carried out using a combination of pCLAMP6 and 8, Axograph (Axon Instruments), Microsoft Excel (Microsoft), DeltaGraph (Polaroid), and Canvas (Deneba software) softwares. All data are present as means \pm SE.

RESULTS

Most of the TN-GnRH neurons showed regular spontaneous pacemaker activity, and it was biphasically modulated by bath application of sGnRH, as we have reported previously (Abe and Oka, 2000; also see Fig. 3Ba~d).

Involvement in the control of pacemaker activity of Apamin-sensitive Ca²⁺-activated K⁺ currents induced by Ca²⁺ released from the intracellular store

First, in order to examine the possible involvement of Ca²⁺ released from the intracellular store in the control of pacemaker activity of TN-GnRH neuron, the cells were dialyzed with ruthenium red, an inhibitor of Ca²⁺-induced Ca²⁺ release, and heparin, that of IP₃-induced Ca²⁺ release, respectively, by including them in the patch pipette solution. After that, the effects of bath application of sGnRH on the pacemaker activity were examined. To ensure the diffusion of ruthenium red and heparin into the cytoplasm of the TN-GnRH neuron, the data collection was started 10 min after the whole-cell recording was established; this time period was determined on the basis of the results of intracellular application of QX-314 (see Abe and Oka, 2000). Fig. 1 shows the pacemaker activity of a cell recorded with patch pipette containing ruthenium red (50 μ M) and heparin (300 μ g / ml) in the pipette solution. In Ringer solution, the cells showed slightly irregular beating discharge pattern (Fig. 1Ba). Many cells tended to show such a firing pattern by intracellular application of the blockers of Ca²⁺ release from the intracellular store. In these experiments, we further applied La³⁺ (5 μ M) to the bath solution in order to block some Ca²⁺ channels (see below). Bath application of La³⁺ slowed down the frequency of pacemaker activity (Fig. 1Bb). Further bath application of sGnRH (1 μ M) failed to evoke transient decrease of firing frequency (Fig. 1Bc) but evoked subsequent increase of firing frequency (Fig. 1Bd). Figure 1Bc and Bd (20 and 120 s after the onset of sGnRH perfusion, respectively) were recorded during time periods similar to that of transient decrease and subsequent increase of firing frequency by sGnRH. Fig. 1Bf shows the time course of these changes before and after the application of sGnRH (during the period corresponding to Fig. 1Bb~d).

We compared the degree of transient decrease of firing frequency in the presence and absence of the intracellular Ca²⁺ mobilization blockers. Each effect of the blockers was measured during a 20 s period that was 30 ~ 50 s after the onset of sGnRH perfusion, which corresponded to the early transient phase of pacemaker frequency decrease, and was normalized to the frequency in Ringer solution. The transient decrease that was induced by sGnRH (0.88 \pm 0.06; n=16)

was nullified by heparin (1.01 \pm 0.05; n=5). Similarly, intracellular application of ruthenium red alone or both ruthenium red and heparin nullified the transient decrease or rather increased slightly the firing frequency (1.11 \pm 0.14; n=7 and 1.16 \pm 0.11; n=4, respectively). However, we could not find statistically significant nullifying effects of blockers of Ca²⁺ release from the intracellular store due to the large SE. Anyway, these results suggest that the Ca²⁺ release from the intracellular store may be involved in the transient decrease of firing frequency that was induced by sGnRH.

In our previous studies of voltage-dependent K⁺ currents (Abe and Oka, 1999), we found that a transient and 4AP-sensitive, but not TEA- or charybdotoxin-sensitive Ca²⁺-dependent outward current component was present in TN-GnRH neurons. Because this current could be observed only when Ca²⁺ was present in the external solution, we defined this 4AP-sensitive transient outward current that was dependent on the presence of extracellular Ca²⁺ ions as tentative Ca²⁺-activated K⁺ currents. When the current responses were measured in Ca²⁺-containing extracellular solutions containing 0.75 μ M TTX and 20 mM TEA, a mixture of large transient currents and smaller persistent outward currents could be recorded in response to depolarizing pulses from a holding potential of -100 mV. The activation and steady-state inactivation curves of the mixture current containing tentative Ca²⁺-activated K⁺ current are shifted to more depolarized potentials compared with those of "4AP-sensitive transient current" (data not shown). These tentative Ca²⁺-activated K⁺ currents are only partially inactivated at membrane potentials of -60 to -40 mV, which correspond to the base membrane potentials of the pacemaker activity of TN-GnRH neurons. Thus, it is reasonable to think that tentative Ca²⁺-activated K⁺ current can be activated in the pacemaker range of TN-GnRH neurons.

Next, we examined the effect of sGnRH on the tentative Ca²⁺-activated K⁺ current. Fig. 2A shows the superimposed traces of the tentative Ca²⁺-activated K⁺ currents before, during, and after sGnRH applications elicited during a series of +50 mV test pulses from a holding potential of -100 mV. In this case, bath application of sGnRH (1 μ M) facilitated current amplitude about 18% (Fig. 2A). The current amplitude recovered to the control level after washout by Ringer solution. Fig. 2B shows the time course of the tentative Ca²⁺-activated K⁺ current before, during, and after sGnRH applications. Test pulses were applied at 15 s intervals. The current amplitude showed a tendency to gradually decrease due to the rundown. However, it was rapidly increased by bath application of sGnRH (within 15 to 90 s; Fig. 2B, solid bars). This time course corresponded to the onset of the transient decrease of firing frequency of pacemaker activity induced by sGnRH. Averaged increase of the amplitude of tentative Ca²⁺-activated K⁺ current was about 9% (2 to 19%, n=4).

Finally, we examined the effects of bath application of apamin, a specific SK-type Ca²⁺-activated K⁺ channel blocker. Fig. 3 shows the effect of bath application of




Fig. 1. Inhibition of Ca²⁺ release from the intracellular store disrupts the transient decrease of pacemaker frequency after GnRH application. (A) Continuous recording of the pacemaker activity. Traces indicated by bars a-f are shown in (B) on enlarged time scales. Similar conventions are used in Figs. 3, 5, and 6. Intracellular application of Ruthenium Red (50 μ M), which inhibits the Ca²⁺-induced Ca²⁺ release, and heparin (300 μ g/ml), which inhibits IP₃-induced Ca²⁺ release, blocked transient decrease (Bc) but not subsequent increase in the frequency of pacemaker activities (Bd), both of which should be induced by sGnRH. Compare with normal changes in the frequency of pacemaker activities that are induced by sGnRH shown in Fig. 3Ba-d.




Fig. 2. Tentative Ca^{2+} -activated transient K^+ current is facilitated by sGnRH. (A) Voltage clamp recordings of tentative Ca^{2+} -activated transient K^+ currents. Bath application of 1 mM sGnRH increased the amplitude of tentative Ca^{2+} -activated transient K^+ current. Current traces before, during, and after sGnRH application are shown. (B) shows the time course of the peak amplitude of the tentative Ca^{2+} -activated transient K^+ current during sGnRH applications (bars).

apamin. In Ringer, TN-GnRH neurons showed regular beating discharge (3.02 ± 0.86 Hz; $n=4$). Bath application of sGnRH (20 nM) transiently decreased (2.74 ± 0.83 Hz; Fig. 3Bb) and subsequently increased (4.98 ± 0.60 Hz; Fig. 3Bc) firing frequency of pacemaker activity. After washout of sGnRH, bath application of apamin (100 nM) facilitated the frequency of pacemaker activity (3.60 ± 0.68 Hz; Fig. 3Be). However, further bath application of sGnRH (20 nM) failed to evoke transient decrease of firing frequency (3.70 ± 0.74 Hz; Fig. 3Bf) but evoked subsequent increase of firing frequency (6.59 ± 1.15 ; Fig. 3Bg). Fig. 3Bd and Bh show the time course of these changes before and after the application of sGnRH. We also compared quantitatively the normalized transient decrease of firing frequency in the presence and absence of apamin. The transient decrease that was induced by sGnRH (0.92 ± 0.07 ; $n=4$) was nullified by apamin

(1.02 ± 0.03 ; $n=4$).

From these data, it is suggested that sGnRH triggers the Ca^{2+} release from the intracellular store and activates apamin-sensitive Ca^{2+} -activated K^+ currents, which leads to the transient decrease of the frequency of pacemaker activity.

Involvement of Ca^{2+} currents in the pacemaker activity

Because the preliminary experiments that examined the effect of sGnRH upon $I_{\text{Na}(\text{slow})}$ and/or $I_{\text{K}(\text{V})}$ did not show any noticeable modulation, we investigated the possible involvement of Ca^{2+} currents in the modulation of pacemaker activity. Fig. 4 shows the effects of Ca^{2+} channel blockers on the pacemaker frequency of TN-GnRH neurons. Bath application of various Ca^{2+} channel blockers slowed down the pacemaker frequency of TN-GnRH neurons. Fig. 4Aa shows




Fig. 3. Inhibition of SK-type Ca^{2+} -activated K^+ channel disrupts the transient decrease of pacemaker frequency after GnRH application. (A) Continuous recording of the pacemaker activity. In Ringer solution, bath application of sGnRH (20 nM) transiently decreased (Bb) and subsequently increased (Bc) the frequency of pacemaker activity (the overall time course of these changes is shown in Bd). By bath application of apamin (100 nM), which inhibits the SK type Ca^{2+} -activated K^+ channel, the frequency of pacemaker activity was increased (Be). Furthermore, bath application of apamin blocked transient decrease (Bf) but not subsequent increase in the frequency of pacemaker activities by sGnRH (Bg)(the overall time course of these changes is shown in Bh). Following washout by Ringer solution, the firing frequency decreased again.




Fig. 4. Blockade of Ca^{2+} currents decreases the pacemaker frequency. Aa, b: Effects of blockers of Ca^{2+} currents, La^{3+} and Ni^{2+} , respectively. Both of these treatments decreased the frequency of pacemaker activities. Following washout, the frequency of pacemaker activities recovered. B: Comparison of the effects of Ca^{2+} channel blockers on the pacemaker activity. The numbers in parentheses represent the numbers of cells tested for each blocker. The ordinate indicates the relative pacemaker frequencies after each treatment normalized to those before the treatment; (Frequency of pacemaker activity in Ringer solution containing Ca^{2+} channel blocker) / (Frequency of pacemaker activity in Ringer solution). Bath application of Ca^{2+} channel blockers decreased the pacemaker frequency. It should be noted that simultaneous bath application of both La^{3+} and Ni^{2+} (the right column in B) had an additive effect of blocking the frequency of pacemaker activity (compare with the left and middle columns in B).

the effects of the blocker of high-voltage-activated Ca^{2+} currents, La^{3+} (5 μM , $n=4$), and Fig. 4Ab shows those of the blockers of low-voltage-activated Ca^{2+} current and store-operated Ca^{2+} current, Ni^{2+} (1 mM, $n=4$). The frequencies of pacemaker activities before and after these treatments were 3.7 ± 0.8 and 2.6 ± 0.5 Hz for La^{3+} ($P < 0.05$, Student's two-tailed paired t test) and 4.1 ± 0.7 and 1.7 ± 0.2 Hz for Ni^{2+} ($P < 0.05$, Student's two-tailed paired t test). Furthermore, when the bath application of the combination of La^{3+} (5 μM) and Ni^{2+} (1 mM) was examined, it completely blocked the generation of pacemaker activity (in 7/8 cells, Fig. 6Ba, c and d). The results are presented in Fig. 4B as the relative pacemaker frequencies after each treatment normalized to those before the treatment, which were defined as the ratio (firing frequency in Ringer solution containing Ca^{2+} channel blockers) / (firing frequency in Ringer solution). These results indicate that some kinds of Ca^{2+} currents are present in TN-GnRH neurons and contribute to the generation of pacemaker activity.

Next, we examined the effects of Ca^{2+} channel blockers on the modulation of pacemaker activity by sGnRH. Fig. 5 shows the effects of Ni^{2+} on the modulation of pacemaker activity by sGnRH. In Ringer solution, TN-GnRH neuron showed regular beating discharge (3.5 ± 0.5 Hz; Fig. 5Ba). Bath application of Ni^{2+} (1 mM) decreased the firing frequency of pacemaker activity (1.8 ± 0.2 Hz; Fig. 5Bb). Subsequent bath application of sGnRH (300 nM) transiently decreased (1.5 ± 0.2 Hz; Fig. 5Bc) and then increased (2.4 ± 0.8 Hz; Fig. 5Bd) the firing frequency of pacemaker

activity in a qualitatively similar manner with sGnRH application in Ringer. However, the late-phase increase of pacemaker frequency was less pronounced compared with sGnRH application in Ringer (Fig. 7; compare the left and right columns). Similarly, the late-phase increase of pacemaker frequency was less pronounced under La^{3+} treatment compared with sGnRH application in Ringer (Fig. 7; compare the left and middle columns). During the simultaneous bath application of La^{3+} (5 μM) and Ni^{2+} (1 mM) (Fig. 6), the pacemaker activity was stopped (Fig. 6Bb, Bc), and further bath application of sGnRH (300 nM, in 3/4 cells, Fig. 6Bd) or depolarizing DC current injection (in 2/2 cells, data not shown) did not reinstate the pacemaker activities.

Taken together, these data indicate that some kind(s) of Ca^{2+} channels present in the TN-GnRH neurons may be the target for the modulation by sGnRH, which somehow contributes to the late phase increase of pacemaker frequency.

Voltage-clamp recordings of voltage-dependent Ca^{2+} currents

To reveal the quantitative contribution of Ca^{2+} currents to the generation and modulation of pacemaker activity, it is important to determine the Ca^{2+} channel subtypes and kinetics. Depolarizing steps from a holding potential of -100 mV elicited a mixture of transient and sustained inward current components (Fig. 8Aa). They showed the characteristics of combined low voltage-activated (LVA) and high voltage-activated (HVA) Ca^{2+} currents. The peak inward currents (measured at peak current amplitudes during depo-




Fig. 5. The late-phase increase of pacemaker frequency by sGnRH is less pronounced under Ni^{2+} treatment. Bath application of Ni^{2+} (1 mM) decreased the frequency of pacemaker activity (Bb) but did not qualitatively affect either transient decrease (Bc) or subsequent increase of firing frequency (Bd) by sGnRH. However, the late-phase increase was less pronounced compared with sGnRH application in Ringer (see Fig. 7). Following washout by Ni^{2+} -containing Ringer solution, the firing frequency decreased again (Be).

larizing pulses) began to appear more positive than -60 mV and reached a maximum near -20 mV. The sustained currents (measured at 200 ms after the onset of the test pulse) began to appear more positive than -20 mV and reached a maximum near 0 mV. The currents that were elicited by depolarizing steps from a holding potential of -60 mV are shown in Fig. 8Ab. The transient inward current almost disappeared, and sustained current(s) mainly remained. At this holding potential, the activation of both peak and sustained currents required depolarization more positive than -20 mV,

and the maximum inward current was obtained near -10 mV. In the currents recorded from a different cell, however, the sustained current began to appear more positive than -40 mV and rapidly reached its maximum at -30 mV, when a holding potential was -60 mV (see below).

Fig. 8B shows the averaged current-voltage relationship of transient and sustained Ca^{2+} currents ($n=40$). We defined the current component that was obtained by subtraction of sustained current from peak current, as the 'transient' Ca^{2+} current. On the average, the transient and sustained cur-




Fig. 6. Simultaneous application of La^{3+} and Ni^{2+} stops the pacemaker activity. Bath application of La^{3+} (5 mM) and Ni^{2+} (1 mM) blocked the generation of pacemaker activities (Bb, c). Furthermore, bath application of sGnRH did not induce pacemaker activities of TN-GnRH neurons (Bd). Pacemaker activities recovered after washout by Ringer solution (Bf).

rents measured at -10 mV from a holding potential of -100 mV was -1.41 ± 0.26 and -0.87 ± 0.12 nA, respectively (Fig. 8B and). However, the averaged current-voltage relationship of sustained current elicited from a holding potential of -60 mV showed an activation threshold of -40 mV but also had a shoulder of current at -20 mV (Fig. 8B, arrows).

This shoulder of current may reflect the difference of activation threshold and voltage dependence of sustained currents among different cells. The sustained current measured at -10 mV from a holding potential of -60 mV was -0.80 ± 0.13 nA ($n=18$; Fig. 8B).

These results suggest that a transient inward current




Fig. 7. The late-phase increase of pacemaker frequency by sGnRH is less pronounced in the presence of Ca²⁺ channel blockers. Normalized increase of firing frequency by sGnRH in the presence or absence of Ca²⁺ channel blockers was defined as

$$\frac{\text{Frequency}_{(\text{sGnRH})} - \text{Frequency}_{(\text{Ringer})}}{\text{Frequency}_{(\text{Ringer})}}$$

and was calculated from the data such as those shown in Fig. 5. Normalized increase of pacemaker frequencies was less pronounced under Ni²⁺ or La³⁺ treatment. The numbers in parentheses represent the numbers of cells tested for each Ca²⁺ channel blocker.

component that begins to activate near -60 mV and is largely inactivated when the holding potential is -60 mV represents the LVA Ca²⁺ current. On the other hand, HVA Ca²⁺ current component that are not inactivated at holding potentials of -60 mV consist of two types of sustained currents. The one current begins to activate from -20 mV and reaches its maximum amplitude near -10 mV. The other current begins to activate near -40 mV and rapidly reaches maximum at -30 mV.

Actions of GnRH on Ca²⁺ currents

Previous preliminary studies suggested that neither $I_{\text{Na}(\text{slow})}$ nor $I_{\text{K}(\text{v})}$ is significantly modulated by sGnRH applications (our unpublished observations). Therefore, we suspected that the Ca²⁺ currents may be modulated by sGnRH. Fig. 9 shows an example of the effects of sGnRH on the Ca²⁺ currents. The modulation was examined in 10 cells held at -100 mV. In this case, when $1 \mu\text{M}$ sGnRH was added to the bath solution, the current amplitude and activation threshold of transient current were not changed (Fig. 9Ba and). However, the amplitude of sustained current increased (-0.3 to -1.3 nA) in a wide voltage range. Furthermore, the activation threshold and maximum amplitude of sustained current were shifted to more negative potentials (Fig. 9Bb and). Responses of Ca²⁺ currents to GnRH were variable among the recorded cells. In another case, the amplitudes of both transient and sustained Ca²⁺ currents did not increase. However, the activation threshold and the peak amplitude of sustained Ca²⁺ current were shifted to more negative potentials (data not shown). On the average, the amplitude of the transient Ca²⁺ current measured at its maximum was decreased from $-1.4 \text{ nA} \pm 0.3$ to -1.1 ± 0.4 nA ($P > 0.05$, Student's two-tailed paired t-test; Fig. 10A and

), and those of the sustained Ca²⁺ current was increased from -1.6 ± 0.4 to -1.9 ± 0.4 ($P < 0.05$, Student's two-tailed paired *t* test; Fig. 10B and), before and after bath application of $1 \mu\text{M}$ sGnRH, respectively ($n=10$). Moreover, the activation threshold and the voltage that evokes the maximum amplitude of the sustained Ca²⁺ currents were shifted to more negative potentials (Fig. 10B and), while those of transient Ca²⁺ current were not changed. These results indicate that the sustained HVA Ca²⁺ current is facilitated by sGnRH, i.e., the relative contribution of HVA Ca²⁺ current to the pacemaker activity may be increased by sGnRH. It is then suggested that this increased availability of HVA Ca²⁺-current increases the frequency of pacemaker activity.

DISCUSSION

Involvement of Ca²⁺ release from intracellular store and apamin-sensitive K⁺ current in the transient decrease of firing frequency of TN-GnRH neurons

Intracellular application of ruthenium red and heparin inhibited the transient decrease but not late-phase increase of the pacemaker frequency, both of which were induced by bath application of sGnRH. This result suggests that $[\text{Ca}^{2+}]_i$ mobilization is involved in the transient decrease but not the late-phase increase of firing frequency of pacemaker activity. Then, the presence of tentative Ca²⁺-activated transient K⁺ current was suggested from the result of present voltage-clamp experiments. This current was TEA-insensitive but 4AP-sensitive, and could be only evoked when the Ca²⁺ is present in the extracellular bath solutions. However, the activation and steady-state inactivation curves of this current were shifted to more depolarized potentials compared with the 4AP-sensitive A-like currents of TN-GnRH neurons (Abe and Oka, 1999), and the two currents are considered to be different. The amplitude of the tentative Ca²⁺-activated transient K⁺ current was rapidly increased by bath application of sGnRH. Furthermore, bath application of apamin inhibited the transient decrease but not the late-phase increase of the pacemaker frequency by sGnRH. These results suggest that the apamin-sensitive Ca²⁺-activated transient K⁺ current was activated by Ca²⁺ released from the intracellular store, which had been induced by sGnRH. This may explain the early-phase transient decrease of pacemaker activity by sGnRH.

It has been reported that the Ca²⁺-activated K⁺ current exist in the embryonic GnRH neurons (Kusano *et al.*, 1995) and GT1-7 cells (Spergel *et al.*, 1996; Van Goor *et al.*, 1999). The transient decrease of firing frequency by GnRH, which was induced by the activation of SK type Ca²⁺-activated K⁺ current, has been reported in the GT1-7 cells (Van Goor *et al.*, 1999). It has been generally accepted that the activation of G_{q/11}-coupled GnRH receptor increases the intracellular Ca²⁺ concentration via phosphoinositide signaling pathway (Naor, 1990; Naor *et al.*, 1998; Stojilkovic *et al.*, 1994b), and the increased $[\text{Ca}^{2+}]_i$ may open Ca²⁺-activated K⁺ current (Sah, 1996). All of these reports are in favor of




Fig. 8. Voltage clamp recording reveals the presence of one transient LVA and two types of sustained HVA Ca^{2+} current components. A: Currents evoked during 200 ms voltage steps from -100 to $+50$ mV (holding potential= -100 mV)(a) and from -60 to $+50$ mV (holding potential= -60 mV)(b). B: Current-voltage relations of the Ca^{2+} currents averaged from 40 ($V_h=-100$ mV) and 18 ($V_h=-60$ mV) TN-GnRH neurons. I/V curves were constructed by plotting the transient current amplitudes (filled squares) obtained by subtracting sustained current from peak current, and sustained current amplitudes (filled circles) measured at the end of 200 ms test pulses. The transient current probably correspond to the low-voltage activated (LVA) T type Ca^{2+} current. The I/V curve of the sustained currents evoked from a holding potential of -60 mV (open circles) had a shoulder at -20 mV in addition to the activation threshold of -40 mV (arrows). These may correspond to the activation thresholds of two HVA Ca^{2+} currents.




Fig. 9. sGnRH modulates the Ca²⁺ current components. A: Current traces from a holding potential of -100 mV before (a) and after the addition of 200 nM sGnRH to the bath solutions (b). B: The current-voltage relations of the transient current (a), and the sustained current measured at the end of 400 ms test pulse (b), which were constructed from the traces in A. Symbols are defined in the inset. In this cell, the current amplitude of sustained current was augmented by sGnRH, and the activation threshold of sustained current was shifted to more hyperpolarized potentials.

the above-mentioned mechanisms for the transient decrease of pacemaker activity of TN-GnRH neurons by sGnRH.

Ca²⁺ current component is involved in the pacemaker activity

We have shown that some kind(s) of Ca²⁺ current contribute(s) to the modulation of the pacemaker activity of TN-GnRH neurons. Furthermore, the result that simultaneous bath application of 5 μ M La³⁺ and 1 mM Ni²⁺ completely inhibited the generation of pacemaker activity and its modulation by sGnRH, suggests that Ca²⁺ current components that were blocked by simultaneous application of La³⁺ and Ni²⁺ may be somehow involved in the generation of pacemaker potentials and may be the target of modulation by sGnRH.

The present results may appear to be partly inconsistent with those of the previous study that the Ca²⁺ currents are not essential for the generation of pacemaker potentials (Oka, 1995). One possible explanation may be the difference in the recording methods, intracellular recording (Oka, 1995) vs. whole-cell patch-clamp recording (present study).

It may be possible that the intracellular dialysis of EGTA, which was introduced by the patch pipette solution, may prevent Ca²⁺ channels from Ca²⁺-dependent inactivation (Hille, 2001). Thus, a relatively large proportion of Ca²⁺ channel may be available for the generation of the pacemaker activity in the whole-cell patch-clamp recording. Second possibility is the presence of diffusion barrier. While the ventral meningeal membrane of the brain was not removed in the previous study, it is always completely removed in the patch-clamp experiments. The meningeal membrane may have served as a diffusion barrier for the drug delivery. Thirdly, the bath application of Ca²⁺ channel blockers may have simultaneously blocked other currents (for example, I_{Na(slow)}). Preliminary experiments suggest that Ni²⁺ may inhibit I_{Na(slow)} (data not shown). It has also been reported that Ni²⁺ and La³⁺ also blocks Na⁺/Ca²⁺ exchanger, store-operated Ca²⁺ current, and other voltage-gated ion channels (Nowycky, 1991; Taylor and Brond 1998). In the present study, we used supramaximum concentrations of these Ca²⁺ channel blockers to block the Ca²⁺ influx completely. Especially, the concentrations of Ni²⁺ (1 mM) and La³⁺ (5 μ M) used here are sufficient to block store-operated Ca²⁺ cur-



A Transient Ca^{2+} currentB Sustained Ca^{2+} current

Fig. 10. Averaged I/V relations also show the modulation of Ca^{2+} current by sGnRH. I/V curves were constructed by plotting the averaged current amplitudes ($n=10$) evoked from holding potentials of -100 mV before and after the addition of 200 nM sGnRH to the bath solution. The current-voltage relations of transient current (A) and sustained current (B) are shown.

rents (Skryma *et al.*, 2000). Thus, it may be possible that these Ca^{2+} channel blockers blocked not only voltage-gated Ca^{2+} channels, but also store-operated Ca^{2+} channels, other channels, and/or exchangers. In fact, it has been reported in GT 1-7 cell-lines that store-operated Ca^{2+} current contributes to the modulation of firing activity by GnRH (Van Goor *et al.*, 1999a). Further studies using more specific Ca^{2+} channel blockers are necessary to clarify the quantitative contributions of voltage-dependent Ca^{2+} current and store-operated Ca^{2+} current.

Both LVA and HVA Ca^{2+} currents are present in the TN-GnRH neurons

Voltage-clamp experiments of whole-cell patch-clamp recording suggested that TN-GnRH neuron has at least one LVA Ca^{2+} current component and two HVA Ca^{2+} current components, and the relative prevalence of each current component seems to be different among different TN-GnRH neurons. LVA Ca^{2+} current was a transient current evoked from membrane potentials more positive than -60 mV and reached its maximum at about -20 mV (Figs. 8; Transient current). This current was almost inactivated at a holding potential of -60 mV. It most probably corresponds to the T-type Ca^{2+} current (Nowicky *et al.*, 1985). In contrast, both of the two HVA Ca^{2+} currents inactivated slowly and could be observed at holding potentials of -100 mV and -60 mV. One of the HVA Ca^{2+} current activated at potentials more

positive than -40 mV. The other HVA Ca^{2+} current activated at potentials more positive than -20 mV. Thus, the averaged I/V curve of the sustained current had a shoulder and reached its maximum around 0 mV (Fig. 8B). Preliminary experiments showed that the amplitude of HVA Ca^{2+} current was reduced (= was not completely blocked) by bath application of nifedipine (data not shown). Therefore, at least one of the HVA Ca^{2+} current may correspond to the L-type Ca^{2+} current.

It has been demonstrated that embryonic GnRH neurons (Kusano *et al.*, 1995) and GT1 cells (Bosma, 1993; Hales *et al.*, 1994; Javors *et al.*, 1995; Costantin and Charles, 1999) express several types of plasma membrane Ca^{2+} channels, including transient and sustained voltage-dependent Ca^{2+} channels. Kusano *et al.* (1995) reported that embryonic GnRH neurons possess T- and L-type Ca^{2+} channels, and GT1 cells possess T-, N-, and L-type Ca^{2+} channels. On the other hand, Costantin and Charles (1999) and Van Goor *et al.* (1999b) reported that GT1 cells possess T- and L-type Ca^{2+} channels. The present study used for the first time the adult authentic GnRH neurons, and the results basically agreed well with these reports. However, pharmacological isolation and detailed analysis of kinetic properties of each Ca^{2+} current component have not yet been done in the present study mainly due to the poor space-clamp of the *in vitro* whole-brain preparations of TN-GnRH neurons. Therefore, the use of dissociated TN-GnRH neurons may be

necessary to evaluate quantitatively the contribution of each Ca^{2+} current component to the control of pacemaker and secretory activities of TN-GnRH neurons.

Modulation of HVA type Ca^{2+} channels by GnRH

Bath application of sGnRH shifted the activation threshold of sustained Ca^{2+} current component to more hyperpolarized potentials. Moreover, the increase of sustained current amplitude was observed in some recordings, although complete recovery of Ca^{2+} current amplitude by washout could not be obtained due to the rundown of Ca^{2+} currents. It is generally accepted that GnRH receptors are coupled to $\text{G}_{q/11}$ type G-proteins, and $\text{G}_{q/11}$ type G-proteins are coupled to phosphoinositide-mediated signaling pathways (Stojilkovic *et al.*, 1994a,b). It has also been reported that Ca^{2+} channels (especially, N- and L-type Ca^{2+} channel) are positively modulated by PKC (Bourinet *et al.*, 1992; Dolphin, 1998; Meir *et al.*, 1999; Stea *et al.*, 1995; Swartz, 1993; Yang and Tsien, 1993; Zamponi *et al.*, 1997; Zhu and Ikeda, 1994). Bosma and Hille (1992) reported that immortalized gonadotrope (aT3-1 cell), which also has GnRH receptors, express Ca^{2+} channels, and these Ca^{2+} channel were augmented by application of GnRH or phorbol ester.

From these observations and present results, it is highly possible that certain type(s) of Ca^{2+} currents that contribute to the generation of pacemaker activity are modulated by sGnRH and are involved in the late-phase increase of the pacemaker frequency.

Mechanisms of the generation and modulation of pacemaker activity and possible physiological functions

From the results of the present study, we present a model about the generation and modulation mechanisms of pacemaker activity in the TN-GnRH neuron (Fig. 11). In the intact brain, TN-GnRH neurons show regular beating pacemaker activity. The $I_{\text{Na}(\text{slow})}$, $I_{\text{Na}(\text{fast})}$, and $I_{\text{K}(\text{V})}$ interact in the following manner to generate the pacemaker potentials. The $I_{\text{Na}(\text{slow})}$, which is persistently active in the subthreshold membrane potential range, always supplies the persistent depolarizing drive and gradually depolarizes the membrane potentials. When the membrane potential reaches the activation threshold for $I_{\text{K}(\text{V})}$, outward current develops, and the net flux of current reverses to outward. Then, the membrane potential becomes hyperpolarized and deactivates the K^+ current, and the next cycle begins. This is the subthreshold pacemaker activity, and when the membrane potential reaches the activation threshold for the $I_{\text{Na}(\text{fast})}$, the spiking pacemaker activities ensues. In addition, some kind(s) of Ca^{2+} currents and apamin-sensitive Ca^{2+} -activated K^+ current may be also involved in the generation of pacemaker potentials. Although $I_{\text{Na}(\text{slow})}$ and $I_{\text{K}(\text{V})}$ are essential for the generation of the basic rhythm of pacemaker activities, they do not seem to be modulated by the process described below. When the GnRH peptide binds to the GnRH receptor located on the cell surface of TN-GnRH neurons, $\text{G}_{q/11}$ protein-coupled process starts and activates phospholipase C, which leads to the production of IP_3 and diacylglycerol. IP_3 stimulates the release of Ca^{2+} from the IP_3 -sensitive intracellular Ca^{2+} store, and the increased $[\text{Ca}^{2+}]_i$ then opens




Fig. 11. Model of the biphasic modulation of pacemaker activity of TN-GnRH neuron. The GnRH peptide, which binds to its receptor located at the cell membrane of TN-GnRH neurons, induces biphasic modulation of the pacemaker activities; (1) Facilitates Ca^{2+} release from the intracellular Ca^{2+} store. The increased $[\text{Ca}^{2+}]_i$ activates apamin-sensitive Ca^{2+} -activated K^+ current, $I_{\text{K}(\text{Ca})}$. This, in turn, decreases the frequency of pacemaker activities transiently. (2) Up-regulates the Ca^{2+} current(s), I_{Ca} , and increases the frequency of pacemaker activities in the late phase. $I_{\text{Na}(\text{slow})}$, $I_{\text{K}(\text{V})}$, and $I_{\text{Na}(\text{fast})}$ are involved in the generation of basic pattern of pacemaker activities but are not directly modulated by GnRH.

apamin-sensitive Ca^{2+} -activated K^+ channel. Thus, the pacemaker frequency of TN-GnRH neuron is transiently decreased. On the other hand, diacylglycerol activates protein kinase C, and protein kinase C then phosphorylates and modulates the Ca^{2+} current. Thus, the pacemaker frequency of TN-GnRH neurons is increased in the late phase.

What is the physiological significance of the pacemaker activity of TN-GnRH neurons and its modulation? Unfortunately, the functional link between the electrical activities of peptidergic neurons and the peptide release has not been firmly established thus far. A recent amperometric and RIA studies showed that the membrane depolarization triggers the secretion of GnRH from the TN-GnRH neuron in the teleost brain-pituitary slice preparation (Ishizaki and Oka, 2001 and in preparation). In the intact brain, the pacemaker activity of TN-GnRH neurons is characterized by low frequency (<10 Hz) regular beating discharge. The depolarization that is produced by single action potential of pacemaker activity may not be strong enough to induce GnRH release from the dense cored vesicles. However, it has been reported that such low frequency firing activity enables vasopressin release from rat neurohypophysis (Bondy *et al.*, 1987), substance P or thyrotropin releasing hormone release from rat ventral spinal cord (Iverfeldt *et al.*, 1989), and GnRH release from preganglionic C-neurons of the bullfrog (Peng and Horn, 1991). Furthermore, increase of firing frequency potentiated peptide release from these cells. Thus, it may be possible to think that the frequency of beating pacemaker activity of TN-GnRH neurons may affect the release of GnRH.

What then is the physiological significance of the modulation by sGnRH of pacemaker activity of TN-GnRH neurons? TN-GnRH neurons of the dwarf gourami make tight cell clusters with no intervening glial cells (Oka and Ichikawa, 1991; Oka and Matsushima, 1993; Oka, 1997), and the possibility of active exocytotic release from the cell body and its vicinity has also been suggested (Oka and Ichikawa, 1991). Other studies have shown that GnRH receptors are widely distributed throughout the brain (Jennes *et al.*, 1997; Stojilkovic *et al.*, 1994b). In addition, considerable overlap of the brain areas that contain GnRH-producing cells and GnRH receptor mRNA-expressing cells has been reported (Jennes *et al.*, 1996). Also, cultured hypothalamic GnRH neurons have GnRH receptors (Krsmanovic *et al.*, 1999). Furthermore, it has been reported using an *in vivo* analysis of multiunit activities in ovariectomized rats that GnRH injected in and around the median eminence is able to cause a population of GnRH neurons to fire synchronously (Hiruma and Kimura, 1995). From these observations and the present results, it is suggested that GnRH released from GnRH neurons facilitates the activities of their own (autocrine) and/or neighboring GnRH neurons (paracrine) and may cause synchronized positive feedback facilitation of multiple GnRH neurons. It is well known that in oxytocin neurons the release of oxytocin from single neuron into the brain environment stimulates its own activity and

thus further release (Freund-Mercier and Richard, 1984; Moos *et al.*, 1984). A similar effect has been reported for insulin-stimulated insulin release in pancreatic β cells (Aspinwall *et al.*, 1999). Therefore, this mechanism is probably common to all neurosecretory neurons or secretory cells, whose synchronized facilitation of firing leads to facilitated release.

Comparison with the autoregulation in the other putative GnRH neurons

Possible autoregulation mechanisms of TN-GnRH cells suggested in the present study are discussed here in relation to references that described studies on putative GnRH neurons. The concept of an ultrashort feedback mechanism in the control of neurosecretion was first suggested by Hyyppa *et al.* (1971) in studies on the control of FSH secretion. Similarly, it was revealed from *in vivo* (Valenca *et al.*, 1987) and *in vitro* studies (DePaolo *et al.*, 1987) that GnRH release from hypothalamic GnRH system was negatively autoregulated. It was postulated that negative feedback could be mediated via axo-dendritic/axo-somatic synapses on adjacent GnRH or other types of neurons (DePaolo *et al.*, 1987), and histological evidence for such connections has been reported (Leranth *et al.*, 1985; Witkin and Silverman, 1985).

However, the results using recording of multiunit activity (MUA) in the hypothalamus, which are considered to reflect the secretory activity of GnRH, have been very complicated. Hiruma and Kimura (1995) reported that intravenous injection of GnRH or microinjection of GnRH into the median eminence immediately evoked a MUA volley of the rat. However microinjection of GnRH into the medial preoptic area did not cause these effects. Moreover, intravenous or intracerebroventricular injections of GnRH did not affect MUA volleys of the rhesus monkeys (Kesner *et al.*, 1986; Ordog *et al.*, 1997). Unfortunately, the neuronal elements responsible for the MUA volley, i.e., whether the MUA volley represent the activity of GnRH neurons themselves or other neuronal elements, have not been determined. Therefore, it is difficult to directly compare the results of the present paper with the results of MUA studies.

Exposure of perfused GT1-7 cells to a GnRH agonist (analog) caused a transient elevation of GnRH release and subsequent suppression of the basal pulsatile secretion. During further continuous exposure to the agonist, the cells recovered from inhibition and exhibited infrequent but increasingly prominent peaks, with a net increase in GnRH release (Krsmanovic *et al.*, 1993). Recently, similar autoregulation of GnRH secretion has been reported in hypothalamic culture (Krsmanovic *et al.*, 1999). However, the time courses of such autoregulation are much slower than those of TN-GnRH neurons. The autoregulation of GnRH release in GT1-7 cell and hypothalamic culture takes dozens of minutes to several hours to occur. However, the biphasic autoregulation of pacemaker activity of TN-GnRH neurons took only a few minutes. The former has been suggested to

result from the internalization of receptor molecules and downregulation of the expression of GnRH receptor gene that was induced by GnRH (Park, 1998; Stojilkovic *et al.*, 1994a). On the other hand, the biphasic autoregulation of pacemaker activity of TN-GnRH neurons is considered to be due to the modulations of ionic channels induced by the activation of cell signaling pathways downstream of GnRH receptor activation. Thus, the two phenomena are considered to be based on quite different mechanisms.

Recently, Van Goor *et al.* (1999a, b) reported that the membrane excitability of GT1 cells was modulated by GnRH peptide. In this modulation of membrane excitability, they suggested that the activation of GnRH receptor induces $[Ca^{2+}]_i$ mobilization, and this $[Ca^{2+}]_i$ mobilization activates SK-type Ca^{2+} -activated K^+ channel and activates store-operated Ca^{2+} -channel, which transiently hyperpolarizes and then persistently depolarizes membrane potentials, respectively. They further suggested that this sustained membrane depolarization is explained by complex interplay of Na^+ channel, K^+ channel, store-operated Ca^{2+} channel, and L-type Ca^{2+} channel (Van Goor *et al.*, 2000). Furthermore, LeBeau *et al.* (2000) showed the contribution of cAMP-operated inward current to this modulation in GT1 cells. The mechanism of the transient decrease of firing frequency in TN-GnRH neurons (the present study) basically agrees well with these reports. However, detailed mechanism of the sustained increase of firing frequency seems to differ from each other. To further understand this mechanism, more detailed kinetic and pharmacological investigation of the control of pacemaker activities of TN-GnRH neurons are under way.

ACKNOWLEDGEMENTS

This research was supported by Grants-in-Aid from the Ministry of Education, Culture, Sports, Science and Technology (MEXT) of Japan to Y. Oka (No. 12440237), and the Sasakawa Scientific Research Grant from the Japan Science Society to H. Abe.

REFERENCES

- Abe H, Oka Y (1999) Characterization of K^+ currents underlying pacemaker potentials of fish gonadotropin-releasing hormone cells. *J Neurophysiol* 81: 643–653
- Abe H, Oka Y (2000) Modulation of pacemaker activity by salmon gonadotropin-releasing hormone (sGnRH) in terminal nerve (TN)-GnRH neurons. *J Neurophysiol* 83: 3196–3200
- Adams PRB, Brown DA (1980) Luteinizing hormone-releasing factor and muscarinic agonists act on the same voltage-sensitive K^+ -current in bullfrog sympathetic neurones. *Br J Pharmacol* 68: 353–355
- Aspinwall CA, Lakey JRT, Kennedy RT (1999) Insulin-stimulated insulin secretion in single pancreatic beta cells. *J Biol Chem* 274: 6360–6365
- Bondy CA, Gainer H, Russell JT (1987) Effects of stimulus frequency and potassium channel blockade on the secretion of vasopressin and oxytocin from the neurohypophysis. *Neuroendocrinol* 46: 258–267
- Bosma MM (1993) Ion channel properties and episodic activity in isolated immortalized gonadotropin-releasing hormone (GnRH) neurons. *J Memb Biol* 136: 85–96
- Bosma MM, Hille B (1992) Electrophysiological properties of a cell line of the gonadotrope lineage. *Endocrinology* 130: 3411–3420
- Bourinet E, Fournier J, Nargeot J, Charnet P (1992) Endogenous Xenopus-oocyte Ca-channels are regulated by protein kinases A and C. *FEBS Lett* 299: 5–9
- Costantin JL, Charles AC (1999) Spontaneous action potentials initiate rhythmic intercellular calcium waves in immortalized hypothalamic (GT1-1) neurons. *J Neurophysiol* 82: 429–435
- DePaolo LV, King RA, Carrillo AJ (1987) In vivo and in vitro examination of an autoregulatory mechanism for luteinizing hormone-releasing hormone. *Endocrinology* 120: 2561–2571
- Dolphin AC (1998) Mechanisms of modulation of voltage-dependent calcium channels by G proteins. *J Physiol* 506: 3–11
- Eisthen HL, Delay RJ, Wirsig-Wiechmann CR, Dionne VE (2000) Neuromodulatory effects of gonadotropin releasing hormone on olfactory receptor neurons. *J Neurosci* 20: 3947–3955
- Elmslie KS, Zhou W, Jones SW (1990) LHRH and GTP- γ S modify calcium current activation in bullfrog sympathetic neurons. *Neuron* 5: 75–80
- Freud-Mercier MJ, Richard P (1984) Electrophysiological evidence for facilitatory control of oxytocin neurons by oxytocin during suckling in the rats. *J Physiol* 352: 447–466
- Hales TG, Sanderson MJ, Charles AC (1994) GABA has excitatory actions on GnRH-secreting immortalized hypothalamic (GT1-7) neurons. *Neuroendocrinol* 59: 297–308
- Hille B (2001) *Ionic Channels of Excitable Membranes*, 3rd ed., Sinauer Associates, Sunderland
- Hille B (1994) Modulation of ion-channel function by G-protein-coupled receptors. *Trends Neurosci* 17: 531–536
- Hiruma H, Kimura F (1995) Luteinizing hormone-releasing hormone is a putative factor that causes LHRH neurons to fire synchronously in ovariectomized rats. *Neuroendocrinol* 61: 509–516
- Hyypya M, Motta M, Martini, L (1971) 'Ultrashort' feedback control of follicle-stimulating hormone-releasing factor secretion. *Neuroendocrinol* 42: 392–398
- Ishizaki M, Oka Y (2001) Amperometric recording of gonadotropin-releasing hormone release activity in the pituitary of the dwarf gourami (teleost) brain-pituitary slices. *Neurosci Lett* 299: 121–124
- Iverfeldt K, Serfözö P, Diaz Arnesto L, Bartfai T (1989) Differential release of coexisting neurotransmitters: frequency dependence of the efflux of substance P, thyrotropin releasing hormone and [3H] serotonin from tissue slices of rat ventral spinal cord. *Acta Physiol Scand* 137: 63–71
- Javors MA, King, TS, Chang X, Klein, NA, Schenken, RS (1995) Partial characterization of $K(+)$ -induced increase in $[Ca^{2+}]_{cyt}$ and GnRH release in GT1-7 neurons. *Brain Res* 694(1–2): 49–54
- Jennes L, McShane T, Brame B, Centers A (1996) Dynamic changes in gonadotropin releasing hormone receptor mRNA content in the mediobasal hypothalamus during the rat estrous cycle. *J Neuroendocrinol* 8: 275–281
- Jennes L, Centers A, Eyigor O (1997) GnRH receptors in the rat central nervous system. In "GnRH Neurons: Gene to Behavior", Brain Shuppan, Tokyo, pp 79–95
- Kesner JS, Kaufman JM, Wilson RC, Kuroda G, Knobil E (1986) On the short-loop feedback regulation of the hypothalamic luteinizing hormone releasing hormone 'pulse generator' in the rhesus monkey. *Neuroendocrinology* 42(2): 109–111
- Krsmanovic LZ, Stojilkovic SS, Merelli F, Dufour SM, Virmani MA, Catt KJ (1992) Calcium signaling and episodic secretion of gonadotropin-releasing hormone in hypothalamic neurons. *Proc Natl Acad Sci USA* 89: 8462–8466
- Krsmanovic LZ, Stojilkovic SS, Mertz LM, Tomic M, Catt KJ (1993) Expression of gonadotropin-releasing hormone receptors and

- autocrine regulation of neuropeptide release in immortalized hypothalamic neurons. *Proc Natl Acad Sci USA* 90: 3908–3912
- Krsmanovic LZ, Martinez-Fuentes AJ, Arora KK, Mores N, Navaro CE, Chen HC, Stojilkovic SS, Catt KJ (1999) Autocrine regulation of gonadotropin-releasing hormone secretion in cultured hypothalamic neurons. *Endocrinol* 140: 1423–1431
- Kusano K, Fueshko S, Gainer H, Wray S (1995) Electrical and synaptic properties of embryonic luteinizing hormone-releasing hormone neurons in explant cultures. *Proc Natl Acad Sci USA* 92: 3918–3922
- LeBeau AP, Van Goor F, Stojilkovic SS, Sherman A (2000) Modeling of membrane excitability in gonadotropin-releasing hormone-secreting hypothalamic neurons regulated by Ca^{2+} -mobilizing and adenylyl cyclase-coupled receptors. *J Neurosci* 15: 9290–9297
- Leranth C, Sugura LM, Palkovits M, MacLusky NJ, Shanabrough M, Naftolin F (1985) The LH-RH-containing neuronal network in the preoptic area of the rat: demonstration of LH-RH-containing nerve terminals in synaptic contact with LH-RH neurons. *Brain Res* 345: 332–336
- Meir A, Ginsburg S, Butkevich A, Kachalsky, SG, Kaiserman I, Ahdut R, Demigoren S, Rahamimoff R (1999) Ion channels in presynaptic nerve terminals and control of transmitter release. *Physiol Rev* 79: 1019–1088
- Moos F, Freund-Mercier MJ, Guerne JM, Stoeckel ME, Richard P (1984) Release of oxytocin and vasopressin by magnocellular nuclei *in vitro*: Specific facilitatory effect of oxytocin on its own release. *J Endocrinol* 102: 63–72
- Naor Z (1990) Signal transduction mechanisms of Ca^{2+} mobilizing hormones: The case of gonadotropin-releasing hormone. *Endocrine Rev* 11: 326–353
- Naor Z, Harris D, Shacham S (1998) Mechanism of GnRH receptor signaling: Combinatorial cross-talk of Ca^{2+} and protein kinase C. *Frontiers in Neuroendocrinol* 19: 1–19
- Nowycky MC (1991) Distinguishing between multiple calcium channel types. In "Molecular Neurobiology: A Practical Approach", IPL press, Oxford, pp 27–47
- Nowycky MC, Fox AP, Tsien RW (1985) Three types of neuronal calcium channel with different calcium agonist sensitivity. *Nature* 316: 440–443
- Oka Y, Abe H (2002) Physiology of GnRH neurons and modulation of their activities by GnRH. In: "Neuroplasticity, Development, and Steroid Hormone Action", CRC Press, Boca Raton, pp 191–203
- Oka Y, Ichikawa M (1991) Ultrastructure of the ganglion cells of the terminal nerve in the dwarf gourami (*Colisa lalia*). *J Comp Neurol* 304: 161–171
- Oka Y, Matsushima T (1993) Gonadotropin-releasing hormone (GnRH)-immunoreactive terminal nerve cells have intrinsic rhythmicity and project widely in the brain. *J Neurosci* 13: 2161–2176
- Oka Y (1995) Tetrodotoxin-resistant persistent Na^+ current underlying pacemaker potentials of fish gonadotropin-releasing hormone neurones. *J Physiol* 482: 1–6
- Oka Y (1996) Characterization of TTX-resistant persistent Na^+ current underlying pacemaker potentials of fish gonadotropin-releasing hormone (GnRH) neurons. *J Neurophysiol* 75: 2397–2404
- Oka Y (1997) GnRH neuronal system of fish brain as a model system for the study of peptidergic neuromodulation In "GnRH Neurons: Gene to Behavior", Brain Shuppan, Tokyo, pp 245–276
- Ordog T, Chen MD, Nishihara M, Connaughton MA, Goldsmith JR, Knobil E (1997) On the role of gonadotropin-releasing hormone (GnRH) in the operation of the GnRH pulse generator in the rhesus monkey. *Neuroendocrinology* 65(5): 307–313
- Parhar IS, Iwata M (1994) Gonadotropin-releasing hormone (GnRH) neurons project to growth hormone and somatotactin cells in the steel head trout. *Histochem* 102: 195–203
- Park, MK (1998) Molecular structure and expression divergence of the GnRH receptors In "Brain and Reproduction Evolution and Fitness of GnRH Neuronal System" (in Japanese), Gakkai Shuppan Center, Tokyo, pp 129–152
- Peng Y, Horn JP (1991) Continuous repetitive stimuli are more effective than bursts for evoking LHRH release in bullfrog sympathetic ganglia. *J Neurosci* 11: 85–95
- Sah P (1996) Ca^{2+} -activated K^+ currents in neurones: types, physiological roles and modulation. *Trends Neurosci* 19: 150–154
- Schwanzel-Fukuda M, Silverman AJ (1990) The nervous terminalis of the guinea pig: A new luteinizing hormone-releasing hormone (LHRH) neuronal system. *J Comp Neurol* 191: 213–225
- Skryma R, Mariot P, Le Bourhis X, Van Coppenolle F, Shuba Y, Vanden Abeele F, Legrand G, Humez S, Boilly B, Prevarskaya N (2000) Store depletion and store-operated Ca^{2+} current in human prostate cancer LNCaP cells: involvement in apoptosis. *J Physiol* 527: 71–83
- Spergel DJ, Catt KJ, Rojas E (1996) Immortalized GnRH neurons express large-conductance calcium-activated potassium channels. *Neuroendocrinol* 63: 101–111
- Stea A, Soong TW, Snutch TP (1995) Determinants of PKC-dependent modulation of a family of neuronal calcium channels. *Neuron* 15: 929–940
- Stojilkovic SS, Krsmanovic LZ, Spergel DJ, Catt KJ (1994a) Gonadotropin-releasing hormone neurons: Intrinsic pulsatility and receptor-mediated regulation. *Trends Endocrinol Metab* 5: 201–209
- Stojilkovic SS, Reinhart J, Catt KJ (1994b) Gonadotropin-releasing hormone receptors: Structure and signal transduction pathways. *Endocrine Rev* 15: 462–499
- Stojilkovic SS, Catt KJ (1995a) Expression and signal transduction pathways of gonadotropin-releasing hormone receptors. *Rec Prog in Hormone Res* 50: 161–205
- Stojilkovic SS, Catt KJ (1995b) Novel aspects of GnRH-induced intracellular signaling and secretion in pituitary gonadotrophs. *J Neuroendocrinol* 7: 739–757
- Swartz KJ (1993) Modulation of Ca^{2+} channels by protein kinase C in rat central and peripheral neurons: disruption of G protein-mediated inhibition. *Neuron* 11: 305–320
- Taylor CW, Broad LM (1998) Pharmacological analysis of intracellular Ca^{2+} signaling: problems and pitfalls. *Trends Pharm Sci* 19: 370–375
- Valenca MM, Johnston CA, Ching M, Negro-Vilar A (1987) Evidence for a negative ultrashort loop feedback mechanism operating on the luteinizing hormone-releasing hormone neuronal system. *Endocrinology* 121: 2256–2259
- Van Goor F, Krsmanovic LZ, Catt KJ, Stojilkovic SS (1999a) Coordinated regulation of gonadotropin-releasing hormone neuronal firing patterns by cytosolic calcium and store depletion. *Proc Natl Acad Sci USA* 96: 4101–4106
- Van Goor F, Krsmanovic LZ, Catt KJ, Stojilkovic SS. (1999b) Control of action potential-driven calcium influx in GT1 neurons by the activation status of sodium and calcium channels. *Mol Endocrinol* 13: 587–603
- Van Goor F, LeBraun AP, Krsmanovic LZ, Sherman A, Catt KJ, Stojilkovic SS (2000) Amplitude-dependent spike-broadening and enhanced Ca^{2+} signaling in GnRH-secreting neurons. *Biophys J* 79: 1310–1323
- Witkin JW, Silverman AJ (1985) Synaptology of luteinizing hormone-releasing hormone neurons in rat preoptic area. *Peptides* 6: 263–271
- Yamamoto N, Oka Y, Amano M, Aida K, Hasegawa Y, Kawashima S (1995) Multiple gonadotropin-releasing hormone (GnRH)-immunoreactive systems in the brain of the dwarf gourami, *Colisa lalia*: immunohistochemistry and radioimmunoassay. *J*

- Comp Neurol 355: 354–368
- Yang J, Tsien RW (1993) Enhancement of N- and L- type calcium channel currents by protein kinase C in frog sympathetic neurons. *Neuron* 10: 127–136
- Zamponi GW, Bourinet E, Neison D, Nargeot J, Snutch TP (1997) Crosstalk between G proteins and protein kinase C mediated by the calcium channel α 1 subunit. *Nature* 385: 442–446
- Zhu Y, Ikeda SR (1994) Modulation of Ca^{2+} -channel currents by protein kinase C in adult at sympathetic neurons. *J Neurophysiol* 72: 1549–1560

(Received October 25, 2001 / Accepted November 25, 2001)