

Arachnids (Arachnida: Araneae, Mesostigmata, Pseudoscorpiones) from tropical greenhouses at Rotterdam Zoo (the Netherlands), including a pholcid spider new to Europe

Authors: Bloem, Geron J., and Noordijk, Jinze

Source: Arachnologische Mitteilungen: Arachnology Letters, 61(1) : 36-44

Published By: Arachnologische Gesellschaft e.V.

URL: <https://doi.org/10.30963/aramit6106>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Arachnids (Arachnida: Araneae, Mesostigmata, Pseudoscorpiones) from tropical greenhouses at Rotterdam Zoo (the Netherlands), including a pholcid spider new to Europe

Geron J. Bloem & Jinze Noordijk



doi: 10.30963/aramit6106

Abstract. An inventory of arachnids was conducted in three tropical greenhouses of Rotterdam Zoo, the Netherlands. This was part of a survey in which pitfall traps, banana baits, litter samples, hand searching and beating were used to map arthropod diversity. A total of twenty spider species, one mite taxon and one pseudoscorpion species were collected. Fourteen specimens were used for COI-barcoding, which yielded two of the listed species. Four spider species were recorded for the first time for the Netherlands: *Belisana ambengan* Huber, 2005, *Heteroonops spinimanus* (Simon, 1892), *Nesticella mogera* (Yaginuma, 1972) and *Triaeris stenaspis* Simon, 1891, the first of which is also a new record for Europe and appears to be an extremely rare pholcid species. From the species found, thirteen are non-native to the Netherlands and four of these are considered as expansive, since they have outdoor populations as well. Hand searching yielded the highest number of species, compared to the other four sampling techniques. A species saturation curve shows that in all likelihood not all species present in the surveyed greenhouses have been found during this survey.

Keywords: alien species, *Belisana ambengan*, faunistics, new records, Nesticidae, Oonopidae, Pholcidae

Zusammenfassung. Spinnentiere aus tropischen Gewächshäusern des Zoos in Rotterdam (Niederlande), mit einer Zitterspinne neu für Europa (Arachnida: Araneae, Mesostigmata, Pseudoscorpiones). In drei tropischen Gewächshäusern des Zoos in Rotterdam wurden Spinnentieruntersuchungen durchgeführt. Dies gehörte zu einem Programm in dem mit Bodenfallen, Bananenködern, Streuproben, Handfängen und Klopfproben die Arthropodenvielfalt kartiert wurde. Insgesamt wurden 20 Spinnenarten, ein Milbentaxon und eine Pseudoskorpionart erfasst. 14 Tiere wurde zum COI-Barcoding verwendet, was zwei der aufgeführten Arten ergab. Vier Spinnenarten wurden erstmals für die Niederlande nachgewiesen: *Belisana ambengan* Huber, 2005, *Heteroonops spinimanus* (Simon, 1892), *Nesticella mogera* (Yaginuma, 1972) und *Triaeris stenaspis* Simon, 1891, die erstgenannte ist auch ein Neunachweis für Europa und scheint eine sehr seltene Zitterspinnenart zu sein. Dreizehn der Arten sind nicht in den Niederlanden heimisch und vier davon werden als expansiv eingestuft, weil es von ihnen auch Freilandpopulationen gibt. Mit Handfängen wurden, im Vergleich zu den vier anderen Sammelmethoden, die meisten Arten erfasst. Eine Artensättigungskurve zeigt, dass wahrscheinlich noch nicht alle Arten erfasst wurden, die in den untersuchten Gewächshäusern vorkommen.

Greenhouses can provide a suitable environment for diverse groups of invertebrates. Each tropical greenhouse has its own characteristics caused by various factors, e.g. climate control, management, composition and origins of plant and soil material. Depending on these factors, complex artificial ecosystems can be formed to a certain extent. Greenhouses often import a large number of (tropical) plant species from remote sources, almost always with accompanying soil, resulting in the introduction of species living in or on these substrates. Some of these are able to settle in the greenhouse. Certain alien species that thrive in greenhouses might even become indoor pests or expansive if they manage to establish themselves outdoors (Wang et al. 2015). Additionally, synanthropic species may colonise or be introduced by transport from (nearby) man-made biotopes. Some native species living in natural habitats occasionally enter greenhouses as well (Kielhorn 2008).

A survey of arachnids was carried out in three tropical greenhouses in Rotterdam Zoo. This marks as the first extensive arachnid survey in Dutch tropical greenhouses since 1949 (van der Hammen 1949, 1969a). At that time, Leendert van der Hammen and fourteen other collectors surveyed eleven greenhouses in the midwest and centre of the Netherlands, including Rotterdam Zoo, by hand, searching over a seven-year period. Most greenhouses were, however, visited only once. Thirty-two arachnid species were collected, with only one species, *Zygiella x-notata* (Clerck, 1757), sampled at

Rotterdam Zoo (van der Hammen 1949). Currently, only a handful of studies on arachnid faunas of European tropical greenhouses exist, e.g. the Eden Project in England (Snazell & Smithers 2007), several botanical and zoological gardens in the German states Berlin and Brandenburg (Kielhorn 2008, 2009, 2016), the Botanical Garden of the University of Debrecen in Hungary (Pfliegler 2014), the Botanical Garden of the PJ Šafárik University in Slovakia (Šestáková et al. 2017) and multiple zoological gardens in the Czech Republic (Hula & Pešan 2018).

The aim of the present paper is to give insight into the arachnid fauna of the tropical greenhouses at Rotterdam Zoo and document species found in the Netherlands for the first time. Additionally, a graph is given to illustrate which method proved most effective in collecting arachnids in tropical greenhouses.

Material and methods

Locations

This study was carried out in three tropical greenhouses located at Rotterdam Zoo: Amazonica, Victoria Serre and the Kwekerij.

Amazonica (51.9280°N, 4.4473°E, -1 m alt., Fig. 1) is a large circular greenhouse designed as a butterfly garden, opened in 2013. It has a diameter of 60 m and its highest point is 12 m. A total of 318 plant species are growing in open soil in Amazonica, mostly of South American origin. The average temperature is 25–28°C during daytime and goes down to 18–23°C at night. During warm summer days, the average temperature can be as high as 32–34°C.

The Victoria Serre (51.9260°N, 4.4529°E, -1 m alt., Fig. 2) is a large aviary built in 1945. It consists of three distinct but connected buildings, each hosting a different array of

Geron J. BLOEM, HAS University of Applied Sciences, Onderwijsboulevard 221, 5223 DE 's-Hertogenbosch, The Netherlands; E-mail: g.bloem@live.nl
Jinze NOORDIJK, European Invertebrate Survey – the Netherlands / Naturalis Biodiversity Center, Darwinweg 2, 2300 RA Leiden, The Netherlands; E-mail: jinze.noordijk@naturalis.nl

Academic editor: Petr Dolejš

submitted: 31.1.2020, accepted: 2.3.2021, online: 6.4.2021

Asian bird species. In the first and second part, the vegetation is grown in patches of soil alongside a footpath. In the third part, the vegetation grows around a 10 m diameter pool containing giant water lilies. The diameter of the circular-shaped third part is 24 m and its highest point is circa 8 m. A total of 61 plant species are growing in open soil. The average temperature is 25°C during daytime and 20°C at night.


The Kwekerij ('nursery') (51.9243°N, 4.4544°E, -1 m alt., Fig. 3) is a complex consisting of thirteen small greenhouses built in 1993, which function as nurseries for tropical plants. In contrast to the other greenhouses, it is not open to the public. The arthropod survey was carried out in four greenhouses with heights ranging from 9.8 to 11.6 m. The majority of the planting consists of a mixture of potted plants that can also be found in Amazonica and Victoria Serre. In one of the sampled greenhouses, several plants are (permanently) grown in an elongated patch of soil. Most of the greenhouses have the same temperature fluctuations as in Amazonica.

Sampling methods

The arachnids were surveyed in two separate periods: 4. Jun. – 10. Jul. (week 23–28) and 27. Aug. – 19. Sep. (week 35–38) 2018. Samples were collected each week using five methods. (i) Pitfall traps, composed of a 500 ml yoghurt can filled with circa 50 ml ethylene glycol solution. A total of five pitfall traps were placed per greenhouse, in the most dissimilar habitats possible. The traps were protected by a plastic lid supported by four clout nails. The pitfalls were emptied once a week. (ii) Banana baits, which were obviously primarily used to attract arthropods other than Arachnida, e.g. Drosophilidae and certain Coleoptera. A half peeled ripening banana was placed inside a plastic container on the ground, with 10 mm mesh covering the entrance, at two sites per greenhouse complex. The bananas were sampled once a week. (iii) Leaf litter samples, which consisted of sieving the top soil layer (i.e. 2 cm) within an area of 50 × 39 cm. The samples were sifted through a beetle sieve with a mesh size of 10 mm. The samples were taken weekly at different locations, once per greenhouse complex. In the Kwekerij, the samples were always taken at the only location with permanent vegetation. (iv) Hand searching, which included manually looking for arachnids and moving objects such as rocks, wood and litter. (v) Beating, consisted of shaking branches by hand over a sweep net. A few web-building spider species (i.e. *Parasteatoda tepidariorum* (C. L. Koch, 1841), *Pholcus phalangioides* (Fuesslin, 1775) and *Uloborus plumipes* Lucas, 1846) were counted instead of sampled, because of their unmistakable identification and common occurrence. At the start of the inventory, specimens of these species were collected to verify the identification. The exact sampling locations of hand searching and beating were randomised, with the aim of ultimately surveying as many locations as possible. Hand searching and beating were performed once a week, within a fixed period of time (i.e. 60 minutes). All samples and traps were evenly spread across each greenhouse.

Identification and COI-barcoding

All specimens were collected by transferring them into vials containing 96% ethanol. Identification of spiders was primarily done following Nentwig et al. (2019). The pseudoscorpions were identified with Legg & Farr-Cox (2016). Some specimens were photographed using a Zeiss Axioskop



Figs 1-3: Impression of the three sampling locations at Rotterdam Zoo. **1.** Amazonica; **2.** Victoria Serre; **3.** the Kwekerij

microscope equipped with an AxioCamMRC5 digital camera, creating stacked images with a great depth of field. Male specimens of *Belisana ambengan* were sent to Bernhard Huber (Zoologisches Forschungsmuseum Alexander Koenig, Bonn, Germany) for identification.

From the collected specimens, fourteen were subsampled for molecular analysis of the cytochrome c oxidase subunit I gene (COI) (Tab. 1). DNA extraction and amplification were performed at the Naturalis DNA Barcoding Facility. DNA primers LCO1490 and HCO2198 were used to amplify a 658-bp region of the mitochondrial COI gene. The resulting PCR products were sequenced at BaseClear using the Sanger sequencing method (for standardised protocols on extraction, amplification and sequencing, see Ivanova et al. 2009). The COI-sequences were matched against the BOLD database

with the basic local alignment search tool (BLAST). Sequences with at least 99.0% similarity were declared a match (Hebert et al. 2003). The voucher specimens and parts of the collected material were deposited in the spider collection (RMNH.ARA) of Naturalis Biodiversity Center. Another part of the material is stored in the private collection of the first author.

Data analysis

In order to evaluate the course of sampling, a growth curve illustrating the cumulative percentage of additional arachnid and arthropod species was used. To compare the differences in species richness between the sampling methods, a univariate analysis of variance was performed. Differences were considered to be significant at $p < 0.05$ values.

Tab. 1: List of arachnids recorded in the greenhouses at Rotterdam Zoo. Abbreviations: * – species recorded in the Netherlands for the first time; A – alien species (established) in the Netherlands; BB – banana baits; BE – beating; E – expansive alien species in the Netherlands; HS – hand searching; j – juvenile; LS – litter sieving; N – native species in the Netherlands; PT – pitfall traps; s – subadult

| Order/Family/Species | Status | Methods of collection | | | | |
|--|--------|-----------------------|---------------------|--------------------------------|----------------------------------|-----------------------------------|
| | | BB | BE | HS | LS | PT |
| Araneae | | | | | | |
| Agelenidae | | | | | | |
| 1 <i>Eratigena atrica</i> (C. L. Koch, 1843) | N | | | 1 ♂ | | |
| Araneidae | | | | | | |
| 2 <i>Araneus diadematus</i> Clerck, 1757 | N | | | 1 j | | |
| 3 <i>Nuctenea umbratica</i> (Clerck, 1757) | N | | | 1 j | | |
| Dysderidae | | | | | | |
| 4 <i>Dysdera cf. crocata</i> C. L. Koch, 1838 | | | | | | 1 s♀, 3 jj |
| Linyphiidae | | | | | | |
| 5 <i>Erigone atra</i> Blackwall, 1833 | N | | | | 2 ♀♀ | |
| 6 <i>Mermessus trilobatus</i> (Emerton, 1882) | E | 1 ♀ | | | | 2 s♀♀ |
| 7 <i>Ostearius melanopygius</i> (O. Pickard-Cambridge, 1880) | E | | | 1 ♀ | | |
| Nesticidae | | | | | | |
| 8 * <i>Nesticella mogera</i> (Yaginuma, 1972) | A | | | | | 2 ♂♂, 3 jj |
| Oonopidae | | | | | | |
| 9 * <i>Heteroonops spinimanus</i> (Simon, 1892) | A | | | | 2 ♀ | |
| 10 * <i>Triaeris stenaspis</i> Simon, 1891 | A | | | | | 7 ♀♀, 1 j |
| Pholcidae | | | | | | |
| 11 * <i>Belisana ambengan</i> Huber, 2005 | A | | 2 ♂♂, 1 s♀, 4 jj | | | |
| 12 <i>Pholcus phalangioides</i> (Fuesslin, 1775) | A | | | 119 (observations), 1 ♂ | | |
| 13 <i>Psilochorus simoni</i> (Berland, 1911) | A | | | 1 ♂ | 1 s♀ | 1 ♀ |
| Salticidae | | | | | | |
| 14 <i>Hasarius adansoni</i> (Audouin, 1826) | A | 1 ♂, 1 ♀, 1 j | | 1 ♂, 1 ♀, 4 s♀♀, 2 jj | 1 ♂, 2 s♂♂, 3 s♀♀, 12 jj | 2 ♂♂, 5 s♂♂, 1 ♀, 9 s♀♀, 21 jj |
| Segestriidae | | | | | | |
| 15 <i>Segestria bavaria</i> C. L. Koch, 1843 | N | | | 1 ♀ | | |
| Theridiidae | | | | | | |
| 16 <i>Coleosoma floridanum</i> Banks, 1900 | A | | | 1 ♂, 1 s♂, 1 ♀, 10 s♀♀, 1 j | 6 ♂♂, 1 s♂, 6 ♀♀, 14 s♀♀, 1 j | 4 s♀♀, 2 jj |
| 17 <i>Parasteatoda tepidariorum</i> (C. L. Koch, 1841) | E | | 1 ♀ | 42 (observations), 2 ♀ | 1 j | 1 ♂ |
| 18 <i>Steatoda grossa</i> (C. L. Koch, 1838) | E | | 1 s♀ | 1 ♀, 1 j | | 2 ♂♂ |
| Uloboridae | | | | | | |
| 19 <i>Uloborus plumipes</i> Lucas, 1846 | A | | 3 jj | 407 (observations) | | |
| Zodariidae | | | | | | |
| 20 <i>Zodarion italicum</i> (Canestrini, 1868) | N | | | | 2 jj | 3 jj |
| Mesostigmata | | | | | | |
| Phytoseiidae | | | | | | |
| 1 <i>Typhlodromini</i> gen. sp. | | | | | 3 ex. | 3 ex. |
| Pseudoscorpiones | | | | | | |
| Chthoniidae | | | | | | |
| 1 <i>Ephippiochthonius tetrachelatus</i> (Preysslner, 1790) | N | | | | 2 ex. | 1 ex. |
| Total number of different taxa | | 2 | 4 | 12 | 10 | 12 |
| Total number of specimens | | 4 | 11 | 601 | 60 | 74 |

Tab. 2: List of the sequenced specimens that could be assigned to a Barcode Index Number (BIN) in the BOLD database, which verified the identification

| Order/Family/Species | Sequenced specimens | Voucher ID | Assigned BIN | % Match with BIN | GenBank Accession |
|--|---------------------|--------------------------------|--------------|------------------|------------------------------|
| Araneae | | | | | |
| Nesticidae | | | | | |
| <i>Nesticella mogera</i> | 1 ♂, 2 jj | RMNH.5105512, 5105515, 5105516 | BOLD:ACQ6894 | 99.02–99.35 | MW664906, MW664907, MW664908 |
| Oonopidae | | | | | |
| <i>Heteroonops spinimanus</i> | 1 ♀ | RMNH.5105513 | BOLD:AAN6310 | 99.81 | MW664913 |
| Salticidae | | | | | |
| <i>Hasarius adansoni</i> | 1 ♂ | RMNH.5105521 | BOLD:AAW0165 | 100 | MW664912 |
| Theridiidae | | | | | |
| <i>Coleosoma floridanum</i> | 1 ♂, 1 ♀ | RMNH.5105519, 5105520 | BOLD:AAV1727 | 99.82–100 | MW664909, MW664910 |
| Zodariidae | | | | | |
| <i>Zodarion italicum</i> | 1 j | RMNH.5105524 | BOLD:ACX0947 | 100 | MW664911 |
| Pseudoscorpiones | | | | | |
| Chthoniidae | | | | | |
| <i>Ephippiochthonius tetrachelatus</i> | 1 ex | RMNH.5105525 | BOLD:AAJ0945 | 100 | MW664914 |

Results

Twenty-two taxa were identified: twenty spider species, one mite taxon and one pseudoscorpion species (Tab. 1). Nine specimens could be identified by matching their COI-sequence against the BOLD database (Tab. 2). The majority of the species recorded in the greenhouses are alien to the Netherlands, the remaining ones are (semi)synanthropic.

Spiders (Araneae)

In total, twenty taxa (734 ex.) from twelve families were recorded (Tab. 1). *Uloborus plumipes* was the most dominant species overall, but completely absent from the Victoria Serre. Other very common species included *Pholcus phalangioides*, *Hasarius adansoni* (Audouin, 1826), *Parasteatoda tepidariorum* and *Coleosoma floridanum* Banks, 1900. Four species, *Nesticella mogera*, *Heteroonops spinimanus*, *Triaeris stenaspis* and *Belisana ambengan*, represent new records for the Netherlands. These and several other remarkable species are discussed in detail below.

Nesticidae

Nesticella mogera (Fig. 4)

Material examined. 4.–11. Jun. 2018 – 1 ♂, pitfall trap, Kwekerij; 25. Jun. – 2. Jul. 2018 – 3 jj, pitfall trap, Kwekerij (RMNH.5105515, 5105516, 5105517); 10.–18. Sep. 2018 – 1 ♂, pitfall trap, Amazonica (RMNH.5105512).

Identification. Nentwig et al. (2019). The identification of two out of three identical juvenile specimens found in Kwekerij was confirmed by COI barcoding.

The specimens sampled at Rotterdam Zoo represent the first record of this species in the Netherlands. In Europe, *N. mogera* was previously found in several heated buildings in Great Britain (Snazell & Smithers 2007), Denmark (Lissner & Scharff 2021), Germany (Kielhorn 2009, Reiser 2013), Poland (Bielak-Bielecki & Rozwałka 2011), Finland (Koponen et al. 2016), Italy (Pantini et al. 2020), Hungary (Pfliegler 2014) and the Czech Republic (Růžička 2016). The presence of this East-Asian troglophile species was expected, because the specimen collected in 2009 in Poland was found at a building-garden centre in Lublin in potted plants deriving



Fig. 4: Habitus of adult male *Nesticella mogera* from the Amazonica greenhouse, 10.–18. Sep. 2018; body length: 2.1 mm

from the Netherlands (Bielak-Bielecki & Rozwałka 2011). This suggests that *N. mogera* probably has been present in the Netherlands since at least 2009.

Oonopidae

Heteroonops spinimanus (Fig. 5)

Material examined. 11. Sep. 2018 – 1 ♀, litter sieving, Amazonica; 19. Sep. 2018 – 1 ♀, litter sieving, Amazonica (RMNH.5105513).

Identification. The identity of this species was verified by analysis of the COI gene of one female. Alternatively, females are distinguished from other nonscutate oonopids by the characteristic spination of the palp (Saaristo 2001).

The current survey produced the first record of this species for the Netherlands. *Heteroonops spinimanus* has been sampled from greenhouses in Germany (Kielhorn 2008) and the



Fig. 5: Adult female *Heteroonops spinimanus* from the Amazonica greenhouse, 19. Sep. 2018; body length: 1.8 mm

Czech Republic (Hula & Pešan 2018) as well. It is a Central-American litter-dwelling species, hence its presence in the litter sieving samples at Rotterdam Zoo (Pfeiffer 1996, Kielhorn 2008). Males are either rare or extremely short-lived, and were initially assigned to a different species (Platnick & Dupérré 2009). It is nevertheless possible that at least some of the introduced populations consist of parthenogenetic individuals (Platnick & Dupérré 2009).

Triaeris stenaspis (Fig. 6)

Material examined. 25. Jun. – 2. Jul. 2018 – 1 ♀, 1 j, pitfall trap, Kwekerij (RMNH.5105518); 2.–9. Jul. 2018 – 1 ♀, pitfall trap, Kwekerij; 27. Aug. – 3. Sep. 2018 – 1 ♀, pitfall trap, Victoria Serre; 3.–10. Sep. 2018 – 1 ♀, pitfall trap, Kwekerij; 3.–10. Sep. 2018 – 3 ♀♀, pitfall trap, Victoria Serre (RMNH.5105514).

Identification. Nentwig et al. (2019). *Triaeris stenaspis*, easily recognizable by the characteristic shape of its large dorsal scutum and the relatively small ventral scutum on the opisthosoma, is a parthenogenetic species (Korenko et al. 2009).

The current arachnid survey marks the first time that specimens of *T. stenaspis* were collected from the Netherlands. This oonopid spider of African origin was to be expected, as it was collected from several other greenhouses in Europe (Korenko et al. 2007, Pfliegler 2014, Rozwałka et al. 2016,



Fig. 6: Habitus of adult female *Triaeris stenaspis* from the Victoria Serre greenhouse, 27. Aug. – 3. Sep. 2018; body length: 1.7 mm

Šestáková et al. 2017, Rembold et al. 2020, Lissner & Scharff 2021). Furthermore, *T. stenaspis* was recently recorded at a tropical butterfly house (in Whipsnade Zoo, England) in which the trees and shrubs all originate from the Netherlands (Telfer 2020). Because of its synanthropic and parthenogenetic nature, it can easily invade greenhouses.

Pholcidae

Belisana ambengan (Figs 7-8)

Material examined. 9. Jul. 2018 – 1 j, beating, Kwekerij; 11. Sep. 2018 – 2 ♂♂, 1 s♀, 1 j, beating, Amazonica; 19. Sep. 2018 – 2 jj, beating, Victoria Serre (RMNH.ARA.18236).

Identification. These specimens were identified by Bernhard Huber as *Belisana ambengan*, based on his own first description (Huber 2005).

This find is surprising, because it represents the first record of a *Belisana* species in Europe and the species was solely known from its holotype (a male) and a prosoma of a female. Even more surprisingly, one male and one female were also found by the second author in a tropical greenhouse of Wildlands Adventure Zoo (Emmen, province of Drenthe) on 2. Jul. 2019 and an additional female was collected a few days later on 6. Jul. 2019 in the same greenhouse by D. Sies (all coll. Naturalis Biodiversity Center). The male from Emmen was used to make Figs 6-7, since the specimens from Rotterdam were already in the collection of B. Huber (Bonn, Germany). The type material of *B. ambengan* was collected in a secondary forest in Ambengan, Bali, Indonesia (Huber 2005). The record of this species at Rotterdam Zoo suggests that it might be more widely distributed in Indonesia than its type locality, since the chance of an isolated population to translocate to the Netherlands is negligibly small. A large part of the tropical plants at Rotterdam Zoo is imported from Indonesia.



Fig. 7: Habitus of adult male *Belisana ambengan* from Wildlands Adventure Zoo Emmen, 2. Jul. 2019; body length: 1.3 mm




Fig. 8: Right male palp in retrolateral view of *Belisana ambengan* from Wildlands Adventure Zoo Emmen, 2. Jul. 2019. Hairs not drawn and bulbapophysis is bent inwards and not visible, see Huber (2005) for its morphology

Theridiidae

Coleosoma floridanum

Material examined. 4.–11. Jun. 2018 – 1 s♀, pitfall trap, Victoria Serre; 11.–18. Jun. 2018 – 2 s♀, pitfall trap, Victoria Serre; 2.–9. Jul. 2018 – 1 s♀, 2 jj; 3. Jul. 2018 – 1 ♀, 1 ♂, litter sieving, Victoria Serre (RMNH.5105519, 5105520); 19. Sep. 2018 – 1 ♀, 1 ♂, litter sieving, Victoria Serre (RMNH.ARA.18233). The remaining specimens mentioned in Tab. 1 were collected at both Amazonica and Victoria Serre in nearly all sampling weeks, but not stored in a collection.

Identification. Nentwig et al. (2019) and COI barcoding.

Females of this Pantropical species carrying egg sacs were a common sight in the greenhouses at Rotterdam Zoo. Despite its common occurrence at the Rotterdam greenhouses, the presence of *C. floridanum* was unnoticed in the Netherlands since their first and only previous record by van Helsdingen (1995), probably due to their small size. In 2019 it was also found by the second author in a tropical greenhouse of Wildlands Adventure Zoo (Emmen, province of Drenthe) and it is suggested that this spider is more common than previously suspected.

Uloboridae

Uloborus plumipes

Material examined. 26. Jun. 2018 – 3 jj, beating, Kwekerij. Due to their high visibility and easily recognizable traits, specimens belonging to this species were counted instead of sampled during hand searching.

Identification. Nentwig et al. (2019). Their high visibility obviously accounted for the relatively high numbers of records.

Uloborus plumipes was the most abundant arachnid species from Amazonica and Kwekerij. In fact, it is a very widely distributed species in tropical and commercial greenhouses and garden centres in the Netherlands (Noordijk et al. 2018). It was previously stated that this species might be parthenogenetic in urban areas, but Oxford (2011) convincingly contradicted this statement. Its notable absence from Victoria Serre might be attributed to limiting, site-specific factors, e.g. use or avoidance of certain pesticides (Suvák 2013).

Zodariidae

Zodarion italicum

Material examined. 11. Jun. 2018 – 1 j, litter sieving, Kwekerij (RMNH.5105524); 11.–18. Jun. 2018 – 1 j, pitfall trap, Victoria Serre; 19. Jun. 2018 – 1 j, litter sieving, Kwekerij;

3.–10. Sep. 2018 – 1 j, pitfall trap, Kwekerij; 10.–18. Sep. 2018 – 1 j, pitfall trap, Amazonica.

Identification. As no adult specimens were collected, the identity of *Z. italicum* was determined by analysing the COI gene of one juvenile.

The presence of this native myrmecophagous species seemed unusual, because it was previously solely known from a natural site in the far south of the country (province of Limburg) (van Helsdingen 2018). *Zodarion italicum* is a South European species extending its range to the north, and has been recorded in greenhouses at the Botanic Garden Berlin-Dahlem and other places with anthropogenic habitats like railway stations and airports in Western and Central Europe (Malten et al. 2005, Pekár et al. 2005, Kielhorn 2008). It is therefore highly likely that the population at Rotterdam Zoo originates from other greenhouses or urban transport areas rather than the surrounding natural habitat.

Mesostigmata

Phytoseiidae

Typhlodrominae

Typhlodromini

Material examined. 26. Jun. 2018 – 2 ex., litter sieving, Victoria Serre; 25. Jun. – 2. Jul. 2018 – 2 ex., pitfall trap, Victoria Serre; 17. Sep. 2018 – 1 ex., litter sieving, Victoria Serre; 10.–18. Sep. 2018 – 1 ex., pitfall trap, Victoria Serre.

Identification. Six morphologically identical mites were sampled at Victoria Serre. Two of these specimens were sent to Henk Siepel (Radboud University, Nijmegen), who identified them as the tribus Typhlodromini (Tab. 1). No decisive answer could be given about what genus or species the specimens belong to, but they are likely the same species.

The number of mites is remarkably low. Phytophagous species were absent in the samples. In most greenhouses the numbers of phytophagous mites are high, regularly at such levels that it urges for pest control (Gerson & Weintraub 2012). Since herbivorous mites are absent, the predatory mites found at Rotterdam Zoo probably prey upon collembolans (Petrova et al. 2004).

Pseudoscorpiones

Only one, native pseudoscorpion species (3 ex.) was found inside the greenhouses (Tab. 1).

Chthoniidae

Ephippiochthonius tetrachelatus

Material examined. 4.–11. Jun. 2018 – 1 ♀, pitfall trap, Victoria Serre; 19. Jun. 2018 – 1 ♀, litter sieving, Victoria Serre (RMNH.5105525); 25. Jun. – 2. Jul. 2018 – 1 ♀, pitfall trap, Victoria Serre.

Identification. Legg & Farr-Cox (2016) and COI barcoding.

This pseudoscorpion species is native to Europe and hemisynanthropic, although its association with anthropogenic habitats differs mainly by latitude (Štáhlavský 2001). In the Netherlands, it can be found at a variety of dry and weakly humid locations and is also common in greenhouses, foraging under stones, logs and in leaf-litter (van der Hammen 1969b). In Scandinavia and northeast Poland, however, it lives almost exclusively in heated buildings or other synanthropic habitats, while in the Mediterranean it occurs widely independent of soil humidity (Štáhlavský 2001). *Ephippiochthonius tetrache-*

latus is frequently translocated to remote localities far from its native Western Palearctic and Nearctic range (Muchmore 2000). It is therefore suggested that the population of this species in Rotterdam Zoo, like *Zodarion italicum*, originates from other (European) greenhouses rather than the outside surroundings. Recently, Kaňuchová et al. (2016) also found it in many compost heaps, which could possibly serve as a means of transport between the greenhouses.

General evaluation

The growth of the cumulative percentage of additional species slightly decreased over time in both arachnid and arthropod species (Fig. 9). Although the last survey week did not yield any additional arachnid species, the number of additional arthropod species (e.g. Diptera, Isopoda, Myriapoda) slightly increased. The arachnid species curve seems saturated, but the presumed asymptote is reached only in the last survey week. However, the arthropod species curve does not reach a clear saturation point.

The mean species richness per sample was found to significantly differ between the sampling methods. Out of the five collecting methods, hand searching yielded the significantly highest mean number of arachnid species per sample, followed by both pitfall trapping and litter sieving, while both beating and the banana baits had the lowest mean species




Fig. 9: The growth of the cumulative percentage of additional species over time.




Fig. 10: The mean arachnid species richness per sample per sampling method, including SE-bars and statistical significance markings (with a level set at 0.05)

richness (Fig. 10). Hand searching yielded four species not caught by the other methods, while the pitfall traps and the litter sieving resulted both in two unique and beating in one unique species.

Discussion

Impact on the natural environment

A subset of the alien species might be able to thrive outdoors. *Mermessus trilobatus* (Emerton, 1882) established successful populations in a wide variety of open natural habitats in the Netherlands and other European countries (van Helsdingen 2009, Holec et al. 2012). *Ostearius melanopygius* (O. Pickard-Cambridge, 1880) is frequently found in manure heaps, and also in natural areas (Roberts 1998, Nentwig et al. 2019). *Parasteatoda tepidariorum* and *Steatoda grossa* (C. L. Koch, 1838) are regularly found on outer walls of artificial structures in the Netherlands (pers. obs.). Apparently, these species are able to survive outside of buildings in the European temperate zone. All other alien species from Tab. 1 remain very strongly associated with heated buildings in the current Dutch climate, although in 2005 one stray individual of *Hasarius adansoni* was found in debris in a riverine flood area (van Helsdingen 2009). Their impact on natural habitats is therefore now insignificant, but it remains to be seen which species will protrude to the outdoors due to the increasing warming of the climate.

Species to be expected

The sampling stopped at the point when no further species were collected (Fig. 9): the presumed asymptote. However, Gotelli & Colwell (2011) state that the asymptote of a growth curve can be reached only when at least 20 samples are taken. Moreover, future imports of plants and accompanying soil can introduce additional arachnid species to Rotterdam Zoo. Therefore, it is expected that a subsequent survey will confirm the presence of additional species.

The composition of species found at Rotterdam Zoo shows overlap with other surveys of arachnid fauna in greenhouses (Snazell & Smithers 2007, Kielhorn 2008, Pfliegler 2014, Šestáková et al. 2017, Hula & Pešan 2018). In most studies, the dominant species is either *Parasteatoda tepidariorum* or *Pholcus phalangioides*, while *Uloborus plumipes* is the most ubiquitous species at Rotterdam Zoo. However, the unexpected discovery of a new pholcid spider for Europe at two zoos in the Netherlands is remarkable. Another similar example is the recent discovery of a small pholcid in a Dutch tropical greenhouse, *Spermophora kerinci* Huber, 2005 (Noordijk 2020); this species is further only known from Indonesian rainforests and a tropical greenhouse in England and Germany (Snazell & Smithers 2007, Kielhorn 2009). It is therefore suggested that arachnid communities in European tropical greenhouses are interconnected, probably due to similar importing routes for the planting material or due to exchange of plants or other products. For this reason, it is expected that some of the additional alien species sampled at other tropical greenhouses in Europe will be recorded in Dutch tropical greenhouses in the near future as well, e.g. *Nesticodes rufipes* (Lucas, 1846), *Theotima minutissima* (Petrunkevitch, 1929), *Eukoenia florenxia* (Rucker, 1903) and *Pseudanapis aloha* Forster, 1959 (Snazell & Smithers 2007, Kielhorn 2008, Šestáková et al. 2017, Hula & Pešan 2018).

Comparison of collecting methods

The results of the current study are similar to other studies of arachnofauna in greenhouses involving comparable methods, e.g. Šestáková (2017), where hand collection yielded the highest total number of species and unique species. These results were to be expected, since hand searching covers most strata of the vegetation. Except for the banana baits, each method however yielded unique species (Tab. 1), and they are therefore complementary. The most surprising find, i.e. *Belisana ambengan*, was collected only by beating.

In future arachnid surveys in greenhouses, it is advised to use at least a combination of methods that sample the vegetation (hand searching or beating) and that yield epigeic arachnids (pitfall trapping or litter sieving) (for similar comparisons in the field, see Churchill & Arthur 1999, Jiménez-Valverde & Lobo 2005, Cardoso et al. 2008). Although the number of species did not differ significantly, pitfall trapping is favoured above litter sieving due to its passive and continuous nature. In order to gain more knowledge about arachnid communities in greenhouses, and their potential source function for invasive alien species, it is recommended to carry out more arachnid surveys in this relatively understudied habitat.

Acknowledgements

We are indebted to Ninte Noordijk who found *Coleosoma floridanum* in Rotterdam Zoo in 2018, the second record for the Netherlands at that time, and thereby triggering this survey. We would like to thank Berry van der Hoorn for granting the budget for molecular analyses, and lab personnel Roland Butôt and Frank Stokvis for executing the DNA extraction and performing the molecular analyses at Naturalis Biodiversity Center. Louwerens-Jan Nederlof and his colleagues at Rotterdam Zoo made this study possible and assisted in every way they could. Bruce Schoelitz (HAS University of Applied Sciences, 's Hertogenbosch, the Netherlands) helped in designing the experimental set-up and Peter van Helsdingen (Naturalis Biodiversity Center, Leiden, the Netherlands) is thanked for his aid in the identification laboratory. We thank Henk Siepel (Radboud University Nijmegen, the Netherlands) for examining the mites. Special thanks belong to Dr. Bernhard A. Huber (Zoologisches Forschungsmuseum Alexander Koenig, Bonn, Germany) who identified *Belisana ambengan*. We are thankful to Petr Dolejš, Karl-Heinrich Kielhorn and the editors Theo Blick and Konrad Wiśniewski for their constructive comments and useful suggestions to improve our manuscript.

References

- Bielak-Bielecki P & Rozwałka R 2011 *Nesticella mogera* (Yaginuma, 1972) (Araneae: Nesticidae) in Poland. – *Zeszyty Naukowe Uniwersytetu Szczecińskiego, Acta Biologica* 18: 137-141
- Cardoso P, Scharff N, Gaspar C, Henriques SS, Carvalho R, Castro PH, Schmidt JB, Silva I, Szűts T, De Castro A & Crespo LC 2008 Rapid biodiversity assessment of spiders (Araneae) using semi-quantitative sampling: a case study in a Mediterranean forest. – *Insect Conservation and Diversity* 1: 71-84 – doi: [10.1111/j.1752-4598.2007.00008.x](https://doi.org/10.1111/j.1752-4598.2007.00008.x)
- Churchill TB & Arthur JM 1999 Measuring spider richness: effects of different sampling methods and spatial and temporal scales. – *Journal of Insect Conservation* 3: 287-295 – doi: [10.1023/A:1009638706789](https://doi.org/10.1023/A:1009638706789)
- Gerson U & Weintraub PG 2012 Mites (Acari) as a factor in greenhouse management. – *Annual Review of Entomology* 57: 229-247 – doi: [10.1146/annurev-ento-120710-100639](https://doi.org/10.1146/annurev-ento-120710-100639)
- Gotelli NJ & Colwell RK 2011 Estimating species richness. In: Magurran AE & McGill BJ (eds) *Biological Diversity: Frontiers in Measurement and Assessment*. Oxford University Press, Oxford. pp. 39-54
- Hammen L van der 1949 On Arachnida collected in Dutch greenhouses. – *Tijdschrift voor Entomologie* 91: 72-82
- Hammen L van der 1969a Aanvullende gegevens over de kassenfauna van Nederland. Bibliografische en arachnologische notities [Additional data on the fauna of Dutch greenhouses]. – *Zoologische Bijdragen* 11: 25-28 [in Dutch, with English summary]
- Hammen L van der 1969b Bijdrage tot de kennis van de Nederlandse bastaardschorpioenen (Arachnida, Pseudoscorpionida) [Contribution to the knowledge about Dutch pseudoscorpions (Arachnida, Pseudoscorpionida)]. – *Zoologische Bijdragen* 11: 15-24 [in Dutch, with English summary]
- Hebert PDN, Ratnasingham S & Dewaard JR 2003 Barcoding animal life: cytochrome c oxidase subunit 1 divergences among closely related species. – *Proceedings of the Royal Society of London B (Suppl.)* 270: 96-99 – doi: [10.1098/rsbl.2003.0025](https://doi.org/10.1098/rsbl.2003.0025)
- Helsdingen PJ van 1995 Een stukje tropen in Nederland [A bit of the tropics in the Netherlands]. – *Nieuwsbrief SPINED* 9: 4-6 [in Dutch, with English summary]
- Helsdingen PJ van 2009 *Mermessus denticulatus* (Banks, 1898) and *Mermessus trilobatus* (Emerton, 1882), adventive species in the Netherlands (Araneae, Linyphiidae). – *Contributions to Natural History* 12: 617-626
- Helsdingen PJ van 2018 Catalogus van de Nederlandse spinnen [Catalogue of spiders of the Netherlands]. – Versie 2018.2. European Invertebrate Survey, Leiden. 241 pp. [in Dutch, with English summary]
- Holec M, Svobodová Z, Habušová O, Mohamed Hussein H & Sehnal F 2012 First record of spider *Mermessus trilobatus* (Araneae) in South Bohemia. – *Studia Oecologica* 6: 15-18
- Huber BA 2005 High species diversity, male-female coevolution, and metaphyly in Southeast Asian pholcid spiders: the case of *Belisana Thorell* 1898 (Araneae, Pholcidae). – *Zoologica* 155: 1-126
- Hula V & Pešan V 2018 Spiders of tropical pavilions of the zoological gardens in the Czech Republic. In: Mezöfi L & Szita É (eds) *Program and Abstracts of the 31st European Congress of Arachnology, 8-13 July, 2018, Vác, Hungary*. Hungarian Ecological Society & Centre for Agricultural Research of the Hungarian Academy of Sciences, Budapest. pp. 65
- Ivanova NV, Borisenko AV & Hebert PDN 2009 Express barcodes: racing from specimen to identification. – *Molecular Ecology Resources* 9 (Suppl. 1): 35-41 – doi: [10.1111/j.1755-0998.2009.02630.x](https://doi.org/10.1111/j.1755-0998.2009.02630.x)
- Jiménez-Valverde A & Lobo J M 2005 Determining a combined sampling procedure for a reliable estimation of Araneidae and Thomisidae assemblages (Arachnida, Araneae). – *Journal of Arachnology* 33: 33-42 – doi: [10.1636/M03-10](https://doi.org/10.1636/M03-10)
- Kaňuchová A, Christophoryová J & Krajčovičová K 2016 Pseudoscorpions (Arachnida) collected from the heaps with decomposing material in Slovakia. – *Fragmenta Faunistica* 58 (2015): 111-122 – doi: [10.3161/00159301FF2015.58.2.111](https://doi.org/10.3161/00159301FF2015.58.2.111)
- Kielhorn K-H 2008 A glimpse of the tropics – Spiders (Araneae) in the greenhouses of the Botanic Garden Berlin-Dahlem. – *Arachnologische Mitteilungen* 36: 26-34 – doi: [10.5431/aramit3605](https://doi.org/10.5431/aramit3605)
- Kielhorn K-H 2009 First records of *Spermophora kerinci*, *Nesticella mogera* and *Pseudanapis aloha* on the European Mainland (Araneae: Pholcidae, Nesticidae, Anapidae). – *Arachnologische Mitteilungen* 37: 31-34 – doi: [10.5431/aramit3706](https://doi.org/10.5431/aramit3706)
- Kielhorn K-H 2016 Beitrag zur Kenntnis der Webspinnen und Weberknechte in Berlin und Brandenburg. – *Märkische Entomologische Nachrichten* 17: 261-286
- Koponen S, Fritzén NR & Pajunen T 2016 Checklist of spiders in Finland (Araneae). 6th version, December 2016. – Internet: http://biolcoll.utu.fi/arach/checklist_of_spiders_in_Finland.htm (11. Dec. 2019)
- Korenko S, Rezáč M & Pekár S 2007 Spiders (Araneae) of the family Oonopidae in the Czech Republic. – *Arachnologische Mitteilungen* 34: 6-8 – doi: [10.5431/aramit3402](https://doi.org/10.5431/aramit3402)
- Korenko S, Šmerda J & Pekár S 2009 Life-history of the parthenogenetic oonopid spider, *Triaeris stenaspis* (Araneae: Oonopidae).

- European Journal of Entomology 106: 217–223 – doi: [10.14411/eje.2009.028](https://doi.org/10.14411/eje.2009.028)
- Legg G & Farr-Cox F 2016 Illustrated key to the British false scorpions (Pseudoscorpions). Field Studies Council Publications, Telford. 12 pp.
- Lissner J & Scharff N 2021 Danish Spiders. – Internet: <http://danmarks-edderkopper.dk> (2. Feb. 2021)
- Malten A, Bönsel D & Zizka G 2005 Erfassung von Flora, Fauna und Vegetation auf dem Flughafen Frankfurt am Main. Senckenberg-Museum, Frankfurt. 116 pp.
- Muchmore WB 2000 The Pseudoscorpionida of Hawaii, Part I. Introduction and Chthonioidea. – Proceedings of the Hawaiian Entomological Society 34: 127–142
- Nentwig W, Blick T, Bosmans R, Gloor D, Hänggi A & Kropf C 2019 araneae – Spiders of Europe, version 01.2019. – Internet: <http://www.araneae.unibe.ch> (23. Jan. 2019) – doi: [10.24436/1](https://doi.org/10.24436/1)
- Noordijk J 2020 Een nieuwe trilspinn in Nederland: *Spermophora kerinci* (Araneae: Pholcidae) [A new pholcid spider in the Netherlands: *Spermophora kerinci* (Araneae: Pholcidae)]. – Entomologische Berichten 80: 106 [in Dutch, with English summary]
- Noordijk J, Bink J & Morssinkhof R 2018 De kaskaardespinn *Uloborus plumipes* heeft zich wijd verspreid gevestigd in Nederland (Araneae: Uloboridae) [*Uloborus plumipes* has established widespread in the Netherlands (Araneae: Uloboridae)]. – Nederlandse Faunistische Mededelingen 50: 5–12 [in Dutch, with English summary]
- Oxford G 2011 Death of an urban myth – parthenogenesis in *Uloborus plumipes*. – Newsletter of the British Arachnological Society 121: 6–8
- Pantini P, Bonelli D & Bonacci T 2020 I ragni epigei (Arachnida, Araneae) di un ambiente litoraneo della Calabria [Epigeic spiders (Arachnida, Araneae) in a littoral environment in Calabria]. – Rivista del Museo Civico di Scienze Naturali “Enrico Caffi”, Bergamo 32: 25–32 [in Italian, with English summary]
- Pekár S, Král J, Malten A & Komposch C 2005 Comparison of natural histories and karyotypes of two closely related ant-eating spiders, *Zodarion hamatum* and *Z. italicum* (Araneae, Zodariidae). – Journal of Natural History 39: 1583–1596 – doi: [10.1080/00222930400016762](https://doi.org/10.1080/00222930400016762)
- Petrova V, Salmane I & Ćudare Z 2004 The predatory mite (Acari, Parasitiformes: Mesostigmata (Gamasina); Acariformes: Prostigmata) community in strawberry agrocenosis. – Acta Universitatis Latviensis 676: 87–95
- Pfeiffer WJ 1996 Litter invertebrates. In: Reagan DP & Waide RB (eds) The food web of a tropical rain forest. University of Chicago Press, Chicago. pp. 138–181 – doi: [10.1017/S0266467400010816](https://doi.org/10.1017/S0266467400010816)
- Pfliegler WP 2014 Records of some rare and interesting spider (Araneae) species from anthropogenic habitats in Debrecen, Hungary. – e-Acta Naturalia Pannonica 7: 143–156
- Platnick NI & Dupérré N 2009 The goblin spider genus *Heteroonops* (Araneae, Oonopidae), with notes on *Oonops*. – American Museum Novitates 3672: 1–72 – doi: [10.1206/690.1](https://doi.org/10.1206/690.1)
- Reiser N 2013 Einschleppung und Einwanderung von Spinnentieren (Araneae; Opiliones) in Deutschland. Bachelor thesis, Hochschule Neubrandenburg. 83 pp.
- Rembold K, Junge A-L, Amiet F, Balzari CA, Bergamini A, Blaser S, Boch S, Bürki M, Eggenberg S, Eicher C, Ensslin A, Etter L, Friedli C, Gattlen A, Germann C, Gygax A, Hänggi A, Hertwig ST, Hirschheydt G von, Hoess R, Wisler Hofer C, Inäbnit T, Keller C, Kneubühler J, Kuchler H, Möhl A, Moser T, Neubert E, Pfarrer B, Schäfer D, Schnyder N, Spasojević T, Stofer S, Senn-Irlet B, Es R van der & Fischer M 2020 Vielfalt bedingt Vielfalt – wildlebende Arten im Botanischen Garten der Universität Bern. – Mitteilungen der Naturforschenden Gesellschaft in Bern 77: 24–68 – doi: [10.5169/seals-869443](https://doi.org/10.5169/seals-869443)
- Roberts MJ 1998 Tirion spinnengids [Tirion spider Guide]. Tirion, Barneveld. 397 pp. [in Dutch]
- Rozwałka R, Rutkowski T & Bielak-Bielecki P 2016 New data on introduced and rare synanthropic spider species (Arachnida: Araneae) in Poland (II). – Annales Universitatis Mariae Curie-Skłodowska, Lublin-Polonia 71: 59–85 – doi: [10.17951/c.2016.71.1.59](https://doi.org/10.17951/c.2016.71.1.59)
- Růžička V 2016 Zavlečení [Imports]. – Pavouk 40: 4–6 [in Czech]
- Saaristo MI 2001 Dwarf hunting spiders or Oonopidae (Arachnida, Araneae) of the Seychelles. – Insect Systematics & Evolution 32: 307–358 – doi: [10.1163/187631201X00236](https://doi.org/10.1163/187631201X00236)
- Šestáková A, Suvák M, Krajčovičová K, Kaňuchová A & Christophoryová J 2017 Arachnids from the greenhouses of the Botanical Garden of the P.J. Šafárik University in Košice, Slovakia (Arachnida: Araneae, Opiliones, Palpigradi, Pseudoscorpiones). – Arachnologische Mitteilungen 53: 19–28 – doi: [10.5431/aramit5304](https://doi.org/10.5431/aramit5304)
- Snazell R & Smithers P 2007 *Pseudanapis aloba* Forster (Araneae, Anapidae) from the Eden Project in Cornwall, England. – Bulletin of British Arachnological Society 14: 74–76 – doi: [10.13156/arac.2007.14.2.74](https://doi.org/10.13156/arac.2007.14.2.74)
- Štáhlavský F 2001 Štírci (Arachnida: Pseudoscorpiones) Prahy [Pseudoscorpions (Arachnida: Pseudoscorpiones) of Prague]. – Klapalekiana 37: 73–121 [in Czech, with English abstract and summary]
- Suvák M 2013 Invasive spider *Uloborus plumipes* Lucas, 1846 (Araneae: Uloboridae), new to Slovakia. – Folia Faunistica Slovaca 18: 39–45
- Telfer MG 2020 *Triaeris stenaspis* Simon, 1892 (Oonopidae) at Whip-snade Butterfly House. – Newsletter of the British Arachnological Society 147: 8–9
- Wang C, Zhang X, Pan X, Li Z & Zhu S 2015 Greenhouses: hot-spots in the invasive network for alien species. – Biodiversity and Conservation 24: 1825–1829 – doi: [10.1007/s10531-015-0876-x](https://doi.org/10.1007/s10531-015-0876-x)