



IDENTIFICATION OF A NOVEL MANNHEIMIA GRANULOMATIS LINEAGE FROM LESIONS IN ROE DEER (CAPREOLUS CAPREOLUS)

Authors: Bojesen, Anders M., Larsen, Jesper, Pedersen, Anders G., Mörner, Torsten, Mattson, Roland, et al.

Source: Journal of Wildlife Diseases, 43(3) : 345-352

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-43.3.345>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

IDENTIFICATION OF A NOVEL *MANNHEIMIA GRANULOMATIS* LINEAGE FROM LESIONS IN ROE DEER (*CAPREOLUS CAPREOLUS*)

Anders M. Bojesen,^{1,4,5,6} Jesper Larsen,^{1,4} Anders G. Pedersen,² Torsten Mörner,³ Roland Mattson,³ and Magne Bisgaard¹

¹ Department of Veterinary Pathobiology, The Royal Veterinary and Agricultural University, Stigbøjlen 4, DK-1870 Frederiksberg C, Denmark

² Center for Biological Sequence Analysis, BioCentrum-DTU, Technical University of Denmark, Building 208, DK-2800 Lyngby, Denmark

³ Department of Wildlife, The National Veterinary Institute, SE-751 89 Uppsala, Sweden

⁴ These authors contributed equally to the present work.

⁵ Current address: Department of Veterinary Pathobiology, Faculty of Life Sciences, University of Copenhagen,

4 Stigbøjlen, DK-1870, Frederiksberg C, Denmark

⁶ Corresponding author (email: miki@life.ku.dk)

ABSTRACT: Eight atypical *Mannheimia* isolates were isolated from lesions in roe deer (*Capreolus capreolus*). Traditional classification based on morphologic and physiologic traits showed that they belong to a distinct biogroup (taxon) within genus *Mannheimia*. Extensive phenotypic characterization suggested that the isolates should be classified as *M. granulomatis*, although the presence of distinct traits justified their classification into a separate biogroup within this species. Phylogenetic analyses based on 16S rRNA sequences from two roe deer isolates and 41 other *Mannheimia* strains supported that the roe deer isolates form a monophyletic group within *M. granulomatis*. The *lktA* genotype was present in all roe deer isolates based on Southern blot analysis, whereas the corresponding β -hemolytic phenotype was absent in one of these isolates.

Key words: β -hemolytic phenotype, *Capreolus capreolus*, leukotoxin, *Mannheimia granulomatis*, roe deer.

INTRODUCTION

The genus *Mannheimia* includes strains previously classified as trehalose-negative [*Pasteurella*] *haemolytica*, and is one of the most well-defined and robust clusters within the family of *Pasteurellaceae* Pohl 1981 (Angen et al., 1999). Members of the genus are gram-negative, nonmotile rods or cocci with a G+C content ranging between 39% and 44%. Ribotyping, multi-locus enzyme electrophoresis (MLEE), DNA-DNA hybridizations, and 16S rRNA sequencing have shown that the strains can be divided into five subclades, representing at least five species: *M. haemolytica*+*M. glucosida*, *M. ruminalis*, *M. granulomatis*, *M. varigena*, and subclade V comprising unclassified strains (Angen et al., 1997c; Davies et al., 1997; Angen et al., 1999; Larsen et al., in press). Strains belonging to these species inhabit the mucous membranes of mammals and have been isolated from the respiratory tract of ruminants, rabbits, and hares (Devriese et al., 1991; Angen et al., 1999), the rumen of

sheep and cattle, the ileum of pigs (Angen et al., 1999), and the skin of cattle (Riet-Correa et al., 2000).

Based on phenotypic traits, ruminant strains classified as trehalose-negative *M. [P.] haemolytica* could be divided into 12 biogroups or taxa (Bisgaard and Muters, 1986; Bisgaard et al., 1986). Interestingly, the 16S rRNA phylogeny indicates that different biogroups are present among two or more species, suggesting that members of the same biogroup are not necessarily closely related (Angen et al., 1999; Larsen et al., 2007). Previous works have shown that the β -hemolytic phenotype correlates with the pathogenic potential of strains belonging to genus *Mannheimia*. The results of Angen et al. (1999) indicated that most strains isolated from lesions were β -hemolytic on bovine blood agar plates (except strains belonging to *M. granulomatis*), whereas most strains belonging to the nonpathogenic species *M. ruminalis* were nonhemolytic. Larsen et al. (2007) used Southern blot analysis, sheep blood agar plates, and a representa-

TABLE 1. *Mannheimia* strains isolated from roe deer (*Capreolus capreolus*).

Strains	Lesions	Geographic origin	Year	Sex	Age (yr)	<i>lktA</i>	
						Genotype	Phenotype ^a
B 153/80	Glossitis	Åseda, Sweden	1980	Male	<1	+	+
P 4737/80 ^b	Purulent necrotizing keratoconjunctivitis	Frövi, Sweden	1980	Female	<1	+	+
B 3768/80	Purulent stomatitis and glossitis	Edsbro, Sweden	1980		<1	+	+
B 1829/84	Purulent pneumonia and pleuritis	Stockholm, Sweden	1984	Female	<1	+	-
B 926/85	Purulent pneumonia and pleuritis	Riddarhyttan, Sweden	1985	Male	<1	+	+
Bd 2017/92	Neck phlegmon/abscess	Uppsala, Sweden	1992	Female	<1	+	+
B 234/94	Acute purulent pneumonia	Stockholm, Sweden	1994	Female	~7	+	+
B 1157/95	Acute purulent pneumonia	Stockholm, Sweden	1995	Female	<1	+	+

^a Based on β -hemolysis on sheep blood agar.

^b This strain was not available for analyses of the *lktA* genotype.

tive strain collection to show that the *lktA* genotype and the corresponding β -hemolytic phenotype were present in all *Mannheimia* species, including all strains belonging to *M. haemolytica* and *M. glucosida*, *M. granulomatis*, and *M. varigena*, but only in a fraction of strains belonging to *M. ruminalis*. Murphy et al. (1995) showed that the β -hemolytic phenotype of *M. haemolytica* correlates with the presence of an active leukotoxin (*lkt*) operon on the chromosome. The structural leukotoxin (LkTA) protein of *M. haemolytica*, encoded by *lktA* (Lo et al., 1987), belongs to the *E. coli* HlyA-like subfamily of cytotoxic RTX (repeats in toxin) proteins and is essential in both evasion and exploitation of the adaptive immune system during pulmonary infection (Petras et al., 1995; Tatum et al., 1998; Highlander et al., 2000). These results support the predicted correlation between pathogenic potential and the β -hemolytic phenotype.

In this study, we characterize eight atypical *Mannheimia* isolates obtained from lesions in roe deer (*Capreolus capreolus*) by using a polyphasic strategy that combines phenotypic and genotypic methods. The implications of our results for proper

identification and classification of *Mannheimia* strains are discussed.

MATERIALS AND METHODS

Bacterial strains used

The strains used in this study and their geographic origin, time of isolation, and corresponding pathological lesions are given in Table 1. A total of eight atypical *Mannheimia* strains were isolated from lesions in roe deer submitted to the Department of Wildlife, The National Veterinary Institute, Uppsala, Sweden for post-mortem examination.

Phenotypical characterization

All isolates were phenotypically characterized as previously described by Bisgaard et al. (1991). A total of 82 different phenotypic tests have been used. To determine presence/inactivation of the *lkt* operons, we screened the strains for the β -hemolytic phenotype on sheep blood agar plates as previously described (Murphy et al., 1995).

Reconstruction of phylogenetic trees

The relationships of the strains in this study were inferred with 16S rRNA; these sequences have been used successfully for systematic studies in this group (Angen et al., 1999; Larsen et al., 2007). Sequences for 41 strains were obtained from GenBank (Table 2). The primers previously described by Angen et al.

(1999) were used to amplify and directly sequence 16S rRNA from two roe deer isolates (B 234/94 and B 1829/84). The 16S rRNA sequences were aligned using both Dialign 2 (Morgenstern et al., 1999) and the NAST server (DeSantis et al., 2006), which uses a prealigned set of 16S rRNA sequences as the basis for aligning the user's sequences. The two methods yielded identical alignments. This was expected, because the average pairwise sequence similarity in the full set of 43 sequences was 97.5% and even the least similar pair of sequences was 92.5% identical, supporting that alignment errors were unlikely. Phylogenetic trees were reconstructed with the use of Bayesian techniques as implemented in the program MrBayes version 3.1.1 (Huelsenbeck and Ronquist, 2001; Ronquist and Huelsenbeck, 2003). The best-fitting model was GTR+ invgamma based on the Akaike Information Criterion (AIC) computed by the program MrModeltest (version 2.2, Department of Systematic Zoology, Uppsala University, Uppsala, Sweden; <http://www.ebc.uu.se/systzoo/staff/nylander.html>). Markov chain Monte Carlo (MCMC) was run for 10,000,000 generations with four chains while sampling once every 100 generations. Convergence was confirmed by comparing the results of two independent runs. The program Tracer (version 1.3; Department of Zoology, Oxford University, Oxford, UK; <http://evolve.zoo.ox.ac.uk/software.html?id=tracer>) was used to determine burn-in and also for further confirmation of proper mixing and adequate run length. A burn-in of 1,000,000 generations (10,000 samples) was used in both cases. The distribution of 16S rRNA trees resulting from MCMC was summarized in the form of a consensus tree with all compatible bipartitions included by using the authors' own software in a manner that is essentially identical to what is obtained when using MrBayes' sumt command with the setting `contype=allcompat`, except that branch lengths were averaged over all trees; setting the branch length to zero for those trees that did not contain the corresponding bipartition, as suggested by Felsenstein (2003). The tree was rooted based on a maximum-likelihood analysis reported by Angen et al. (1999).

Analyses of the *lktA* genotype

We analyzed the distribution of the *lktA* gene among roe deer isolates by using Southern blot as previously described by Larsen et al. (2007). The primers `manpop_UP` and `manpop_DOWN` previously described by Larsen et al. (2007) were used to amplify the +845/+1,302

region of *lktA*. The reaction conditions were 2.5 U *Taq* polymerase, 16 mM $(\text{NH}_4)_2\text{SO}_4$, 67 mM Tris-HCl, 0.01% Tween-20, 2.5 mM Mg_2SO_4 , each primer at 0.5 mM, and each nucleotide at 0.1 mM. The cycling conditions were initial denaturation at 94 C followed by 25 cycles of 94 C for 30 s, 52 C for 30 sec, and 72 C for 30 sec, finishing with extension at 72 C for 10 min.

Sequence data

Sequences have been deposited in the GenBank database (accession numbers (AY425296 and AY425297).

RESULTS

Bacterial strains used

The strains used in this study and their geographic origin, time of isolation, and corresponding lesions are given in Table 1. It should be noted that the strains were isolated from six different locations within a 300-km radius from Norrköping (58°36'N, 16°11'E) in southeastern Sweden. The isolates were obtained during a period from 1980 to 1995. Six of the isolates were obtained from animals <1 yr old, whereas a single isolate originated from a 7-yr-old female. Lesions varied from acute purulent bronchopneumonia ($n=2$), purulent pneumonia and pleuritis ($n=2$), glossitis ($n=1$), purulent stomatitis and glossitis ($n=1$), and phlegmon and abscess formation ($n=1$).

Phenotypic characterization

All isolates were gram-negative, non-motile, catalase and oxidase positive, and decomposed glucose fermentatively in Hugh and Leifson medium. All isolates were porphyrin, alanine aminopeptidase and phosphatase positive, and reduced nitrate. In addition, acid was produced from glycerol, D-ribose, D-xylose, meso-inositol, D-mannitol, D-fructose, D-galactose, D-glucose without gas formation, lactose, sucrose, and raffinose. Positive reactions were also obtained for all isolates in ONPG (β -galactosidase) and PGUA (β -glucuronidase). The β -hemolytic phenotype on bovine and ovine blood agar plates

TABLE 2. Isolates of *Mannheimia*, host species, origin, GenBank accession numbers for 16S rRNA sequences used in this study.

Taxon ^a	Strain ID	Host Species	Host Family	Country	GenBank accession no.
<i>M. haemolytica</i>					
Biogroup 1	PHL213	<i>Bos taurus</i>	Ruminantia	UK	DQ301920
Biogroup 1	CCUG 12392 ^T	<i>Ovis aries</i>	Ruminantia		AF060699
<i>M. glucosida</i>					
Biogroup 3A	P731	<i>O. aries</i>	Ruminantia	USA	AF053888
Biogroup 3B	P925 ^T	<i>O. aries</i>	Ruminantia	Scotland	AF053889
Biogroup 3C	UT18	<i>O. aries</i>	Ruminantia	Scotland	AF053890
Biogroup 3D	H62	<i>O. aries</i>	Ruminantia	Belgium	DQ301921
Biogroup 3E	P741	<i>O. aries</i>	Ruminantia	USA	DQ301922
Biogroup 3F	P933	<i>O. aries</i>	Ruminantia	Scotland	DQ301923
Biogroup 3G	P737	<i>O. aries</i>	Ruminantia	USA	AF053891
Biogroup 3H	P733	<i>O. aries</i>	Ruminantia	USA	AF053892
Biogroup 9	P730	<i>O. aries</i>	Ruminantia	USA	AF053897
<i>M. granulomatis</i>					
Biogroup 3J	W4672/1	<i>B. taurus</i>	Ruminantia	Australia	DQ301924
Bt 20 biovar 1	Ph13	<i>Lepus capensis</i>	Leporidae	France	AF053901
Bt 20 biovar 2	BJ1680.3	<i>L. capensis</i>	Leporidae	Belgium	DQ301925
[<i>P.</i>] <i>granulomatis</i>	P1135/26 ^T	<i>B. taurus</i>	Ruminantia	Brazil	AF053902
<i>M. varigena</i>					
Biogroup 6	177 ^T	<i>B. taurus</i>	Ruminantia	Germany	AF053893
Biogroup 6	V1835	<i>B. taurus</i>	Ruminantia	Australia	AY425282
Bt 15 biovar 1 <i>orn</i> ⁺	P655	<i>Sus scrofa</i>	Suidae	Denmark	AF053899
Bt 15 biovar 2 <i>orn</i> ⁻	3997/82	<i>S. scrofa</i>	Suidae	Denmark	DQ301926
Bt 36	H39	<i>B. taurus</i>	Ruminantia	Belgium	DQ301927
<i>M. ruminalis</i>					
Biogroup 1	UT26	<i>O. aries</i>	Ruminantia	Scotland	AF053887
Biogroup 8D	HPA98	<i>O. aries</i>	Ruminantia	Scotland	AF053896
Biogroup 10	HPA95	<i>O. aries</i>	Ruminantia	Scotland	AY425289
Biogroup 10	HPA114	<i>O. aries</i>	Ruminantia	Scotland	AY425290
Biogroup 10	UT27	<i>O. aries</i>	Ruminantia	Scotland	AY425291
Bt 18 biovar 1	HPA92 ^T	<i>O. aries</i>	Ruminantia	Scotland	AF053900
Bt 18 biovar 1	HPA81	<i>O. aries</i>	Ruminantia	UK	U57077
Bt 18 biovar 2	HPA113	<i>O. aries</i>	Ruminantia	UK	AY425283
Bt 18 biovar 2	HPA90	<i>O. aries</i>	Ruminantia		AY425284
Bt 18 biovar 2	UT38	<i>O. aries</i>	Ruminantia		AY425285
Bt 18 biovar 3 <i>xyl</i> ⁺	HPA109	<i>O. aries</i>	Ruminantia	UK	AY425286
Bt 18 biovar 3 <i>xyl</i> ⁻	HPA93	<i>O. aries</i>	Ruminantia	UK	AY425287
Bt 18 biovar 4	HPA88	<i>O. aries</i>	Ruminantia	UK	AY425288
Subclade V					
Biogroup 7	R19.2	<i>B. taurus</i>	Ruminantia	Scotland	AF053894
Biogroup 8A ^c	HPA102	<i>O. aries</i>	Ruminantia	Scotland	AF053895
Biogroup 8B	274	<i>B. taurus</i>	Ruminantia	Germany	AY425292
Biogroup 8B	R108B(3)	<i>B. taurus</i>	Ruminantia	Scotland	AY425293
Biogroup 8C	M14.4	<i>B. taurus</i>	Ruminantia	Scotland	AY425294
Biogroup 10	BJ3956.1	<i>O. aries</i>	Ruminantia	Belgium	AY425295
Biogroup 10	HPA121	<i>B. taurus</i>	Ruminantia	Scotland	AF053898
Bt 39	BNO311	<i>B. taurus</i>	Ruminantia	Australia	AF216870

^a Bt=Bisgaard taxon.

TABLE 3. Phenotypic characters separating roe deer isolates from related taxa of genus *Mannheimia*.^a

Character	Roe deer isolates	<i>M. granulomatis</i>			str. P1135/26 ^e	<i>M. ruminalis</i> str. HPA92 ^e
		Bt 20 biovar 1 ^b	Bt 20 biovar 2 ^c	Biogroup 3J ^d		
β-hemolysis ^f	+/w	+	+	+	+	—
Glycerol	(+)	(+)	(+)	(+)	(+)	—
D (+) xylose	+/(+)	+	+/(+)	+	—	—
Meso-inositol	(+)	(+)/—	(+)/w	—	—	—
D (—) sorbitol	—	+	+	+	+	—
Cellobiose	—	+/(+)	—	(+)	(+)	—
Maltose	—	+/(+)	(+)	(+)	(+)	—
Raffinose	(+)	+/w	(+)	—	—	—
Dextrin	—	(+)/—	(+)/w	(+)	w	—
Aesculin	—	+/(+)	—	+/(+)	+	—
Amygdalin	—	(+)	—	(+)	+	—
Arbutin	—	+/(+)	—	(+)	+	—
Gentiobiose	—	+/(+)	—	(+)	+	—
Salicin	—	+/(+)	—	(+)	+	—
NPG (β-glucosidase)	—	+	+	+	+	—
PGUA (β-glucuronidase)	+	—	—	—	+	—

^a +, ≥90% strains are positive; —, <10% strains are positive; w, <90 and ≥10 positive; (+), late positive (3–14 days).

^b Based on 19 isolates from *Lepus capensis* and *Oryctolagus cuniculus*.

^c Based on seven isolates from *Lepus capensis*.

^d Based on three isolates from *Bos taurus*.

^e T=Type strain.

^f Based on bovine blood.

was present in six isolates, whereas isolate B 1828/84 was nonhemolytic. Late acid production from L-fucose was observed for two isolates, whereas a single isolate showed a positive reaction in the ONPX test (β-xylosidase).

A negative reaction was observed for all isolates in the following tests: symbiotic growth, Simmons citrate, malonate, H₂S/TSI, KCN growth, MR 37 C, VP 37 C, formation of gas from nitrate, urease, arginine dihydrolase, lysine and ornithine decarboxylase, phenylalanine deaminase, indole, gelatinase, hydrolysis of Tween 20 and Tween 80, McConkey agar growth, pigment formation, and production of acid from meso-erythritol, adonitol, D-arabitol, xylitol, L-arabinose, D-arabinose, L-xylose, dulcitol, D-sorbitol, D-fucose, D-mannose, L-rhamnose, L-sorbose, cellobiose, maltose, D-melibiose, trehalose, D-melezitose, dextrin, D-glycogen, inulin, aesculin,

amygdalin, arbutin, gentiobiose, salicin, D-turanose, and β-N-CH₃-glucosamid. Negative reactions were also observed in NPG (β-glucosidase), PNPG (α-glucosidase), ONPF (α-fucosidase), α-galactosidase, and α-mannosidase. Phenotypic characters separating the isolates from biogroups belonging to *M. granulomatis* and the type strain of *M. ruminalis* (HPA92^T) are given in Table 3.

16S rRNA phylogeny

BLAST searches of the NCBI non-redundant (nr) database with the 16S rRNA sequences of isolates B 234/94 and B 1829/84 as queries suggested that they are most closely related to *M. granulomatis*. Because the closest BLAST hits are not necessarily each other's closest relative (Koski and Golding, 2001), we reconstructed the phylogeny with the use of these sequences, along with sequences of




FIGURE 1. Distribution of 16S rRNA trees from Markov chain Monte Carlo (MCMC) summarized in the form of a consensus tree with all compatible bipartitions included and with branch lengths averaged over all trees, setting the branch length to zero for those trees that did not contain the corresponding bipartition. The tree was rooted based on a maximum-likelihood analysis reported by Angen et al. (1999). Posterior probability values are shown for all compatible bipartitions. Presence/absence of the *lktA* genotype and the corresponding β -hemolytic phenotype is indicated by $\{\pm;\pm\}$. Monophyly of the five subclades reported by Angen et al. (1999) is indicated by black circles. Nomenclature: Sequence names contain the corresponding strain ID as listed in Table 2.

41 *Mannheimia* strains. The 16S rRNA data set included 1,257 bp after removal of ambiguous bases. Both runs converged and the consensus tree with all compatible bipartitions included and with branch lengths averaged over all trees, setting the branch length to zero for those trees that did not contain the corresponding

bipartition, is shown in Figure 1, along with posterior probabilities (PP). This phylogram is largely topologically concordant with the Bayesian analysis reported by Larsen et al. (2007) and suggest that the two roe deer isolates form a cluster within *M. granulomatis* (PP=0.90%). There was strong PP support for mono-

phyly of the roe deer isolates (PP=100%), supporting that they belong to a distinct lineage of *M. granulomatis* (PP=100%).

Analyses of the *lktA* genotype

The *lktA* genotype and the corresponding β -hemolytic phenotype are given in Table 1. It should be noted that isolate P 4737/80 died during storage and could not be included in these analyses. Southern blot analysis indicated that the *lktA* gene was present in all seven isolates. The +845/+1,302 region of the *lktA* gene was amplified by PCR in the six β -hemolytic isolates. We could not amplify *lktA* from the nonhemolytic isolate B 1829/84, presumably because substitutions have resulted in imperfect matches of the primers.

DISCUSSION

Traditional classification based on morphologic and physiologic traits showed that the eight roe deer isolates belonged to a distinct biogroup (taxon) within genus *Mannheimia* (data not shown). Extensive phenotypic characterization suggested that the isolates should be classified as *M. granulomatis*, although the presence of distinct traits justified their classification into a separate biogroup within this species (Table 3). The 16S rRNA tree supported that the roe deer isolates form a monophyletic group within *M. granulomatis*. The 16S rRNA sequences from the roe deer isolates were almost identical despite having been isolated in 1984 and 1994, respectively (data not shown), supporting that they constitute a clonal complex within *M. granulomatis*.

The bovine *M. granulomatis* strains W4672/1 (biogroup 3J) and P1135/26^T (*M. [P.] granulomatis*) were their closest neighbors on the 16S rRNA tree, whereas the leporine strains Ph13 (Bisgaard taxon 20 biovar 1) and BJ1680.3 (Bisgaard taxon 20 biovar 2) constituted a more distantly related group (Fig. 1). The results presented here suggest the divergence of at least three distinct *M. granulomatis*

lineages that may have adapted to cervine, bovine, and leporine hosts, respectively.

Prior to this work, nonhemolytic *Mannheimia* strains had never been isolated outside the bovine and ovine rumen and they had never been associated with disease conditions, supporting the predicted correlation between pathogenic potential and the β -hemolytic phenotype. However, our analyses showed that isolate B 1829/84 was nonhemolytic, although the *lktA* gene was present on the chromosome. The presence of an inactivated *lkt* operon has been observed in a single *M. ruminalis* strain (HPA88), whereas chromosomal deletions of $\sim 7,400$ bp, corresponding to the entire *lkt* operon, have occurred in the remaining nonhemolytic *M. ruminalis* strains (Larsen et al., in press). If one assumes that the evolving *M. granulomatis* populations adapted to different ecologic niches (host species and anatomic niches) offering different conditions for growth, then this diversity could imply that different pools of unused genes have been lost in those populations. This hypothesis predicts the existence of variable morphologic and physiologic traits between strains belonging to the same species and constitutes the most parsimonious explanation for the observed phenotypic variation within *M. granulomatis* and other *Mannheimia* species (Angen et al., 1997a, b). These results support the use of a polyphasic strategy, combining phenotypic and genotypic methods, for proper identification and classification of *Mannheimia* strains.

ACKNOWLEDGMENTS

We thank C. Pedersen and C. Buerholt for technical assistance. This work was supported by a grant from the Danish Agricultural and Veterinary Research Council (9702797).

LITERATURE CITED

- ANGEN, Ø., J. E. OLSEN, AND M. BISGAARD. 1997a. Further studies of the relationships among strains classified as taxon 15, taxon 18, taxon 20, [*Pasteurella*] *granulomatis* of the [*Pasteurella*] *haemolytica*-complex in ruminants using

- quantitative evaluation of phenotypic data. *Zentralblatt für Bakteriologie* 286: 317–332.
- , B. AALBAEK, E. FALSEN, J. E. OLSEN, AND M. BISGAARD. 1997b. Relationships among strains classified with the ruminant *Pasteurella haemolytica*-complex using quantitative evaluation of phenotypic data. *Zentralblatt für Bakteriologie* 285: 459–479.
- , D. A. CAUGANT, J. E. OLSEN, AND M. BISGAARD. 1997c. Genotypic relationships among strains classified under the (*Pasteurella*) *haemolytica*-complex as indicated by ribotyping and multilocus enzyme electrophoresis. *Zentralblatt für Bakteriologie* 286: 333–354.
- , R. MUTTERS, D. A. CAUGANT, J. E. OLSEN, AND M. BISGAARD. 1999. Taxonomic relationships of the [*Pasteurella*] *haemolytica* complex as evaluated by DNA-DNA hybridizations and 16S rRNA sequencing with proposal of *Mannheimia haemolytica* gen. nov., comb. nov., *Mannheimia granulomatis* comb. nov., *Mannheimia glucosida* sp. nov., *Mannheimia ruminalis* sp. nov. and *Mannheimia varigena* sp. nov. *International Journal of Systematic Bacteriology* 49 Pt 1: 67–86.
- BISGAARD, M., AND R. MUTTERS. 1986. Re-investigations of selected bovine and ovine strains previously classified as *Pasteurella haemolytica* and description of some new taxa within the *Pasteurella haemolytica*-complex. *Acta Pathologica, Microbiologica et Immunologica Scandinavica B* 94: 185–193.
- , J. E. PHILLIPS, AND W. MANNHEIM. 1986. Characterization and identification of bovine and ovine *Pasteurellaceae* isolated from the oral cavity and rumen of apparently normal cattle and sheep. *Acta Pathologica, Microbiologica et Immunologica Scandinavica B* 94: 9–17.
- , S. B. HOUGHTON, R. MUTTERS, AND A. STENZEL. 1991. Reclassification of German, British and Dutch isolates of so-called *Pasteurella multocida* obtained from pneumonic calf lungs. *Veterinary Microbiology* 26: 115–124.
- DAVIES, R. L., S. ARKINSAW, AND R. K. SELANDER. 1997. Evolutionary genetics of *Pasteurella haemolytica* isolates recovered from cattle and sheep. *Infection and Immunity* 65: 3585–3593.
- DESANTIS, T. Z. JR., P. HUGENHOLTZ, K. KELLER, E. L. BRODIE, N. LARSEN, Y. M. PICENO, R. PHAN, AND G. L. ANDERSEN. 2006. NAST: A multiple sequence alignment server for comparative analysis of 16S rRNA genes. *Nucleic Acids Research* 34: W394–399.
- DEVRIESE, L. A., M. BISGAARD, J. HOMMEZ, E. UYTTEBROEK, R. DUCATILLE, AND F. HAESEBROUCK. 1991. Taxon 20 (Fam *Pasteurellaceae*) infections in European brown hares (*Lepus europaeus*). *Journal of Wildlife Disease* 27: 685–687.
- FELSENSTEIN, J. 2003. *Inferring phylogenies*. Sinauer Associates, Sunderland, Massachusetts, 664 pp.
- HIGHLANDER, S. K., N. D. FEDOROVA, D. M. DUSEK, R. PANCIERA, L. E. ALVAREZ, AND C. RINEHART. 2000. Inactivation of *Pasteurella (Mannheimia) haemolytica* leukotoxin causes partial attenuation of virulence in a calf challenge model. *Infection and Immunity* 68: 3916–3922.
- HUELSENBECK, J. P., AND F. RONQUIST. 2001. MRBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics* 17: 754–755.
- KOSKI, L. B., AND G. B. GOLDING. 2001. The closest BLAST hit is often not the nearest neighbor. *Journal of Molecular Evolution* 54: 540–542.
- LARSEN, J., A. G. PEDERSEN, H. CHRISTENSEN, M. BISGAARD, Ø. ANGEN, P. AHRENS, AND J. E. OLSEN. 2007. Evidence for vertical inheritance and loss of the leukotoxin operon in genus *Mannheimia*. *Journal of Molecular Evolution*. <http://dx.doi.org/10.1007/s00239-006-0065.3>.
- LO, R. Y., C. A. STRATHDEE, AND P. E. SHEWEN. 1987. Nucleotide sequence of the leukotoxin genes of *Pasteurella haemolytica* A1. *Infection and Immunity* 55: 1987–1996.
- MORGENSTERN, B. 1999. DIALIGN 2: Improvement of the segment-to-segment approach to multiple sequence alignment. *Bioinformatics* 15: 211–218.
- MURPHY, G. L., L. C. WHITWORTH, K. D. CLINKENBEARD, AND P. A. CLINKENBEARD. 1995. Haemolytic activity of the *Pasteurella haemolytica* leukotoxin. *Infection and Immunity* 63: 3209–3212.
- PETRAS, S. F., M. CHIDAMBARAM, E. F. ILLYES, S. FROSHAUER, G. M. WEINSTOCK, AND C. P. REESE. 1995. Antigenic and virulence properties of *Pasteurella haemolytica* leukotoxin mutants. *Infection and Immunity* 63: 1033–1039.
- RIET-CORREA, F., S. L. LADEIRA, G. B. ANDRADE, AND G. R. CARTER. 2000. Lechiguana (focal proliferative fibrogranulomatous panniculitis) in cattle. *Veterinary Research Communications* 24: 557–572.
- RONQUIST, F., AND J. P. HUELSENBECK. 2003. MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19: 1572–1574.
- TATUM, F. M., R. E. BRIGGS, S. S. SREEVATSAN, E. S. ZEHR, H. S. LING, L. O. WHITELEY, T. R. AMES, AND S. K. MAHESWARAN. 1998. Construction of an isogenic leukotoxin deletion mutant of *Pasteurella haemolytica* serotype 1: Characterization and virulence. *Microbial Pathogenesis* 24: 37–46.

Received for publication 6 July 2006.